

Correspondence

INSTRUCTIONS FOR LETTERS TO THE EDITOR

Editorial comment in the form of a Letter to the Editor is invited; however, it should not exceed 800 words, with a maximum of 10 references and no more than 2 figures (submitted as camera ready hard copy per Journal Guidelines) or tables and no subdivision for an Abstract, Methods, or Results. Letters should have no more than 3 authors. Full name(s) and address of the author(s) should accompany the letter as well as the telephone number, fax number, or E-mail address.

Contact: The Managing Editor, The Journal of Rheumatology, 365 Bloor Street East, Suite 901, Toronto, ON CANADA M4W 3L4. Tel: 416-967-5155; Fax: 416-967-7556; E-mail: jrheum@jrheum.com Financial associations or other possible conflicts of interest should always be disclosed.

Macrophage Activation Syndrome and Etanercept in Children with Systemic Juvenile Rheumatoid Arthritis

To the Editor:

We read with interest the article by Ramanan, *et al*¹ describing a patient with systemic onset juvenile rheumatoid arthritis (JRA) who developed a clinical and laboratory picture consistent with macrophage activation syndrome (MAS) following etanercept therapy. We describe a patient with a similar presentation who developed MAS while taking etanercept; however, the cause of disease activation was found to be Epstein-Barr viral (EBV) infection.

A 10.5-year-old Caucasian girl presented in December 1995 with an 8-week history of intermittent fever, characteristic rash, polyarthritis, and morning stiffness and was diagnosed with systemic onset JRA. She was initially treated with prednisone, naprosyn, and methotrexate (MTX) with partial response. In June 1999, she started taking etanercept 0.4 mg/kg subcutaneous biweekly to achieve full clinical remission. She tolerated it well with no complication, and JRA manifestations completely resolved within 6 months of therapy. She remained in remission for the next 2 years while taking MTX and etanercept.

In June 2001, she presented to a local hospital with high grade fever of 106°F. Examination revealed an enlarged spleen and normal liver span, with no lymphadenopathy and no evidence of arthritis. Laboratory evaluation showed low hemoglobin concentration (6.8 g/dl), low platelet count (43,000/ μ l), low white blood cell count (2300/ μ l); and high prothrombin time (17.9 s), high international normalized ratio (1.92) and elevated erythrocyte sedimentation rate (31 mm/h). Her liver profile showed elevated liver enzymes including ALT 306 U (normal 24 U), AST 140 U (normal 36 U), and alkaline phosphatase 831 U. A bone marrow biopsy showed erythrophagocytosis with normal iron stores. Her MTX and etanercept were stopped at the time of admission.

Over the next few days, she continued to have moderate to high grade fever with deterioration of her general condition, and she was transferred to the intensive care unit for further care. She was given antibiotics for possible sepsis, although her blood and urine cultures obtained at the time of admission were negative.

Serological tests for viral infections including hepatitis and cytomegalovirus were negative. On Day 4, the IgM antibody titers against viral capsid antigen became positive, indicating a recent EBV infection. A diagnosis of MAS was made on the basis of pancytopenia, elevated liver enzymes, coagulopathy, persistent fever, and bone marrow findings. She was treated with intravenous methylprednisolone and cyclosporine followed by prednisone. Her clinical and laboratory measures showed marked improvement over the next few days. She was discharged home on a tapering dose of oral prednisone. In a followup visit 2 weeks later she was doing well and the MTX and etanercept were restarted.

This case bears many similarities to the case described by Ramanan and colleagues; however, EBV infection and not etanercept was found to be a possible trigger for MAS in our patient. In the absence of a definitive causal relation between etanercept and MAS, we are reluctant to accept that etanercept alone may induce MAS. Although Ramanan and colleagues ruled out EBV infection, other uncommon viral infections might have been a trigger in their patient.

Few studies have actually found etanercept to be useful in the treatment of MAS^{2,3}. Etanercept continues to be invaluable for therapy of JRA as well as many other rheumatic disorders, and while it is important to encourage continuous surveillance and report side effects, it is also important and prudent to avoid generating undue alarm regarding uncertain and not well proven effects.

HAROON SARWAR, MD, Clinical Rheumatology Fellow, Section of Rheumatology, Department of Medicine; LUIS R. ESPINOZA, MD, Professor of Medicine, Chief, Rheumatology Section, Department of Medicine; ABRAHAM GEDALIA, MD, Professor of Pediatrics, Chief, Division of Pediatric Rheumatology, Department of Pediatrics, LSU Health Sciences Center and Children's Hospital of New Orleans, 200 Henry Clay Avenue, New Orleans, Louisiana 70118, USA.

REFERENCES

1. Ramanan AV, Schneider R. Macrophage activation syndrome following initiation of etanercept in a child with systemic onset juvenile rheumatoid arthritis. *J Rheumatol* 2003;30:401-3.
2. Prahala S, Bove KE, Dickens D, Lovell DJ, Gromm AA. Etanercept in the treatment of macrophage activation syndrome. *J Rheumatol* 2001;28:2120-4.
3. Aeberli D, Oertle S, Mauron H, Reichenbach S, Jordi B, Villiger PM. Inhibition of the TNF pathway: use of infliximab and etanercept as remission-inducing agents in cases of therapy-resistant chronic inflammatory disorders. *Swiss Med Wkly* 2002;132:414-22.

Drs. Ramanan and Schneider reply

To the Editor:

We thank Sarwar, *et al* for their interest in our article. The patient they describe developed macrophage activation syndrome (MAS) after being maintained on etanercept for 2 years. A similar result has been reported by others as well. Our patient was different in that she developed MAS within 2 weeks (4 doses) of initiation of etanercept, which she had never received before. Our patient also developed giant urticaria at the injection site that heralded the onset of MAS.

We were not able to identify any of the typical viral triggers that have been associated with MAS. The viral serology was negative for Epstein-Barr virus, cytomegalovirus, adenovirus, and parvovirus B19. As we did not do a bone marrow aspirate, testing for viruses by polymerase chain reaction (PCR) in the marrow could not be done.

We did acknowledge in our report that etanercept has been used in the acute management of MAS. However, the same center that published this case report has now described 2 patients who developed MAS while taking etanercept. It is unclear whether MAS occurred when these patients were

maintained on etanercept or after initiation of therapy, as seen in our patient.

Children with systemic onset JRA seem to be susceptible to developing MAS after initiation of a number of therapies, including second injection of gold, nonsteroidal antiinflammatory drugs, and methotrexate.

We agree with Sarwar and colleagues that etanercept is a valuable agent in the management of JRA. While we concur that it is important not to generate undue alarm about etanercept, we believe that increased vigilance for MAS is required in children who are starting this therapy. Our case report highlights this, and also that giant urticaria at the injection site of etanercept may perhaps herald the onset of MAS.

ATHIMALAIPET V. RAMANAN, MRCP(UK), MRCPC(UK); RAYFEL SCHNEIDER, MBBS, FRCP, Division of Rheumatology, Department of Pediatrics, The Hospital for Sick Children, 555 University Avenue, Toronto, Ontario M5G 1X8, Canada.

REFERENCES

1. Ramanan AV, Schneider R. Macrophage activation syndrome following initiation of etanercept in a child with systemic onset juvenile rheumatoid arthritis. *J Rheumatol* 2003;30:401-3.
2. Aouba A, De Bendt M, Aslangul E, Atkhen N, Patri B. Hemophagocytic syndrome in a rheumatoid arthritis patient treated with infliximab. *Rheumatology Oxford* 2003;42:800-2.
3. Grom AA, Lee S, Villanueva J, Passo M, Filipovich A. Natural killer cell dysfunction with macrophage activation syndrome in systemic onset juvenile rheumatoid arthritis [abstract]. *Arthritis Rheum* 2002;46 Suppl:S472.
4. Prahald S, Bove KE, Dickens D, Lovell DJ, Grom AA. Etanercept in the treatment of macrophage activation syndrome. *J Rheumatol* 2001;28:2120-4.
5. Silverman ED, Miller JJ 3rd, Bernstein B, Shafai T. Consumption coagulopathy associated with systemic juvenile rheumatoid arthritis. *J Pediatr* 1983;103:872-6.
6. Hadchouel M, Prieur AM, Griscelli C. Acute hemorrhagic, hepatic, and neurologic manifestations in juvenile rheumatoid arthritis: possible relationship to drugs or infection. *J Pediatr* 1985;106:561-6.
7. Ravelli A, Caria MC, Buratti S, Malattia C, Temporini F, Martini A. Methotrexate as a possible trigger of macrophage activation syndrome in systemic juvenile idiopathic arthritis. *J Rheumatol* 2001;28:865-7.
8. Stephan JL, Kone-Paut I, Galambrun C, Mouy R, Bader-Meunier B, Prieur AM. Reactive haemophagocytic syndrome in children with inflammatory disorders. A retrospective study of 24 patients. *Rheumatology Oxford* 2001;40:1285-92.
9. Sawhney S, Woo P, Murray KJ. Macrophage activation syndrome: a potentially fatal complication of rheumatic disorders. *Arch Dis Child* 2001;85:421-6.

Therapeutic Implications for Interferon- α in Arthritis

To the Editor:

We read with interest the article by Wong, *et al*¹ on their experiments indicating that interferon- α (IFN- α) might be useful in the treatment of rheumatoid arthritis (RA) through the effects it exerts on cytokines and soluble cytokine receptors by augmenting osteoprotegerin (OPG) gene expression and the production of interleukin 1 receptor antagonist (IL-1Ra) and soluble tumor necrosis factor receptor (sTNFR) in synovial tissue or synovial fluid. IFN- α is a "pleiotropic" cytokine and has been shown to exert immunostimulatory as well as immunosuppressive effects. Reports by Wong, *et al* and others have described the immunosuppressive and anti-inflammatory effects of IFN- α . It decreases the levels of IL-8², whereas

sTNFR-I (p55), sTNFR-II (p75), and IL-1Ra are increased and probably antagonize the action of the respective cytokines, as shown in patients with hepatitis C and in *in vitro* experiments³⁻⁷, and reviewed by Kaser and Tilg⁸. Matrix metalloproteinase 9 (MMP-9) gene expression (MMP-9 being one of the most important mediators of cartilage destruction) has recently been shown to be decreased by IFN- β , also a type I interferon⁹.

Our own observations in patients with Behçet's disease treated with IFN- α , in concordance with the data of Wong, *et al*, also revealed an increase in sTNFR-II and IL-1Ra in serum (in press). However, Wong, *et al* do not discuss that IFN- α also induces HLA class I expression, increases the numbers of B cells, and activates monocytes/macrophages^{10,11}. Further, TNF- α gene expression and serum levels (in press) are increased by IFN- α ¹². Thus it not only improves antigen presentation and elimination of foreign antigens, but may also induce autoimmune phenomena. Autoimmune thyroiditis and SLE have been described to occur as adverse effects of IFN treatment^{13,14}. As for systemic lupus erythematosus (SLE), it has become clear that plasmacytoid dendritic cells, which are the naturally occurring main producers of IFN- α *in vivo*, play a major role in the pathogenesis of SLE¹⁵.

The authors might also have discussed that cases of RA occurred during IFN- α treatment for diseases such as carcinoma, lymphoma, chronic myelogenous leukemia, and hepatitis B and C^{16,17}. Thus, it must be considered that if patients with RA were treated with IFN- α , the proinflammatory actions of this cytokine (mainly the monocyte activation with the resulting increase in TNF- α and B cell activation) might outweigh the anti-inflammatory ones (such as the increase in sTNFR-II and IL-1Ra and inhibition of OPG gene expression), resulting in increased disease activity. It might be possible that IFN- α exerts differential effects depending on the immunological background (e.g., Th1, Th2, or Th0).

The results of the preclinical studies announced by the authors will be of great interest.

INA KÖTTER, MD, Senior Physician, Department of Internal Medicine II, Haematology, Oncology, Immunology and Rheumatology;
NICOLE STÜBIGER, MD, Department of Ophthalmology I, University Hospital, Otfried-Müller-Str. 10, D-72076 Tübingen, Germany.
E-mail: ina.koetter@med.uni-tuebingen.de

REFERENCES

1. Wong T, Majchrzak B, Bogoch E, Keystone EC, Fish EN. Therapeutic implications for interferon- α in arthritis: a pilot study. *J Rheumatol* 2003;30:934-40.
2. Reznikov LL, Puren AJ, Fantuzzi G, et al. Spontaneous and inducible cytokine responses in healthy humans receiving a single dose of IFN-alpha 2b: increased production of interleukin-1 receptor antagonist and suppression of IL-1-induced IL-8. *J Interferon Cytokine Res* 1998;18:897-903.
3. Dinarello CA. Induction of interleukin-1 and interleukin-1 receptor antagonist. *Semin Oncol* 1997;24:S9-81-93.
4. Taylor JL, Grossberg SE. The effects of interferon-alpha on the production and action of other cytokines. *Semin Oncol* 1998;25:23-9.
5. Corssmit EP, Heijligenberg R, Hack CE, Endert E, Sauerwein HP, Romijn JA. Effects of interferon-alpha administration on leucocytes in healthy humans. *Clin Exp Immunol* 1997;107:359-63.
6. Itoh Y, Okanou T, Ohnishi N, et al. Serum levels of soluble tumor necrosis factor receptors and effects of interferon therapy in patients with chronic hepatitis C virus infection. *Am Gastroenterol* 1999;94:1332-40.
7. Tilg H, Vogel W, Dinarello CA. Interferon-alpha induces circulating tumor necrosis factor receptor p55 in humans. *Blood* 1995;85:433-5.
8. Kaser A, Tilg H. Interferon-alpha in inflammation and immunity. *Cell Mol Biol (Noisy-le-grand)* 2001;47:609-17.
9. Nelissen I, Martens E, Ven Den Steen PE, Proost P, Ronsse I,

Opdenakker G. Gelatinase B/matrix metalloproteinase-9 cleaves interferon-beta and is a target for immunotherapy. *Brain* 2003; 126:1371-81.

10. Biron CA. Interferons alpha and beta as immune regulators — a new look. *Immunity* 2001;14:661-4.
11. Brassard DL, Grace MJ, Bordens RW. Interferon-alpha as an immunotherapeutic protein. *J Leukoc Biol* 2002;71:565-81.
12. Stylianou E, Yndestad A, Sikkeland LI, et al. Effects of interferon-alpha on gene expression of chemokines and members of the tumour necrosis factor superfamily in HIV-infected patients. *Clin Exp Immunol* 2002;130:279-85.
13. Ronnblom LE, Alm GV, Oberg KE. Autoimmunity after alpha-interferon therapy for malignant carcinoid tumors. *Ann Intern Med* 1991;115:178-83.
14. Ioannou Y, Isenberg DA. Current evidence for the induction of autoimmune rheumatic manifestations by cytokine therapy. *Arthritis Rheum* 2000;43:1431-42.
15. Ronnblom L, Alm GV. A pivotal role for the natural interferon alpha-producing cells (plasmacytoid dendritic cells) in the pathogenesis of lupus. *J Exp Med* 2001;194:F59-63.
16. Neshet G, Ruchlemer R. Alpha-interferon-induced arthritis: clinical presentation, treatment, and prevention. *Semin Arthritis Rheum* 1998; 27:360-5.
17. Kotter I, Müller CA, Einsele H, Mohren M, Kanz L. Interferon-alpha-associated polyarthritis. Possible induction of seropositive rheumatoid arthritis by interferon-alpha: two case reports and review of the literature. *Z Rheumatol* 1999;58:185-95.

Dr. Fish replies

To the Editor:

Drs. Kötter and Stübiger are correct in their discussion of the heterogeneity of response to interferon (IFN). The development of autoimmune phenomena during IFN treatment of patients for hematological diseases and hepatitis C virus (HCV), albeit at a low incidence, is a concern that was duly noted in our report. Interpatient variability within a specific disease population certainly confounds the explanation for susceptibility for these adverse events.

The chronic inflammation in rheumatoid arthritis (RA) that leads to joint destruction is characterized by the accumulation of leukocytes within synovial tissues. In RA, the character of the inflammatory infiltrate that defines the disordered synovial environment fluctuates during the course of disease in affected joints, with regard to cell makeup and activation state. Thus cellular interactions that determine cytokine production are also in a constant state of flux. It is the very nature of IFN- γ as pleiotropic biological response modifiers that suggests their therapeutic potential: their ability to target different cell populations, thereby affecting cellular communication and immunological control at multiple levels.

The challenge, as indicated by Drs. Kötter and Stübiger, is to develop a treatment protocol for IFN intervention that will specifically and preferentially promote the antiinflammatory events we reported. Certainly, Kötter, *et al* recently reported that IFN- α may be superior to the standard immunosuppression treatments for ocular Behçet's disease, yet they identified incidents of autoimmune phenomena in 16% of their patient population. Undoubtedly, the nature of the underlying disease, whether it is a hematological malignancy, a chronic viral infection or an autoimmune disease, the specific target tissues, and the immune-genetic background define the spectrum of potential IFN-inducible outcomes. Thus, IFN- α treatment may predictably invoke autoimmune phenomena in selected patients with HCV that might not develop in patients with RA. As with Behçet's disease, the potential beneficial effects of IFN therapy in a majority of patients with RA may outweigh adverse or exacerbated disease in another specific subset. The intent would be to predetermine responsiveness in each patient, by defining inclusion and exclusion criteria for IFN treatment, based on established

leukocyte and fibroblast responses *ex vivo*. These criteria are under consideration, initially in preclinical studies.

ELEANOR N. FISH, PhD, Director, Arthritis and Autoimmunity Research Centre, University Health Network, 67 College Street, Room 4-424, Toronto, Ontario M5G 2M1, Canada. E-mail: en.fish@utoronto.ca

REFERENCE

1. Kotter I, Zierhut M, Eckstein AK, et al. Human recombinant interferon alpha-2a for the treatment of Behçet's disease with sight threatening posterior or panuveitis. *Br J Ophthalmol* 2003;87:381-2.



False Positive Parvovirus Serology

To the Editor:

In July 1999 we submitted for publication a case report that described the association of an acute symmetrical polyarthropathy with cranial mononeuritis. Parvovirus B19 IgM was present in the serum and no other cause for the arthropathy or neuropathy was found. Our report was provisionally accepted, but at the suggestion of a reviewer, the parvovirus B19 IgM and IgG titers were repeated 14 months after the original assay, as well as polymerase chain reaction (PCR) analysis for parvovirus B19 DNA, at the Central Public Health Laboratory (CPHL) in London, UK. The results are given in Table 1.

In view of these findings, a further opinion was sought from Prof. Klaus Hedman at the University of Helsinki, Finland, whose assay results are illustrated in Table 2.

These results are consistent with past parvovirus B19 infection, and the anomalous parvovirus B19 IgM results are probably false-positive due to nonspecific reactions, and are not related to recent infection (personal communications, Dr. B.J. Cohen, CPHL, London, UK, and Prof. K. Hedman, University of Helsinki, Helsinki, Finland). On reviewing the initial serum sample results showing persistent parvoviral IgM titer, we withdrew our report. We now wish to draw attention to this case as an example of the doubtful specificity of parvovirus B19 IgM serology that may be encountered by investigators. We would recommend that, where a positive parvoviral IgM is detected, B19 DNA PCR may be necessary to confirm recent infection, where this is clinically necessary.

Table 1. Parvovirus B19 results at CPHL.

	Date of Serum Sample	
	September 25, 1998	November 10, 1999
B19 IgM in-house EIA ¹ (T/CO)	3.0 (positive)	5.2 (positive)
B19 IgG Biotrin EIA (T/CO)	8.5 (positive)	8.3 (positive)
B19 DNA PCR	Negative	Negative

EIA: enzyme immunoassay, PCR: polymerase chain reaction. CPHL: Central Public Health Laboratory.

Table 2. Parvovirus B19 IgG avidity and epitope type-specific antibody results.

	Date of Serum Sample	
	September 25, 1998	November 10, 1999
B19 IgG avidity	34%	42%
Acute < 15%		
Past > 25%		
Second-generation ETS-EIA	475	120
Acute < 10		
Past > 10		
B19-DNA	Negative	Negative

ETS-EIA: epitope type-specific enzyme immunoassay.

ALICE H. JOHNSON, MRCP; ANDREW GOUGH, MD, MRCP, Department of Rheumatology, Harrogate District Hospital, Lancaster Park Road, Harrogate HG2 7SX, United Kingdom.
E-mail: andrew.gough@hhc-tr.northy.nhs.uk

REFERENCE

1. Brown KE, Buckley MM, Cohen BJ, Samuel D. An amplified ELISA for the detection of parvovirus B19 IgM using monoclonal antibody to FITC. *J Virol Methods* 1989;26:189-98.