

Extent of Followup Care After Elective Total Hip Replacement

PAOLA de PABLO, ELENA LOSINA, NIZAR MAHOMED, JOHN WRIGHT, ANNE H. FOSSEL, JANE A. BARRETT, and JEFFREY N. KATZ

ABSTRACT. *Objective.* To estimate the extent of radiographic and orthopedic followup among recipients of total hip replacement (THR), and to identify patients who are less likely to have consistent followup over 6 years postoperatively.

Methods. We studied a population-based sample of 622 patients who received THR in 1995. We developed a multivariate ordinal regression model with the extent of radiographic followup as the dependent variable (none, early, and consistent), adjusting for demographic and preoperative clinical characteristics, and hospital and surgeon volume.

Results. Ninety-four (15%) patients indicated that they had no followup radiographs, 269 (43%) had early followup only, and 259 (42%) had consistent followup radiographs over 6 years. Ninety percent of those with consistent followup orthopedic visits also had consistent followup radiographs over 6 years. Multivariate analyses revealed that older patients were less likely to have radiographic followup than younger patients (OR 0.76, 95% CI 0.65, 0.89) per each 5-year increase in age. Subjects with no college education were less likely to have radiographic followup than those with more education (OR 0.58, 95% CI 0.41, 0.83), and those with lower income were less likely to have radiographic followup than those with a higher income (OR 0.50, 95% CI 0.27, 0.92).

Conclusion. Only 42% of THR recipients reported consistent radiographic followup. Older patients, patients with lower income, and those with lower education level were less likely to have consistent radiographic followup over 6 years after THR. These population groups can serve as targets for interventions to improve followup after elective THR. (J Rheumatol 2006;33:1159–66)

Key Indexing Terms:

TOTAL HIP REPLACEMENT ORTHOPEDIC RADIOGRAPHIC FOLLOWUP

From the Department of Medicine, Division of Rheumatology, Immunology and Allergy, Section of Clinical Sciences; and Department of Orthopedic Surgery, Brigham and Women's Hospital, Harvard Medical School; Department of Biostatistics, Boston University School of Public Health; Department of Medicine, Boston University Medical School; Department of Orthopedic Surgery, Toronto Western Hospital, University Health Network, University of Toronto, Toronto, Ontario, Canada; and Department of Community and Family Medicine, Dartmouth Medical School, Hanover, New Hampshire, USA.

Supported by NIH Grants P01 AR47782 and K24 AR02123, AHRQ Grant 1R01 HS09775-01, and a Clinical Science Grant from the Arthritis Foundation. Dr. de Pablo was supported by a Post Doctoral Fellowship from the Arthritis Foundation.

P. de Pablo, MD, MPH, Post-doctoral Fellow; A.H. Fossel, Division of Rheumatology, Immunology and Allergy, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School; E. Losina, PhD, Division of Rheumatology, Immunology and Allergy, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Assistant Professor of Biostatistics, Department of Epidemiology and Biostatistics, Boston University School of Public Health and Department of Medicine, Boston University Medical School; N. Mahomed, MD, ScD, Associate Professor of Orthopedic Surgery, Department of Orthopedic Surgery, Toronto Western Hospital, University of Toronto; J. Wright, MD, Assistant Professor of Orthopedic Surgery, Department of Orthopedic Surgery, Brigham and Women's Hospital, Harvard Medical School; J.A. Barrett, MS, Assistant Research Professor, Department of Community and Family Medicine, Dartmouth Medical School; J.N. Katz, MD, MS, Assistant Professor of Medicine and Orthopedic Surgery, Division of Rheumatology, Immunology and Allergy, Department of Medicine, and Department of Orthopedic Surgery Brigham and Women's Hospital, Harvard Medical School.

Address reprint requests to Dr. J.N. Katz, Section of Clinical Sciences, Division of Rheumatology, Brigham and Women's Hospital, 75 Francis Street, Boston, MA 02115. E-mail: jnkatz@partners.org

Accepted for publication January 26, 2006.

Total hip replacement (THR) predictably relieves pain and improves function for patients with painful arthritic hips¹⁻⁶. Joint replacement surgery remains one of the most effective treatments for severe arthritis affecting the hips, with excellent longterm results exceeding 20 years in many cases⁷⁻¹². However, aseptic loosening due to osteolysis (inflammatory reaction to wear debris)¹³ represents the largest threat to longterm survival of the implants. Other reasons for failure include dislocation, periprosthetic fracture, stem breakage, cup malposition, dissociated insert, dissociated femoral head, and infection¹⁴.

The incidence of osteolysis increases progressively with time¹⁵. A recent study found osteolysis in 41% of hips at 7 years of followup¹⁶. While osteolysis may ultimately manifest as component loosening associated with pain, it may be silent, particularly in its early phases¹⁷⁻²⁴. The evaluation and treatment of osteolytic defects identified in asymptomatic patients is controversial. However, osteolysis may cause enough bone loss to require revision THR⁶. Revision procedures are more costly and often more complicated than the index surgery^{17,23,25-29}. Moreover, if unrecognized, osteolysis can lead to significant bone stock deficiency, making revision THR much more complex³⁰⁻³³, often requiring bone allograft to restore adequate support^{34,35}. Osteolysis can be detected as a progressive radiolucent line or cavity at the implant-bone or cement-bone interface on radiographs^{24,36-39}. If osteolysis is

identified early, frequent monitoring strategies and, eventually, implementation of novel therapies may help delay or even prevent the need for more complex revision surgery if failures are detected sooner⁴⁰⁻⁴⁶.

An essential feature of any surgical procedure is postoperative followup. Regular followup with an orthopedic surgeon, including radiographs, enables the surgeon to assess the result of surgery, recognize early signs of osteolysis, and identify the possible need for revision at an early stage^{47,48}. Lack of effective followup may make any necessary surgery more difficult and potentially less successful, as conditions that may require a revision operation are not identified early.

Since bone loss may be asymptomatic, longterm followup is essential to determining outcomes and pathological processes related to THR^{6,49}. However, there are no explicit guidelines for longterm followup care after THR following discharge.

The ideal extent of orthopedic and radiographic followup after THR has not been clearly defined. One published clinical pathway recommends visits to the orthopedic surgeon at 6 weeks postsurgery, 3 months, 6 months, and 1 year after THR, and then every 2 years⁵⁰. In addition, evaluation with screening followup radiographs is typically performed in some settings in the immediate postoperative period and at followup every 2 years, with shorter intervals between visits if osteolysis becomes apparent³¹. Other guidelines suggest periodic longterm followup⁴⁹ or a longterm followup visit at a minimum after 5 years and every 5 years thereafter, with a minimum requirement of an anteroposterior and lateral radiograph^{47,51}.

Despite these recommendations, the frequency and duration of outpatient followup appointments after THR appear to vary greatly across orthopedic surgeons and hospitals⁵². In a recent survey of the American Association of Hip and Knee Surgeons, 80% of respondents recommended annual or biennial orthopedic clinical and radiographic examinations after THR, with more frequent followup times for clinical or radiographic signs of failure, previous revision arthroplasty, previous joint sepsis, and subnormal periprosthetic bone quality⁵³. However, to our knowledge there has not been a population-based study examining the extent of clinical orthopedic and radiographic followup after THR.

The purpose of our study was to estimate the extent of radiographic and orthopedic followup among recipients of primary THR, and to identify patients who are less likely to have consistent followup over 6 years postoperatively. We hypothesized that longterm radiographic and orthopedic followup is not performed in all THR recipients consistently, and that the extent of followup varies according to patient characteristics.

MATERIALS AND METHODS

Patients. To select the study cohort, we used Medicare claims to identify a random sample of patients aged 65 years or older that were resident in 3 US states (Ohio, Pennsylvania, and Colorado) and had elective primary THR in calendar year 1995, as reported^{54,55}. The analyses here are built upon surveys administered to this population-based cohort of patients 3 and 6 years after they received elective primary THR.

Data sources

Medical record. Medical record reviews were performed by trained personnel working in peer review organizations using a standardized data abstraction form. The medical record included data on patients' preoperative clinical characteristics such as age, sex, weight and height, the primary underlying joint disease (osteoarthritis, rheumatoid arthritis, or avascular necrosis), the surgical procedure performed (index vs revision), history of previous orthopedic surgery in other joints, and comorbid medical conditions. Comorbid medical conditions were extracted and aggregated using the Charlson Index⁵⁶. Body mass index (BMI) was computed as weight (kg)/height (m)² and was dichotomized for obesity (overweight grade II: BMI > 30)⁵⁷.

Medicare claims. Medicare claims provided data on the volume of primary and revision THR performed in 1995 by the surgeon and the hospital.

Survey questionnaires. The 3-year followup questionnaire included questions about patients' recalled preoperative functional status (including use of supportive devices, limp, stair climbing, and walking distance)⁵⁸. These items were adapted from the Harris Hip Score^{4,5,59}. Weights were assigned as in the Harris Score, summed, and standardized to a 0-100 scale. This variable was dichotomized at the highest quartile. Patients also indicated their socioeconomic status, living arrangement, and number of years of formal education completed. Income consisted of the total household income in the past year (including all sources of income such as wages, social security, pensions, investments, etc.). The 3 and 6-year followup questionnaires assessed patients' pain and functional status with the Western Ontario and McMaster University Osteoarthritis Index (WOMAC)⁶⁰ and patients' mental health status with the 5-item Mental Health Inventory Questionnaire (MHI-5)⁶¹. The questionnaires asked whether patients had followup visits with an orthopedic surgeon and followup radiographs after primary THR.

Statistical methods. The primary outcome was the extent of radiographic followup visits in THR recipients. The dependent variable categories were defined as follows: None: no followup radiographs reported (3 or 6 years) after THR. Early: at least one followup radiograph reported at 0-3 years after THR, but no radiographs at 3-6 years after THR. Consistent: at least one followup radiograph reported at 0-3 years and one followup radiograph reported at 3-6 years after THR.

Definitions of none, early, and consistent orthopedic surgeon followup appointments were made analogously. Demographic data were summarized using descriptive statistics. The independent variables included patient sociodemographic characteristics (age, sex, race, living arrangement, level of income, level of education, type of residence, distance from home to hospital⁶²), preoperative factors (obesity, comorbidities, underlying disease, mental health, and preoperative functional status), and hospital and surgeon characteristics (hospital and surgeon volume and surgeon's age). Analyses of the independent variable "income" included "missing" as a category.

We analyzed the bivariate relationship of each predictor variable to the outcome in order to select candidate predictors for inclusion in the multivariate model. A significance level of 0.10 was applied for selecting predictors. We developed ordinal regression models (also known as proportional odds models) using the 3-level dependent variable noted above, with those having consistent followup as the reference group. In these models, the odds ratio can be interpreted as the effect of an explanatory variable on the odds of having early followup versus no followup, and the odds of having consistent followup versus early followup⁶³. These analyses were adjusted for all predictors that were retained in the model in a stepwise selection with a criterion of $p = 0.10$. These variables included age, sex, annual income, education level, and hospital volume. In addition, other relevant variables such as obesity, comorbidities, preoperative functional status, and distance from home to hospital were also included in the model.

We performed a sensitivity analysis in which the primary outcome measure was defined as a dichotomous indicator for the extent of followup in THR recipients (consistent vs early/none). For this outcome we built a multivariate logistic regression model that adjusted for all predictors that were retained in the model in a stepwise selection, with a selection criterion of $p = 0.10$. These variables included age, sex, income, education level, obesity, comorbidities,

preoperative functional status, distance from home to hospital, and hospital volume.

RESULTS

Patients. We selected a cohort of 1939 patients with primary THR, using the sampling procedures described above. The algorithms for case identification in this cohort are published^{54,55}. Briefly, 956 patients of all those eligible to be involved in the sample returned questionnaires at 3 years of followup. Of these, 177 (19%) patients indicated that they had no followup visits with the orthopedic surgeon in the first 3 years following surgery, 382 (40%) had visits less than yearly, and 336 (35%) had yearly followup visits over this first 3-year period; 61 (6%) had missing values. On the other hand, 163 (17%) patients indicated that they had no followup radiographs in the first 3 years following surgery, 417 (44%) had followup radiographs less than yearly, and 317 (33%) had consistent followup radiographs over the first 3-year period; 59 (6%) had missing values.

At 6 years, we approached 907 patients (all those who agreed at 3 years except those who had died). A total of 792 (87%) returned completed questionnaires at 6 years of followup. Among these, 170 (21%) patients with missing values on the dependent variable were excluded from the analysis.

The 334 patients not included in the study sample at 6 years because they had died, did not respond, or had missing data on the dependent variable were similar to those included in the analyses except that they were 3 years older than those who were included (75 vs 72 yrs; $p = 0.0001$). Also, a lower proportion of subjects not included had incomes $> \$20,000$ US than those who were included (36% vs 46%; $p = 0.005$). There was no difference in race.

The study sample included 622 patients who had a primary THR in 1995 and completed the survey at 3 and 6 years postoperatively. The demographic characteristics of the study sample are listed in Table 1, as are preoperative clinical factors and surgeon and hospital characteristics. The mean age was 72 years, 62% were female, 55% had an education level of a high school degree or less, 34% had an annual income $< \$20,000$, less than 10% were employed, and 30% were living alone at the time of completing the 6-year survey.

Extent of radiographic and orthopedic followup. Ninety five (16%) patients indicated that they had no followup visits with the orthopedic surgeon, 256 (43%) had visits only in the first 3 years after surgery (early followup), and 247 (41%) had consistent followup visits over a 6-year period. Ninety-four (15%) patients indicated that they had no followup radiographs, 269 (43%) had followup radiographs only in the first 3 years after surgery (early followup), and 259 (42%) had consistent followup radiographs over 6 years. Ninety percent of those with consistent followup orthopedic visits also had consistent followup radiographs over 6 years after THR. We give results based on followup radiographs as the primary outcome measure.

Ordinal regression results are shown in Table 2. Multivariate ordinal regression analyses included age, sex, income, education level, distance from home to hospital, obesity, comorbidities, preoperative functional status, and hospital volume as potential predictors of radiographic followup. These analyses revealed that older patients were less likely to have radiographic followup than younger patients (OR 0.76, 95% CI 0.65, 0.89, per each 5-year increase in age). Subjects with no college were less likely to have radiographic followup than those with college or at least some college education (OR 0.58, 95% CI 0.41, 0.83). Finally, those with lower income were less likely to have radiographic followup than those with a higher income (OR 0.50, 95% CI 0.27, 0.92 for subjects with an income $< \$20,000$; and OR 0.50, 95% CI 0.29, 0.88 for those with an income in the range of $\$20$ – $50,000$).

Other factors such as sex, race, living arrangement, distance from home to hospital, obesity, comorbidities, mental health, and functional status were not associated with radiographic followup (Table 2).

A sensitivity analysis using logistic regression showed similar results to those of the ordinal regression models. These logistic regression analyses confirmed that older patients were less likely to have radiographic followup than younger patients (OR per each 5-year increase in age 0.72, 95% CI 0.61, 0.87). Those with a lower level of education (OR 0.62, 95% CI 0.43, 0.91) compared with those with greater than high school education, and those with lower income were less likely to have radiographic followup than those with a higher income (OR 0.55, 95% CI 0.29, 1.05 for subjects with an income $< \$20,000$; and OR 0.48, 95% CI 0.27, 0.86 for those with income in the range of $\$20$ – $50,000$).

DISCUSSION

We used patient survey data in a population-based sample of Medicare beneficiaries to describe practice patterns for radiographic and orthopedic followup after primary elective THR. Our results showed that among those who responded to the survey, 15% of THR recipients get no followup care and that only about 40% have consistent followup over 6 years, with the remainder having followup visits with an orthopedic surgeon as well as radiographic followup only in the first 3 years after THR. In addition, we identified factors related to reduced followup frequency. Older patients and patients with lower level of education and lower income were less likely to have consistent followup with radiographs over 6 years.

Other factors such as sex, race, living arrangement, distance from home to hospital, obesity, comorbidities, mental health, and functional status did not appear to influence the extent of followup. However, other factors may influence the patients' lack of understanding of why followup evaluation might be helpful, even if their orthopedic surgeon had suggested followup at the time of surgery.

Several authors have suggested that even asymptomatic patients should have followup care at least biennially follow-

Table 1. Study sample characteristics and radiographic followup.

Patient Characteristics	No Followup, N = 94 (%)	Early Followup, N = 269 (%)	Consistent Followup, N = 259 (%)	p
Sociodemographic factors				
Age, 5 yr categories				
65–70	26 (12)	86 (39)	106 (49)	0.005
70–75	36 (17)	90 (42)	89 (41)	
75–80	20 (16)	65 (51)	43 (34)	
≥ 80	12 (20)	28 (46)	21 (34)	
Female	57 (15)	163 (42)	166 (43)	0.44
Male	37 (16)	106 (45)	93 (39)	
Race				
White	92 (15)	262 (43)	251 (42)	0.27
Non-white	1 (7)	6 (40)	8 (53)	
Living arrangement				
Alone	28 (14)	88 (47)	73 (39)	0.52
Not alone	66 (15)	181 (42)	186 (43)	
Income, \$US				
High (≥ 50,000)	5 (7)	26 (37)	39 (56)	0.01
Medium (20–50,000)	30 (14)	104 (49)	80 (37)	
Low (≤ 20,000)	42 (20)	91 (43)	79 (37)	
Missing data	17 (14)	48 (38)	61 (48)	
Education level				
College and at least some college	28 (10)	113 (41)	132 (48)	< 0.001
No college	65 (19)	149 (44)	123 (36)	
Distance to hospital, miles				
Low (≤ 5)	25 (13)	80 (41)	88 (46)	0.27
Medium (5–20)	40 (16)	117 (46)	99 (39)	
High (≥ 20)	29 (17)	72 (42)	72 (42)	
Preoperative clinical factors				
Obesity (BMI ≥ 30 kg/m ²)				
Yes	17 (11)	79 (53)	52 (35)	0.57
No	77 (16)	190 (40)	207 (44)	
Comorbidities (≥ 1)				
Yes	41 (15)	130 (47)	103 (38)	0.22
No	53 (15)	139 (40)	156 (45)	
Underlying disease				
OA	78 (14)	240 (44)	227 (42)	0.44
Other	16 (21)	29 (38)	32 (42)	
Functional status (highest quartile)				
High	48 (15)	132 (42)	133 (42)	0.82
Low	46 (15)	137 (44)	126 (41)	
Surgeon and hospital factors				
Surgeon's age, yrs				
≥ 60	10 (8)	53 (45)	56 (47)	0.03
< 60	84 (17)	216 (43)	203 (40)	
Surgeon volume				
High (≥ 15)	30 (14)	100 (47)	85 (40)	0.61
Medium (5–15)	25 (13)	81 (42)	88 (45)	
Low (≤ 5)	39 (18)	88 (41)	86 (40)	
Hospital volume				
High (≥ 50)	37 (18)	91 (44)	77 (38)	0.51
Medium (15–50)	23 (11)	91 (43)	99 (46)	
Low (≤ 15)	34 (17)	87 (43)	83 (41)	
Teaching hospital				
Yes	57 (15)	166 (43)	163 (42)	0.64
No	35 (16)	95 (43)	89 (41)	

BMI: body mass index, OA: osteoarthritis.

ing THR⁵³. While most cases of clinically significant osteolysis are identified later than 6 years after the procedure, and followup would be expected to decrease with time from surgery,

detection of clinically silent problems may be enhanced by early, regular, and consistent followup. This strategy permits identification of potential complications at an earlier stage and

Table 2. Radiographic followup after THR: ordinal regression results.

Patient Characteristics	N (%)	Crude OR	Adjusted OR*
Sociodemographic factors			
Age (ordinal, 5 yr categories)		0.80 (0.69, 0.93)	0.76 (0.65, 0.89)
Sex, female	386 (62)	1.13 (0.84, 1.54)	1.33 (0.95, 1.87)
Race, White	605 (97)	0.58 (0.21, 1.59)	
Living arrangement, alone	189 (30)	0.88 (0.63, 1.22)	
Income, \$US			
High (\geq 50,000)	70 (11)	1	1
Medium (20–50,000)	214 (34)	0.43 (0.20, 0.73)	0.50 (0.29, 0.88)
Low (\leq 20,000)	212 (34)	0.49 (0.29, 0.83)	0.50 (0.27, 0.92)
Missing data	126 (20)	0.69 (0.39, 1.23)	0.80 (0.43, 1.48)
Education level			
College and some college	273 (45)	1	1
No college	337 (55)	0.57 (0.42, 0.78)	0.58 (0.41, 0.83)
Distance to hospital, miles			
Low (\leq 5)	193 (31)	1	1
Medium (5–20)	256 (41)	0.77 (0.54, 1.10)	0.74 (0.51, 1.07)
High (\geq 20)	173 (28)	0.82 (0.56, 1.21)	0.88 (0.58, 1.35)
Preoperative clinical factors			
Obesity (BMI \geq 30 kg/m ²)	148 (24)	0.86 (0.61, 1.21)	0.91 (0.63, 1.31)
Comorbidities (\geq 1)	274 (44)	0.81 (0.60, 1.09)	0.82 (0.60, 1.13)
Underlying disease, OA	545 (88)	1.15 (0.74, 1.81)	
Functional status (highest quartile)			
High (\geq 21)	313 (50)	1	1
Low ($<$ 21)	309 (50)	1.04 (0.78, 1.40)	1.03 (0.75, 1.41)
Surgeon and hospital factors			
Surgeon's age (\geq 60 yrs)	119 (19)	1.44 (0.98, 2.12)	
Surgeon volume			
High (\geq 15)	215 (35)	1	
Medium (5–15)	194 (31)	1.22 (0.84, 1.76)	
Low (\leq 5)	213 (34)	0.93 (0.65, 1.33)	
Hospital volume			
High (\geq 50)	205 (33)	1	
Medium (15–50)	213 (34)	1.52 (1.06, 2.19)	
Low (\leq 15)	204 (33)	1.13 (0.79, 1.63)	
Teaching hospital	386 (64)	1.08 (0.79, 1.47)	

* Adjusted for age, sex, race, income, education level, distance from home to hospital, obesity, comorbidities, preoperative functional status, and hospital volume.

hence may reduce the likelihood of complex revision procedures^{16,21}. Our results indicate that these goals are not met in the majority of patients in a 6-year postoperative period.

Previous studies found that low income was associated with less favorable outcomes after total joint arthroplasty⁶⁴. We found that older people and people with lower income and less education were less likely to receive regular followup. It is possible that healthcare providers offer these patients less information, and/or that the patients are less able to understand the information. Also, Medicare does not cover all medical costs, and these people may not have supplementary insurance or the ability to pay the uncovered amounts for followup appointments. In any case, our results suggest that intervention strategies are necessary to improve followup, specifically in patients who are older and have lower levels of education and income.

Major advantages of this study include the large sample

size and population-based sampling strategy. Further, the study considered the level of function prior to surgery (albeit retrospectively)^{58,65,66}. Also, the response rate is 87% of all those eligible to be involved in the sample at 6 years of followup. However, a major limitation of the study is the low response rate to the 3-year survey. Several authors have suggested that patients who do not respond to followup surveys have worse outcomes of total joint arthroplasty⁶⁷⁻⁶⁹. In contrast, a recent study found that patients who did not attend followup visits with the orthopedic surgeon after total knee replacement did not have significant differences in outcome variables or surgical procedures compared with patients who had complied with a followup protocol⁷⁰. We can only speculate to what extent nonresponse influenced our results. It is impossible to tell from our study whether inconsistent followup is cause or effect of nonresponse. Future studies should attempt to disentangle these complex mechanisms.

In addition, we acknowledge that this cohort is older than 65 years and therefore we are uncertain what happens in younger patients. Orthopedic surgeons might be more aggressive in following younger, more active patients who are at higher risk of failure and hence more likely to need a revision procedure.

Data on followup orthopedic visits and radiographs were recalled and thus were subject to recall bias. We were unable to address this potential bias since we could not perform a medical record confirmation of followup visits. In general, immediate postoperative radiographs are obtained, since they serve as a baseline for identification of osteolysis on later studies, and comparison radiographs are important for decision-making. Also, most institutions follow a standard protocol for postoperative care following THR including a routine visit and radiographs. Our surveys collected information on whether patients had followup appointments with an orthopedic surgeon and followup radiographs after primary THR, but we were unable to distinguish whether they had no followup visits at all or if they had none after the routine postoperative visit. However, by potentially including early routine postoperative visits and radiographs in our definition of the dependent variable, we may overestimate the extent of followup care after primary elective hip replacement. Yet our data show a concerning lack of followup care. Thus the bias toward greater followup care is conservative. In addition, this ambiguity does not arise in the questionnaire at 6 years after the procedure, which is more important in assessing longterm followup care.

Our survey did not include questions on how orthopedic surgeons or health professionals communicate longterm followup care recommendations, nor on patient recall strategies that could facilitate compliance with followup visits. Indeed we do not know what percentage of patients actually comply with recommendations given at discharge from the acute care setting and subsequent followup visits, nor whether these recommendations follow a clinical pathway or not. However, 2 surveys of surgeons in the UK and the US suggested that consistent followup is recommended for all patients^{52,53}. Further research is needed to correlate followup care type and frequency with outcomes and complications.

The principal finding of our survey is that a majority of patients in these series did not receive consistent followup care after total hip arthroplasty over a 6-year period. Healthcare providers may not be aware of the importance of longterm orthopedic and radiographic followup care in every recipient of THR. Our results may be useful in the development of practice guidelines for followup after total hip arthroplasty that may be used by healthcare providers involved in the care of THR recipients, such as orthopedic surgeons, rheumatologists, primary care physicians, radiologists, nurses, and physical therapists. Further, the results suggest that low income, low education, and older age may be independent predictors of inconsistent followup. Hence, specific efforts targeted to these

patient groups may be necessary to improve followup and thus less complicated revision surgeries after THR.

REFERENCES

1. Liang MH, Cullen KE, Larson MG, et al. Cost-effectiveness of total joint arthroplasty in osteoarthritis. *Arthritis Rheum* 1986;29:937-43.
2. Laupacis A, Bourne R, Rorabeck C, et al. The effect of elective total hip replacement on health-related quality of life. *J Bone Joint Surg Am* 1993;75:1619-26.
3. Rissanen P, Aro S, Slati P, Sintonen H, Paavolainen P. Health and quality of life before and after hip or knee arthroplasty. *J Arthroplasty* 1995;10:169-75.
4. Harris WH, Sledge CB. Total hip and total knee replacement (1). *N Engl J Med* 1990;323:725-31.
5. Harris WH, Sledge CB. Total hip and total knee replacement (2). *N Engl J Med* 1990;323:801-7.
6. National Institutes of Health. National Institutes of Health consensus conference: Total hip replacement. NIH Consensus Development Panel on Total Hip Replacement. *JAMA* 1995;273:1950-6.
7. Kawamura H, Dunbar MJ, Murray P, Bourne RB, Rorabeck CH. The porous coated anatomic total hip replacement. A ten to fourteen-year follow-up study of a cementless total hip arthroplasty. *J Bone Joint Surg Am* 2001;83:1333-8.
8. Brown SR, Davies WA, DeHeer DH, Swanson AB. Long-term survival of McKee-Farrar total hip prostheses. *Clin Orthop Relat Res* 2002;157-63.
9. Hamadouche M, Boutin P, Daussange J, Bolander ME, Sedel L. Alumina-on-alumina total hip arthroplasty: a minimum 18.5-year follow-up study. *J Bone Joint Surg Am* 2002;84:69-77.
10. Callaghan JJ, Albright JC, Goetz DD, Olejniczak JP, Johnston RC. Charnley total hip arthroplasty with cement. Minimum twenty-five-year follow-up. *J Bone Joint Surg Am* 2000;82:487-97.
11. Huo MH. What's new in hip arthroplasty. *J Bone Joint Surg Am* 2002;84:1894-905.
12. Berry DJ, Harmsen WS, Cabanela ME, Morrey BF. Twenty-five-year survivorship of two thousand consecutive primary Charnley total hip replacements: factors affecting survivorship of acetabular and femoral components. *J Bone Joint Surg Am* 2002;84:171-7.
13. Archibeck MJ, Jacobs JJ, Black J. Alternate bearing surfaces in total joint arthroplasty: biologic considerations. *Clin Orthop Relat Res* 2000;379:12-21.
14. Berend KR, Lombardi AV Jr, Mallory TH, Adams JB, Russell JH, Groseth KL. The long-term outcome of 755 consecutive constrained acetabular components in total hip arthroplasty examining the successes and failures. *J Arthroplasty* 2005;20 Suppl 3:93-102.
15. Harris WH. "The lysis threshold": an erroneous and perhaps misleading concept? *J Arthroplasty* 2003;18:506-10.
16. Orishimo KF, Claus AM, Sychterz CJ, Engh CA. Relationship between polyethylene wear and osteolysis in hips with a second-generation porous-coated cementless cup after seven years of follow-up. *J Bone Joint Surg Am* 2003;85:1095-9.
17. Berry DJ. Management of osteolysis around total hip arthroplasty. *Orthopedics* 1999;22:805-8.
18. Lonner JH, Siliski JM, Scott RD. Prodromes of failure in total knee arthroplasty. *J Arthroplasty* 1999;14:488-92.
19. O'Rourke MR, Callaghan JJ, Goetz DD, Sullivan PM, Johnston RC. Osteolysis associated with a cemented modular posterior-cruciate-substituting total knee design: five to eight-year follow-up. *J Bone Joint Surg Am* 2002;84:1362-71.

20. Harris WH. Wear and periprosthetic osteolysis: the problem. *Clin Orthop* 2001;393:66-70.
21. Lavernia CJ. Cost-effectiveness of early surgical intervention in silent osteolysis. *J Arthroplasty* 1998;13:277-9.
22. Jacobs JJ, Shanbhag A, Glant TT, Black J, Galante JO. Wear debris in total joint replacements. *J Am Acad Orthop Surg* 1994;2:212-20.
23. Khalily C, Whiteside LA. Predictive value of early radiographic findings in cementless total hip arthroplasty femoral components: an 8- to 12-year follow-up. *J Arthroplasty* 1998;13:768-73.
24. Iwase T, Wingstrand I, Persson BM, Kesteris U, Hasegawa Y, Wingstrand H. The ScanHip total hip arthroplasty: radiographic assessment of 72 hips after 10 years. *Acta Orthop Scand* 2002;73:54-9.
25. Lavernia CJ, Drakeford MK, Tsao AK, Gittelsohn A, Krackow KA, Hungerford DS. Revision and primary hip and knee arthroplasty. A cost analysis. *Clin Orthop* 1995;311:136-41.
26. Barrack RL, Hoffman GJ, Tejeiro WV, Carpenter LJ Jr. Surgeon work input and risk in primary versus revision total joint arthroplasty. *J Arthroplasty* 1995;10:281-6.
27. Barrack RL, Sawhney J, Hsu J, Cofield RH. Cost analysis of revision total hip arthroplasty. A 5-year followup study. *Clin Orthop* 1999;369:175-8.
28. Barrack RL. The evolving cost spectrum of revision hip arthroplasty. *Orthopedics* 1999;22:865-6.
29. Crowe JF, Sculco TP, Kahn B. Revision total hip arthroplasty: hospital cost and reimbursement analysis. *Clin Orthop* 2003;413:175-82.
30. Kershaw CJ, Atkins RM, Dodd CA, Bulstrode CJ. Revision total hip arthroplasty for aseptic failure. A review of 276 cases. *J Bone Joint Surg Br* 1991;73:564-8.
31. Roberson JR. Proximal femoral bone loss after total hip arthroplasty. *Orthop Clin North Am* 1992;23:291-302.
32. Eldridge JD, Smith EJ, Hubble MJ, Whitehouse SL, Learmonth ID. Massive early subsidence following femoral impaction grafting. *J Arthroplasty* 1997;12:535-40.
33. Meding JB, Ritter MA, Keating EM, Faris PM. Impaction bone-grafting before insertion of a femoral stem with cement in revision total hip arthroplasty. A minimum two-year follow-up study. *J Bone Joint Surg Am* 1997;79:1834-41.
34. Gie GA, Linder L, Ling RS, Simon JP, Slooff TJ, Timperley AJ. Impacted cancellous allografts and cement for revision total hip arthroplasty. *J Bone Joint Surg Br* 1993;75:14-21.
35. Slooff TJ, Buma P, Schreurs BW, Schimmel JW, Huiskes R, Gardeniers J. Acetabular and femoral reconstruction with impacted graft and cement. *Clin Orthop* 1996;324:108-15.
36. DeLee JG, Charnley J. Radiological demarcation of cemented sockets in total hip replacement. *Clin Orthop* 1976;121:20-32.
37. Charnley J. *Low friction arthroplasty of the hip*. Berlin: Springer-Verlag; 1979.
38. Gruen TA, McNeice GM, Amstutz HC. "Modes of failure" of cemented stem-type femoral components: a radiographic analysis of loosening. *Clin Orthop* 1979;141:17-27.
39. Schmalzried TP, Kwong LM, Jasty M, et al. The mechanism of loosening of cemented acetabular components in total hip arthroplasty. Analysis of specimens retrieved at autopsy. *Clin Orthop* 1992;274:60-78.
40. Clohisy JC TS, Ross FP, et al. Blockade of TNF-activation of NF-kB in osteoclast precursors prevents implant osteolysis. *J Bone Miner Res* 1999;14 Suppl:S489.
41. Schwarz EM, Looney RJ, O'Keefe RJ. Anti-TNF-alpha therapy as a clinical intervention for periprosthetic osteolysis. *Arthritis Res* 2000;2:165-8.
42. Childs LM, Goater JJ, O'Keefe RJ, Schwarz EM. Efficacy of etanercept for wear debris-induced osteolysis. *J Bone Miner Res* 2001;16:338-47.
43. Schwarz EM, Campbell D, Totterman S, Boyd A, O'Keefe RJ, Looney RJ. Use of volumetric computerized tomography as a primary outcome measure to evaluate drug efficacy in the prevention of peri-prosthetic osteolysis: a 1-year clinical pilot of etanercept vs. placebo. *J Orthop Res* 2003;21:1049-55.
44. Millett PJ, Allen MJ, Bostrom MP. Effects of alendronate on particle-induced osteolysis in a rat model. *J Bone Joint Surg Am* 2002;84:236-49.
45. Thadani PJ, Waxman B, Sladek E, Barmada R, Gonzalez MH. Inhibition of particulate debris-induced osteolysis by alendronate in a rat model. *Orthopedics* 2002;25:59-63.
46. Nehme A, Maalouf G, Tricoire JL, Giordano G, Chiron P, Puget J. Effect of alendronate on periprosthetic bone loss after cemented primary total hip arthroplasty: a prospective randomized study [French]. *Rev Chir Orthop Reparatrice Appar Mot* 2003;89:593-8.
47. UK National Audit Office. Hip replacements: Getting it right first time. Report by the Comptroller and Auditor General, 19 April 2000. Available from: http://www.nao.org.uk/publications/nao_reports/9900417.pdf. Accessed March 10, 2006.
48. Mehin R, Yuan X, Haydon C, et al. Retroacetabular osteolysis: When to operate? *Clin Orthop Relat Res* 2004;428:247-55.
49. Ritter MA, Albohm MJ. Overview: maintaining outcomes for total hip arthroplasty. The past, present, and future. *Clin Orthop Relat Res* 1997;344:81-7.
50. Howe JG. Critical pathways in total hip arthroplasty. In: Callaghan JJ, Rosenberg AG, Rubash HE, editors. *The adult hip*. Philadelphia: Lippincott-Raven; 1998.
51. British Orthopaedic Association. *Total hip replacement: a guide to best practice*. London: British Orthopaedic Association; 1999.
52. Veysi VT, Jones S, Stone MH, Limb D. Out-patient follow-up after total hip replacement in one health region. *J R Coll Surg Edinb* 1998;43:95-6.
53. Teeny SM, York SC, Mesko JW, Rea RE. Long-term follow-up care recommendations after total hip and knee arthroplasty: results of the American Association of Hip and Knee Surgeons' member survey. *J Arthroplasty* 2003;18:954-62.
54. Katz JN, Losina E, Barrett J, et al. Association between hospital and surgeon procedure volume and outcomes of total hip replacement in the United States medicare population. *J Bone Joint Surg Am* 2001;83:1622-9.
55. Katz JN, Phillips CB, Baron JA, et al. Association of hospital and surgeon volume of total hip replacement with functional status and satisfaction three years following surgery. *Arthritis Rheum* 2003;48:560-8.
56. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chron Dis* 1987;40:373-83.
57. US National Institutes of Health. Health implications of obesity. National Institutes of Health Consensus Development Conference Statement. *Ann Intern Med* 1985;103:147-51.
58. Mancuso CA, Charlson ME. Does recollection error threaten the validity of cross-sectional studies of effectiveness? *Med Care* 1995;33 Suppl:AS77-88.
59. Mahomed NN, Arndt DC, McGrory BJ, Harris WH. The Harris hip score: comparison of patient self-report with surgeon assessment. *J Arthroplasty* 2001;16:575-80.
60. Bellamy N, Buchanan WW, Goldsmith CH, Campbell J, Stitt LW. Validation study of WOMAC: a health status instrument for

measuring clinically important patient relevant outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip or knee. *J Rheumatol* 1988;15:1833-40.

61. American Psychiatric Association. Diagnostic and statistical manual, text revision. 4th ed. Arlington, VA: American Psychiatric Association; 2000.
62. Losina E, Barrett J, Baron JA, Levy M, Phillips CB, Katz JN. Utilization of low-volume hospitals for total hip replacement. *Arthritis Rheum* 2004;51:836-42.
63. Snedker K, Glynn P, Wang C. Ordered/ordinal logistic regression with SAS and STATA; 2002. Available from: <http://staff.washington.edu/~glynn/olr.pdf>. Accessed March 10, 2006.
64. Mahomed NN, Barrett JA, Katz JN, et al. Rates and outcomes of primary and revision total hip replacement in the United States medicare population. *J Bone Joint Surg Am* 2003;85:27-32.
65. Fortin PR, Clarke AE, Joseph L, et al. Outcomes of total hip and knee replacement: preoperative functional status predicts outcomes at six months after surgery. *Arthritis Rheum* 1999;42:1722-8.
66. Lingard EA, Wright EA, Sledge CB. Pitfalls of using patient recall to derive preoperative status in outcome studies of total knee arthroplasty. *J Bone Joint Surg Am* 2001;83:1149-56.
67. Wildner M. Lost to follow-up. *J Bone Joint Surg Br* 1995;77:657.
68. Murray DW, Britton AR, Bulstrode CJ. Loss to follow-up matters. *J Bone Joint Surg Br* 1997;79:254-7.
69. Kim J, Lonner JH, Nelson CL, Lotke PA. Response bias: effect on outcomes evaluation by mail surveys after total knee arthroplasty. *J Bone Joint Surg Am* 2004;86:15-21.
70. King PJ, Malin AS, Scott RD, Thornhill TS. The fate of patients not returning for follow-up five years after total knee arthroplasty. *J Bone Joint Surg Am* 2004;86:897-901.