Drs. Marasini and Massarotti reply

To the Editor:

The hypothesis suggested by Day and colleagues, consistent with recent data on the relationship between adipose tissue, inflammation, and the renin-angiotensin system (RAS), is relevant and deserves attention¹. However, data on uric acid excretion and serum urate concentrations and the effects brought about by different RAS-blocking drugs are controversial^{2,3}.

RAS blockade is probably the best therapeutic strategy for patients who are overweight and hypertensive, but because it is associated with a reduced risk of new-onset diabetes, clinical trials are now needed to ascertain (1) if RAS blockade can reduce the risk of gout; (2) if the reduced risk applies to every hyperuricemic subject or only obese subjects; and (3) if the usefulness in reducing urate concentration is limited to certain RAS-blocking drugs or to certain patients of a particular genetic background.

BIANCA MARASINI, MD; MARCO MASSAROTTI, MD, Rheumatology Unit, IRCCS Humanitas Clinical Institute, Via Manzoni 56, Rozzano, Milan 20089, Italy. Address correspondence to Prof. Marasini; E-mail: bianca.marasini@humanitas.it

REFERENCES

1. Sharma AM, Engeli S. The role of renin-angiotensin system blockade in the management of hypertension associated with the cardiometabolic syndrome. *J Cardiometab Syndr* 2006;1:29-35.
2. Iason BE, Martin DE, Boike SC, Jorkasky DK. The effects of eprosartan, an angiotensin II AT1 receptor antagonist, on uric acid excretion in patients with mild to moderate essential hypertension. *J Clin Pharmacol* 1998;38:437-41.
3. Puig JG, Torres R, Ruilope LM. AT1 blockers and uric acid metabolism: are there relevant differences? *J Hypertens Suppl* 2002;20:S29-31.

J Rheumatol 2009;36:12; doi:10.3899/jrheum.090649