

Correction

Chung SJ, Kim JK, Park MC, Park YB, Lee SK. Reactivation of hepatitis B viral infection in inactive HBsAG carriers following anti-tumor necrosis factor- α therapy. *J Rheumatol* 2009;36: 2416-20. Results presented in Table 2 should appear as follows. We regret the error.

Table 2. Serologic profile and viral load of Patient 1.

	Prior to Anti-TNF Therapy	At Week 6	At Week 14	At Week 24
AST, IU/l	21	14	457	28
ALT, IU/l	14	10	1,054	23
Total bilirubin, mg/dl	0.7	0.6	1.5	0.8
Prothrombin time, INR	1.14	Not tested	1.27	1.14
Serologic profile				
HBsAg	Positive	Not tested	Positive	Not tested
Anti-HBs antibody	Negative	Not tested	Negative	Not tested
Anti-HBc antibody IgM	Negative	Not tested	Negative	Not tested
HBeAg	Negative	Not tested	Negative	Not tested
Anti-HBe antibody	Positive	Not tested	Positive	Not tested
Viral load				
HBV PCR, IU/ml	Undetectable	Not performed	3,130,000	Undetectable

HBV: hepatitis B virus. For other abbreviations, see Table 1. PCR: polymerase chain reaction.

doi:10.3899/jrheum.081324C1