Fibromyalgia and Enthesitis in Psoriatic Arthritis. Time to “Characterize” and “Refine” Clinical Definitions

Psoriatic arthritis (PsA) is a complex disease characterized by the presence of polyarthritis, spinal disease, dactylitis, enthesitis, and cutaneous involvement¹. Due to considerable research efforts in PsA in recent years, criteria for classification of PsA² (the CASPAR criteria) have been developed, as well as definitions of clinical features and how to assess them³. In terms of evaluating disease activity, efforts are being made to determine whether composite measures of disease activity and response to therapy can be developed to cover all the domains that characterize PsA⁴.

At present a patient can be classified as having PsA by meeting the CASPAR criteria; nevertheless, in applying the criteria, it is mandatory that the patient must have an inflammatory articular disease (joint, spine, or enthesal). The lack of definition of these entry criteria, crucial for applying the CASPAR criteria, may explain some of the difficulties in clinical research in PsA; for instance, validation of the CASPAR criteria for diagnostic purposes encounters difficulty when criteria are used in a general setting⁵.

Two recent studies^{5,6} testing the sensitivity of the CASPAR criteria in early PsA found different results (99.1% vs 77.3%); apart from the methodological differences in the 2 studies, it is important to emphasize that the clinical diagnosis of PsA relied on clinical expertise as the “gold standard.” Moreover, as many as 43% of patients, in the study with the lowest sensitivity, had a clinical pattern of tenosynovitis, enthesitis, or dactylitis. When tested against noninflammatory diseases⁷ (osteoarthritis and fibromyalgia, FM), in established PsA the sensitivity was 96.1% and the specificity 87.5%.

Recent evidence shows a great divergence between clinical and image findings (total body scintigraphy and ultrasound, US) in early PsA⁸, such that subclinical disease was demonstrated and the pattern of arthritis was reclassified. Again in the same study, clinical diagnosis was the gold standard and as many as three-quarters of patients had oligo-enthesoarthritis.

The presence of “occult” or “painless” disease⁹ is another

important issue when considering a clinical diagnosis of PsA. Several studies have demonstrated, especially in spinal disease¹⁰, the presence of disease without symptoms. Studies using new imaging techniques such as magnetic resonance imaging (MRI) and high-resolution US found subclinical signs of arthritis¹¹ or lower limb enthesal abnormalities¹² in patients with psoriasis. A high prevalence (~70%) of enthesal abnormalities was also found in patients with PsA using power Doppler US (PDUS)^{13,14}.

The lack of a definition of inflammatory articular disease and the recent findings with the newer imaging techniques create problems with the diagnosis of PsA, especially when the spine and entheses are involved¹⁵.

Enthesitis is considered a major feature of PsA and can sometimes be the sole manifestation of early PsA^{8,16}, yet the clinical diagnosis of enthesitis may be difficult. Enthesitis is widely acknowledged as a domain that must be included in the assessment of disease activity in PsA^{3,4}; however, clinical exploration, based on the presence or absence of pain by applying enough pressure to blanch the tip of the finger, is equivocal. FM is a condition¹⁷ characterized by the presence of widespread pain associated with at least 11 of 18 tender points sites on digital palpation; its prevalence in the general population varies between 2% and 4%, and in some cases is associated with rheumatic diseases. The presence of FM in a rheumatic disease can confuse accurate evaluation of disease¹⁸.

Given that enthesitis and FM are characterized by the presence of tender points, it is relevant to be able to distinguish both conditions in a patient with cutaneous psoriasis; using current CASPAR criteria (without a definition of enthesitis) there is a probability of misclassifying some of these patients.

In this issue of *The Journal* Marchesoni and colleagues¹⁹ report the clinical features that can help a physician on clinical grounds to differentiate PsA from FM. A patient has the highest probability of having FM if ≥ 6 somatic manifestations (according to the new diagnostic criteria for FM) and

See Clinical features of PsA versus FM, page 849

≥ 8 tender points are present; thus, if a patient has both criteria, a diagnosis of primary or secondary FM can be made.

Their study, however, raises many questions: the Maastricht Ankylosing Spondylitis Enthesitis Index (MASES), used for measuring enthesitis, showed overlap between the 2 diseases. On the other hand, recent work with PDUS has shown a high prevalence of enthesal abnormalities (~70%) in patients with PsA and a poor correlation between PDUS and MASES^{14,15}. These data emphasize the difficulties in assessing and understanding the clinical relevance of “at least” some enthesal involvement in PsA.

Studies with US and PDUS have demonstrated a high prevalence of enthesal abnormalities in patients with psoriasis without PsA^{12,20}; these findings raise the question of their specificity and clinical meaning. If the lesions found were considered to be specific, many of these patients could be classified as enthesal PsA and would therefore meet the CASPAR criteria.

There is thus a need to clarify several questions concerning enthesal involvement in PsA: (1) to find an accepted and agreed-on method of clinical exploration; (2) to develop a reliable definition of enthesopathy using PDUS, in other words, based on active lesions; (3) to establish correlations between clinical and PDUS findings; (4) to determine the clinical relevance of PDUS findings, not only in psoriasis patients without arthritis, but also in patients with PsA; and (5) to explore the utility of MRI in assessing the enthesal involvement in PsA.

When all these issues have been resolved, a clear definition of enthesal involvement may emerge, and clinicians will be more confident about making decisions, whether for the purpose of diagnosis or for indicating clinical treatments based on objective findings.

JOSÉ LUIS FERNÁNDEZ-SUEIRO, MD,
Rheumatology Division,
Complejo Hospitalario Universitario,
La Coruña, Spain

Supported by a grant from the Fondo de Investigaciones Sanitarias, FIS 08/0789, Ministry of Health, Spain. Address correspondence to Dr. Fernández-Sueiro, Rheumatology Division, Complejo Hospitalario Universitario, C/. Xubias, 84, 15006, La Coruña, Galicia, Spain.
E-mail: L.sueiro@canalejo.org

REFERENCES

- Gladman D, Antoni C, Mease P, Clegg D, Nash P. Psoriatic arthritis: Epidemiology, clinical features, course and outcome. *Ann Rheum Dis* 2005;64 Suppl 2:ii14-17.
- Taylor W, Gladman D, Helliwell P, Marchesoni A, Mease P, Mielants H and the CASPAR study group. Classification criteria for psoriatic arthritis: Development of new criteria from a large international study. *Arthritis Rheum* 2006;54:2665-73.
- Mease P. Measures of psoriatic arthritis. *Arthritis Care Res* 2011;63 Suppl 11:S64-S85.
- Mumtaz A, Gallagher P, Kirby B, Waxman R, Coates L, Veale D, et al. Development of a preliminary composite disease activity index in psoriatic arthritis. *Ann Rheum Dis* 2011;70:272-7.
- Chandran V, Schentag C, Gladman D. Sensitivity of the classification of psoriatic arthritis criteria in early psoriatic arthritis. *Arthritis Rheum* 2007;57:1560-3.
- D'Angelo S, Mennillo G, Cutro M, Leccese P, Nigro A, Padula A. Sensitivity of the classification of psoriatic arthritis criteria in early psoriatic arthritis. *J Rheumatol* 2009;36:368-70.
- Fernández-Sueiro JL, González Díaz de Rábago E, Fernández Lopez JC, Oreiro Villar N, Pinto Tasende, Lema Gontad JM, et al. Validation of CASPAR criteria in a Spanish population: Comparison with potential confounding inflammatory diseases [abstract]. *Ann Rheum Dis* 2010;69 Suppl 3:699.
- Scarpa R, Cuocolo A, Peluso R, Attenu M, Gisonni P, Iervolino S, et al. Early psoriatic arthritis: The clinical spectrum. *J Rheumatol* 2008;35:137-41.
- Palazzi C, Lubrano E, D'Angelo S, Olivieri I. Beyond early diagnosis: Occult psoriatic arthritis. *J Rheumatol* 2010;37:1556-8.
- Queiro R, Belzunegui J, Gonzalez C, De Dios J, Sarasqueta C, Torre J, et al. Clinically asymptomatic axial disease in psoriatic spondyloarthritis. A retrospective study. *Clin Rheumatol* 2002;21:10-3.
- Offidani A, Cellini A, Valeri G, Giovagnoni A. Subclinical joint involvement in psoriasis: Magnetic resonance imaging and x-rays findings. *Acta Dermatol Venereol* 1998;58:2248-56.
- Gisonni P, Tinazzi I, El-Dalati G, Gallo M, Biasi D, Barbar L. Lower limb enthesopathy in patients with psoriasis without clinical signs of arthropathy: A hospital-based case control study. *Ann Rheum Dis* 2008;67:26-30.
- Fernández Sueiro JL, Pinto Tasende JA, González Díaz de Rábago E, Pértiga S, Fernández Lopez JC, et al. Prevalence of enthesitis and concordance of enthesitis exploration with ultrasound versus clinical exploration in psoriatic arthritis [abstract]. *Ann Rheum Dis* 2010;69:580.
- Iagnocco A, Spadaro A, Marchesoni A, Cauli A, De Lucia O, Gabba A, et al. Power Doppler ultrasonographic evaluation of enthesitis in psoriatic arthritis. A multi-center study. *Joint Bone Spine* 2011 Nov 25. [Epub ahead of print]
- Fernandez-Sueiro JL. The challenge and the need of defining axial psoriatic arthritis. *J Rheumatol* 2009;36:2633-4.
- Godfrin B, Zabraniecki L, Lamboley V, Bertrand-Latour F, Sans N, Fournié B. Spondyloarthritis with enthesal pain. A prospective study in 33 patients. *Joint Bone Spine* 2004;71:557-62.
- Wolfe F, Smythe HA, Yunus MB, Bennett RM, Bombardier C, Goldenberg DL, et al. The American College of Rheumatology 1990 criteria for the classification of fibromyalgia. Report of the multicenter criteria committee. *Arthritis Rheum* 1990;33:160-72.
- Almodovar R, Carmona L, Zarco P, Collantes E, González C, Mulero J, et al. Fibromyalgia in patients with ankylosing spondylitis: Prevalence and utility of the measures of activity, function and radiologic damage. *Clin Exp Rheumatol* 2010;28:S33-S39.
- Marchesoni A, Atzeni F, Spadaro A, Lubrano E, Provenzano G, Cauli A, et al. Identification of the clinical features distinguishing psoriatic arthritis and fibromyalgia. *J Rheumatol* 2012;39:849-55.
- Naredo E, Möller I, de Miguel E, Battle-Gualda E, Acebes C, Brito E, et al. High prevalence of ultrasonographic synovitis and enthesopathy in patients with psoriasis without arthritis: A prospective case-control study. *Rheumatology* 2011;50:1838-48.

J Rheumatol 2012;39:677-8; doi:10.3899/jrheum.120095