Use of Hydroxychloroquine in Japan

To the Editor:

Antimalarial agents have been used for the treatment of inflammatory diseases for the last half century, with hydroxychloroquine (HCQ) being approved in the United States in 1955¹. In addition to its common use in systemic lupus erythematosus (SLE), including cutaneous forms, it has also proven useful in treating rheumatoid arthritis (RA)^{2,3,4} and Sjögren's syndrome⁵. In SLE, antimalarial agents have been shown to decrease disease activity and subsequent flares in both nonpregnant⁶ and pregnant patients⁷. Other potential benefits include a decreased risk of infection⁸ and thrombosis⁹. HCQ has been shown to exert a positive effect on overall survival¹⁰. In a large cohort of multiethnic patients with SLE (the LUMINA cohort), HCQ prevented renal and central nervous system disease^{11,12}. Finally, a recent report suggests that maternal use of HCQ may decrease the risk of cardiac manifestations of neonatal lupus¹³.

Despite these benefits and its current use in over 70 countries, chloroquine and HCQ remain unavailable for clinical use for rheumatology patients in Japan. This unavailability stems from a series of lawsuits in the 1970s as a result of chloroquine retinal toxicity, which was first reported by Cambiaggi in 1957¹⁴ and further confirmed by Hobbs, *et al* in 1959¹⁵. Interestingly, chloroquine was widely used in Japan for a variety of clinical indications from 1955 through the early 1970s, including malaria, RA, and SLE, as well as in diseases such as epilepsy and chronic nephritis, in which baseline risk of retinal toxicity was likely higher to begin with¹⁶. The dangers associated with chloroquine use were compounded by the absence of rigorous safety screening protocols, despite the known potential for retinal toxicity. As a result, chloroquine was withdrawn from the Japanese market in 1974.

In the last decade, the clear benefits of antimalarial agents in rheumatological diseases have been increasingly recognized, by the growing cohort of returnee patients already treated with HCQ overseas as well as US-trained rheumatologists returning to Japan for clinical practice. In 2009, an initiative began to promote the study and introduction of HCQ into clinical care in Japan, appreciating the importance of updated safety screening protocols^{17,18}, including earlier detection of retinal toxicity with newer ophthalmologic modalities such as multifocal electroretinogram, spectral domain optical coherence tomography, or fundus autofluorescence. Given its wide use as a standard of care worldwide, there is little reason to support HCQ's continued absence from the market. Unsurprisingly, a recent small study of HCQ in Japanese patients with SLE showed benefit in cutaneous disease, arthritis, and fatigue¹⁹. To this end, 2012 will see the first clinical trial of HCQ for SLE in Japan.

We hope that an understanding of the history of antimalarial agents in Japan — a legacy that precluded the appropriate use of an efficacious therapy — will soon lead to an era of improvement of patient care, survival, and quality of life for patients with SLE in Japan.

MITSUMASA KISHIMOTO, MD, PhD, Division of Allergy and Rheumatology, St. Luke's International Hospital, Tokyo; GAUTAM A. DESHPANDE, MD, Center for Clinical Epidemiology, St. Luke's International Hospital, Tokyo; NAOTO YOKOGAWA, MD, Department of Rheumatology, Tokyo Metropolitan Tama Hospital, Tokyo; JILL P. BUYON, MD, Department of Rheumatology, New York University — Hospital for Joint Diseases, New York, New York, USA; MASATO OKADA, MD, Division of Allergy and Rheumatology, St. Luke's International Hospital, 9-1 Akashi-cho, Chuo-ku, Tokyo 104-8560, Japan. Address correspondence to Dr. M. Kishimoto; E-mail: mkishimo@luke.or.jp

REFERENCES

- Wallace D. The history of antimalarials. *Lupus* 1996;5 Suppl 1:S2-3.
- O'Dell JR, Haire CE, Erikson N, Drymalski W, Palmer W, Eckhoff PJ, et al. Treatment of rheumatoid arthritis with methotrexate alone, sulfasalazine and hydroxychloroquine or a combination of all three medications. *N Engl J Med* 1996;334:1287-91.
- Dixon JS, Pickup ME, Bird HA, Lee MR, Wright V, Downie WW. Biochemical indices of response to hydroxychloroquine and sodium aurothiomalate in rheumatoid arthritis. *Ann Rheum Dis* 1981;40:480-8.
- Kim WU, Seo YI, Park SH, Lee WK, Lee SK, Paek SI, et al. Treatment with cyclosporine switching to hydroxychloroquine in patients with rheumatoid arthritis. *Ann Rheum Dis* 2001;60:514-7.
- Tishler M, Yaron I, Shirazi I, Yaron M. Hydroxychloroquine treatment for primary Sjogren syndrome: Its effect on salivary and serum inflammatory markers. *Ann Rheum Dis* 1999;58:253-6.
- Ruiz-Irastorza G, Ramos-Casals M, Brito-Zeron P, Khamashta MA. Clinical efficacy and side effects of antimalarials in systemic lupus erythematosus: A systematic review. *Ann Rheum Dis* 2010;69:20-8.
- Clowse MEB, Magder L, Witter F, Petri M. Hydroxychloroquine in lupus pregnancy. *Arthritis Rheum* 2006;54:3640-7.
- Ruiz-Irastorza G, Olivares N, Ruiz-Arruza I, Martinez-Berriotxoa A, Egurbide MV, Aguirre C. Predictors of major infections in systemic lupus erythematosus. *Arthritis Res Ther* 2009;11:R109.
- Jung H, Robba R, Su J, Shariati-Sarabi Z, Gladman DD, Urowitz M, et al. The protective effect of antimalarial drugs on thrombovascular events in systemic lupus erythematosus. *Arthritis Rheum* 2010;62:863-8.
- Shinjo SK, Bonfa E, Wojdyla D, Borba EF, Ramirez LA, Scherbarth HR, et al. Antimalarial treatment may have a time-dependent effect on lupus survival: Data from a multinational Latin America inception cohort. *Arthritis Rheum* 2010;62:855-62.
- Fessler BJ, Alarcon GS, McGwin G Jr, Roseman J, Bastian HM, Friedman AW, et al. Systemic lupus erythematosus in three ethnic groups: XVI. Association of hydroxychloroquine use with reduced risk of damage accrual. *Arthritis Rheum* 2005;52:1473-80.
- Pons-Estel GJ, Alarcon GS, McGwin G Jr, Danila MI, Zhang J, Bastian HM, et al. Protective effect of hydroxychloroquine on renal damage in patients with lupus nephritis: LXV. Data from a multiethnic US cohort. *Arthritis Rheum* 2009;61:830-9.
- Izmirly PM, Kim MY, Llanos C, Le PU, Guerra MM, Askanase AD, et al. Evaluation of the risk of anti-SSA/Ro-SSB/La antibody-associated cardiac manifestations of neonatal lupus in fetuses of mothers with systemic lupus erythematosus exposed to hydroxychloroquine. *Ann Rheum Dis* 2010;69:1827-30.
- Cambiaggi A. Unusual ocular lesions in a case of systemic lupus erythematosus. *Arch Ophthalmol* 1957;57:451-7.
- Hobbs HE, Sorsby A, Freedman A. Retinopathy following chloroquine therapy. *Lancet* 1959;2:478-80.
- Marmor MF, Carr RE, Easterbrook M, Farjo AA, Mieler WF; American Academy of Ophthalmology, et al. Recommendations on screening for chloroquine and hydroxychloroquine retinopathy: A report by the American Academy of Ophthalmology. *Ophthalmology* 2002;109:1377-82.
- College of Ophthalmologists. Chloroquine, hydroxychloroquine and the eye. London: College of Ophthalmologists; 1993.
- Marmor MF, Kellner U, Lai TY, Lyons JS, Mieler WF; American Academy of Ophthalmology. Revised recommendations on screening for chloroquine and hydroxychloroquine retinopathy. *Ophthalmology* 2011;118:415-22.
- Yokogawa N, Kato Y, Sugii S, Inada S. Response to hydroxychloroquine in Japanese patients with systemic lupus erythematosus using the Cutaneous Lupus Erythematosus Disease Area and Severity Index (CLASI). *Mod Rheumatol* 2011 Aug 14 [E-pub ahead of print].

J Rheumatol 2012;39:6; doi:10.3899/jrheum.111569