Psoriatic Arthritis Mutilans: Case Series and Literature Review

To the Editor:

Psoriatic arthritis (PsA) is a seronegative spondyloarthropathy that develops in 1% to 39% of patients with psoriasis¹. It may present clinically in different forms, including distal interphalangeal joint arthritis, asymmetrical oligoarthritis, symmetrical polyarthritis, spondylitis, and arthritis mutilans (AM)². Although AM accounts for less than 5% of all PsA, it undeniably represents the most severe and disabling form. Therefore, AM requires timely diagnosis and therapy aiming to prevent or block the irreversible destruction of small joints, mainly involving distal interphalanges of digits and toes. However, data on patients with such arthritis have been published as anecdotal reports or as very small series, so that the available information is sparse.

We describe a series of patients with delayed diagnosis of AM, and a systematic review of studies published in the last decade.

Case 1. A 77-year-old woman, with a 37-year history of psoriasis, was diagnosed with PsA at age 55 years when a symmetric polyarthritis involving both hands and feet occurred. She received longterm therapy with different nonsteroidal antiinflammatory drugs (NSAID) without significant improvement of arthritis, because she was intolerant to disease-modifying antirheumatic drugs (DMARD). She refused biological therapy. When significant deformity in phalanges of the left hand developed, she was referred to our ambulatory service. Biochemical tests revealed erythrocyte sedimentation rate (ESR) 50 mm/h, C-reactive protein (CRP) 5.84 g/l; clinical assessment showed pain score 70 on 100-cm visual analog scale (VAS), Health Assessment Questionnaire (HAQ) score 1.75, swollen joint count (SJC) = 6, tender joint count (TJC) = 5, and DJC = 5. Figure 1A illustrates AM involving the hands. We suggested biologic

Figure 1. Destructive involvement of distal interphalangeal (DIP) joints and subluxation of the metacarpophalangeal joints (A). Mutilation of fourth and fifth digits of the right hand (B). Destructive involvement of the DIP joints, with telescoping of second finger of the right hand and fourth finger of the left hand (C). Shortening of third finger of right hand and second finger of left hand, and subluxation of DIP joints; onycholysis is also evident (D).

therapy with an anti-tumor necrosis factor- α (anti-TNF- α) agent, but she declined treatment.

Case 2. A 74-year-old woman developed psoriasis at the age of 20 years and arthritis when she was 45 years old, with both axial and peripheral involvement. Therapy with indomethacin 50 mg daily achieved adequate control of arthritis-associated pain, so that no other therapies were performed. However, she was referred to our center with the following measures: ESR 43 mm/h, CRP 2.39 g/l, VAS for pain 50, HAQ score 1.25, SJC = 5, TJC = 3, and DJC = 4. Figure 1B shows the patient's hands with AM. She refused treatment with a biologic therapy despite information on the potential benefits for cutaneous and rheumatic manifestations.

Case 3. A 62-year-old woman with a 25-year history of psoriasis was diagnosed with PsA at age 50 years. She was irregularly treated with DMARD and with some cycles of etanercept, which she discontinued because of an "intolerance." Consequently, she remained without any therapy for several months. At clinical assessment, ESR was 25 mm/h, CRP 1 g/l, VAS for pain = 30, HAQ score = 0.75, SJC = 2, TJC = 2, and DJC = 3. Figure 1C shows her hands. She declined reintroduction of biologic therapy.

Case 4. A 70-year-old man with psoriasis at age 40 years developed arthritis involving the hands when he was 63 years old. He had previously received NSAID with nonsignificant control of pain, but no DMARD therapy. Referred to our clinic, he provided a radiograph with features of AM of the hands (Figure 1D). We undertook biochemical tests to start an appropriate therapy, but the patient was lost to followup.

Literature review. A literature search was performed in PubMed with MeSH terms "psoriatic arthritis mutilans" and "opera-glass hand" for the period January 2002 to December 2012. Only studies in which clinical data of patients with AM were specifically reported were extracted. Full articles of all relevant studies were retrieved, and manual searches of reference lists from these articles were performed, identifying 14 suitable studies (Figure 2)^{3,4,5,6,7,8,9,10,11,12,13,14,15,16}. Including our series, a total of 120 patients with AM are described (Table 1). It was impossible to extrapolate data for 37 other AM cases reported in 5 studies because they were cumulatively

described among other patients with PsA^{17,18,19,20,21}. There was a weak prevalence of the male sex, with male/female ratio being 1.29. Duration of psoriasis and PsA before diagnosis of AM ranged from 10 to 52 years and 6 to 35 years, respectively; acute onset of AM (3 months) was reported in 1 patient⁹. In 4 cases, psoriatic AM was associated with Charcot-like joint disease⁶, while 1 patient presented with an axial bone proliferation causing cervical myelopathy¹². In the largest series³, the AM was significantly associated with arthritis duration, involvement of distal phalanges, and reduced functional capacity. The prevalence (55%) of psoriatic nail involvement in AM did not differ from that of other psoriatic patients³. A study found that bone edema, erosion, and proliferation on magnetic resonance imaging (MRI) were significantly more severe in the patients with AM than in controls⁴.

Therapy with 1 or more DMARD was performed before AM diagnosis in all but 1 patient, although such therapy was carried out irregularly in the majority of cases. Following AM diagnosis, biologic therapy with etanercept was performed in 6 cases (Table 1). The functional capacity improved significantly in 3 cases, with recovery of some movements allowing patients to make a fist and to walk easily⁷. Of note, therapy provided a stable bone disease without onset of new mutilation, as documented by radiographic features at 2-year followup in 2 other cases¹⁴. Unfortunately, no followup is available for the remaining AM patient treated with etanercept¹³.

AM involves fewer than 5% of patients with PsA¹, the disease being particularly infrequent in Asian countries. Recent studies found an AM prevalence ranging from 1.5% to 4.9%^{18,20,21}. AM represents an aggressive form of arthritis characterized by marked erosion of small joints of hands and feet, progressing to irreversible joint disruption. The resulting finger and toe mutilations severely impair patients' functional capacity. Much more effort is required to diagnose those patients whose PsA is progressing toward mutilation in an earlier stage. Development of AM has been associated with a longer arthritis duration, involvement of distal phalanges, and impaired functional capacity³, suggesting these patients should undergo tight disease control and therapy. Radiological assessment of joints may be useful for clinical management of PsA. A study found that as

Figure 2. Identification of studies of arthritis mutilans (AM).

Table 1. Clinical characteristics of patients with arthritis mutilans (AM) reported in the last decade.

Country	No. Cases	M/F	Age, yrs	Psoriasis Duration, yrs	Arthritis Duration, yrs	Site of AM	DMARD Therapy	Biologic Therapy Following AM Diagnosis
Spain ³	24	12/12	56.3	20.8	17.7	Hands and feet	S	No
New Zealand ⁴	11	8/3	52	22	12	NA	M; L; Su; S	No
Norway ⁵	6	5/1	NA	NA	NA	Hands	M; S	No
Italy (current series)	4	1/3	71	38.7	21	Hands	M; Su; E	No
Columbia ⁶	4	2/2	57.8	13	6	Hands and feet	M	No
USA ⁷	3	2/1	39.7	17	15.7	Hands and feet	M, H, L, A	E
Portugal ⁸	2	1/1	54.5	NA	NA	Hands and feet	NA	NA
Ireland ⁹	1	0/1	34	21	0.3	Feet	None	No
Brazil ¹⁰	1	1/0	55	25	18	Hands	M	No
Croatia ¹¹	1	1/0	33	10	10	Hands and feet	S; Su; M; H	No
New Zealand ¹²	1	1/0	65	52	35	Hands and feet	M	No
Spain ¹³	1	1/0	62	10	10	Hands and feet	L	E
Portugal ¹⁴	1	1/0	39	20	20	Hands	M	E
USA ¹⁵	1	1/0	55	28	28	Hands and feet	M	E
Scandinavia ¹⁶	59	30/29	NA	NA	30	Hands and feet	NA	NA

DMARD: disease-modifying antirheumatic drug; A: azathioprine; H: hydroxychloroquine; L: leflunomide; M: methotrexate; S: steroid; Su: sulfasalazine; E: etanercept; NA: not available.

many as 23% of patients with PsA have damage at the initial visit that was not detectable at clinical examination²². Similarly, it has been found that joint erosions are detectable at early radiographic assessment in nearly 50% of patients, i.e., within 2 years of arthritis²³. Another study found that bone edema, erosions, and proliferation assessed with MRI scans were significantly more severe in patients with psoriatic AM than in those with other forms of PsA, suggesting that bone edema visible on MRI may be a pre-erosive feature of arthritis⁴. Thus, it is possible to diagnose early joint alterations in psoriatic patients in advance of their clinical manifestation.

An early treatment may prevent joint damage and irreversible finger destruction. A trial showed that structural joint damage in patients with PsA was significantly prevented by infliximab therapy, and that even a 6-month delay in starting such therapy was associated with a lower therapeutic gain¹⁸. A dramatic repair of joint damage has been observed in a patient with PsA following 2 years of etanercept therapy, with fill-in and disappearance of typical destructive erosions in both fingers and toes²⁴. Similar observations were described by others in clinical practice²⁵.

Thus, biologic therapy may reduce radiologic progression and possibly induce regression of erosive damage in such patients. Such therapy could prevent or significantly delay the development of AM, but data for this remain scarce. Our review included 6 cases in whom the etanercept therapy was performed after AM diagnosis, that is, when both joint and bone alterations were already irreversible. However, the absence of progression at 2-year followup would suggest that such therapy may be beneficial even in these cases. In 1 patient from our series, etanercept was administered before onset of AM, but only a few irregular therapeutic cycles were administered because of patient refusal, so no reliable conclusions can be drawn from that case.

Different guidelines suggest that psoriatic patients with peripheral arthritis should receive anti-TNF- α therapy (grade A) or DMARD (grade B) following failure of NSAID²⁶. Unfortunately, all our patients declined consent for such therapy. In our practice, such diffidence is much less evident in patients with other rheumatic diseases or inflammatory bowel diseases. Speculatively, it cannot be excluded that a general reluctance of these patients to accept effective drugs, including biologic agents, may be a factor for the development of AM.

AM is an infrequent but highly disabling form of PsA. Based on our findings, therapy with DMARD seems unable to prevent its development. A potential role for biologic therapy is expected, and prospective studies are urged.

VINCENZO BRUZZESE, MD; CINZIA MARRESE, MD; LORENZO RIDOLA, MD; ANGELO ZULLO, MD, Departments of Internal Medicine, Rheumatology, and Gastroenterology, Nuovo Regina Margherita Hospital, Via E. Morosini 30, 00153 Rome, Italy.

Address correspondence to Dr. Bruzzese; E-mail: vinbruzzese@tiscali.it

REFERENCES

- Cantini F, Piccoli L, Nannini C, Kaloudi O, Bretoni M, Cassarà E. Psoriatic arthritis: a systematic review. *Int J Rheum Dis* 2010;13:300-17.
- Taylor W, Gladman D, Helliwell P, Marchesoni A, Mease P, Mielants H. The CASPAR Study Group. Classification criteria for psoriatic arthritis: development of new criteria from a large international study. *Arthritis Rheum* 2006;54:2665-73.
- Rodríguez-Moreno J, Bonet M, Del Blanco-Barnusell J, Castaño C, Clavaguera T, Mateo-Soria L, et al. Mutilating/resorptive arthritis. A study of 24 patients in a series of 360 patients with psoriatic arthritis. *Reumatol Clin* 2013;9:38-41.
- Tan YM, Østergaard M, Doyle A, Dalbeth N, Lobo M, Reeves Q, et al. MRI bone oedema scores are higher in the arthritis mutilans form of psoriatic arthritis and correlate with high radiographic scores for joint damage. *Arthritis Res Ther* 2009;11:1-9.
- Nossent JC, Gran JT. Epidemiological and clinical characteristics of psoriatic arthritis in northern Norway. *Scand J Rheumatol* 2009;38:251-5.
- Candia L, Cuellar ML, Marlowe SM, Marquez J, Iglesias A, Espinoza LR. Charcot-like arthropathy: a newly-recognized subset of psoriatic arthritis. *Clin Exp Rheumatol* 2006;24:172-5.
- Pomerantz RG, Mody E, Husni ME, Qureshi AA. Follow-up of psoriasis arthritis mutilans patients treated with anti-TNF- α therapy. *J Drugs Dermatol* 2009;8:406-12.
- Ribeiro A, Costa J, Bogas M, Costa L, Araújo D. Forma mutilante de artrite psoriática [Mutilans psoriatic arthritis]. *Acta Reumatol Port* 2009;34:290-1.
- Bell L, Murphy CL, Wynne B, Cunanne G. Acute presentation of arthritis mutilans. *J Rheumatol* 2011;38:174-5.
- Appel da Silva F, Appel da Silva MC, Romagna ES. Clinical images: psoriatic arthritis mutilans. *Arthritis Rheum* 2010;62:2159.
- Vrogić P, Pašić A, Podobnik-Takac T. Psoriasis vulgaris and arthritis psoriatica gravis mutilans. *Acta Dermatovenol Croat*

- 2003;11:22-8.
12. Ly J, Pinto C, Doyle A, Dalbeth N, McQueen FM. Axial bone proliferation causing cervical myelopathy in the mutilans form of psoriatic arthritis despite peripheral bone erosion. *Ann Rheum Dis* 2009;68:443-4.
13. González-Nieto JA, López-Montes L, Gallego-García F, Tugues-Roure JM. Psoriatic arthritis mutilans. *Rev Clin Esp* 2012;212:e87.
14. Peixoto D, Teixeira F, Bogas M, Costa J, Araújo D. Biologic therapy in psoriatic arthritis mutilans. *Acta Reumatol Port* 2011;36:189-90.
15. Gaffar M. Arthritis mutilans in a patient with psoriasis. *Hospital Physician* 2002;10:46-50.
16. Gudbjornsson B, Ejstrup L, Gran JT, Iversen L, Lindqvist U, Paimela L, et al. Psoriatic arthritis mutilans (PAM) in the Nordic countries: demographics and disease status. The Nordic PAM study. *Scand J Rheumatol* 2013 Mar 21. [E-pub ahead of print]
17. Yamamoto T, Yokozeki H, Nishioka K. Clinical analysis of 21 patients with psoriasis arthropathy. *J Dermatol* 2005;32:84-90.
18. van der Heijde D, Kavanaugh A, Gladman D, Antoni C, Krueger GG, Guzzo C, et al. Infliximab inhibits progression of radiographic damage in patients with active psoriatic arthritis through one year of treatment. *Arthritis Rheum* 2007;56:2698-707.
19. Soy M, Karaca N, Umit EU, Bes C, Piskin S. Joint and nail involvement in Turkish patients with psoriatic arthritis. *Rheumatol Int* 2008;29:223-5.
20. Reich K, Kruger K, Mossner R, Augustin M. Epidemiology and clinical pattern of psoriatic arthritis in Germany: A prospective interdisciplinary epidemiological study of 1511 patients with plaque-type psoriasis. *Br J Dermatol* 2009;160:1040-7.
21. Helliwell PS, Porter G, Taylor WJ, for The CASPAR Study Group. Polyarticular psoriatic arthritis is more like oligoarticular psoriatic arthritis, than rheumatoid arthritis. *Ann Rheum Dis* 2007;66:113-7.
22. Khan M, Schentag C, Gladman DD. Clinical and radiological changes during psoriatic arthritis disease progression. *J Rheumatol* 2003;30:1022-6.
23. Kane D, Pathare S. Early psoriatic arthritis. *Rheum Dis Clin North Am* 2005;31:641-57.
24. Eder L, Chandran V, Gladman DD. Repair of radiographic joint damage following treatment with etanercept in psoriatic arthritis is demonstrable by 3 radiographic methods. *J Rheumatol* 2011;38:1065-70.
25. Garcia-Valladares I, Cuchacovich R, Espinoza LR. Psoriatic arthritis: Radiographic joint repair following etanercept therapy [letter]. *J Rheumatol* 2012;39:185.
26. Ritchlin CT, Kavanaugh A, Gladman DD, Mease PJ, Helliwell P, Boehncke WH, et al. Treatment recommendation for psoriatic arthritis. *Ann Rheum Dis* 2009;68:1387-94.

J Rheumatol 2013;40:7; doi:10.3899/jrheum.130093