

Androgens and Integrins in Salivary Glands in Sjögren's Syndrome

PAULIINA POROLA, MIKAEL LAINE, ISMO VIRTANEN, RAIMO PÖLLÄNEN, BEATA D. PRZYBYLA, and YRJÖ T. KONTTINEN

ABSTRACT. Objective. Laminin α 1-chain normally induces intercalated duct progenitors to differentiate to acinar cells through integrin (INT) α 1 β 1 and α 2 β 1 receptors. Maintenance of acinar cells is impaired in Sjögren's syndrome (SS), which is also characterized by low levels of serum and salivary androgens. We hypothesized that androgens normally support salivary gland remodeling by upregulating either laminin α 1 chain or its cellular α 1 or α 2 INT subunit-containing receptors.

Methods. Intercalated duct and acinar human salivary gland (HSG) cells and labial salivary gland (LSG) biopsies from healthy controls and patients with SS were cultured without or with sex steroids. Laminin α 1 chain and INT α 1 and α 2 subunits were studied using quantitative reverse-transcription real-time polymerase chain reaction and INT α 1 and α 2 subunits using immunofluorescence staining.

Results. INT α 1-subunit and α 2-subunit messenger RNA (mRNA) levels were increased in intercalated duct and acinar cells by DHEA and testosterone. In contrast, laminin α 1-chain mRNA levels were not affected. The upregulating effect of DHEA on INT subunits was also seen at the protein level. DHEA also increased mRNA levels of both INT subunits in healthy but not SS LSG.

Conclusion. Androgens increased INT α 1 and α 2 subunits in tubuloepithelial cells and in healthy LSG, but in SS salivary glands this androgen regulation was defective, which is likely to contribute to defective outside-in signaling, acinar atrophy, and ductal cell hyperplasia. (J Rheumatol First Release May 1 2010; doi:10.3899/jrheum.091354)

Key Indexing Terms:

SJÖGREN'S SYNDROME ANDROGENS INTEGRINS LAMININ SALIVARY GLANDS

Sjögren's syndrome (SS)¹ is a complex women-dominant autoimmune disorder affecting salivary and lacrimal glands. The acinar cells of the severely affected glands become atrophic, but the factors responsible for their impaired main-

tenance are unclear. Normal salivary glands undergo continuous remodeling. To be able to do so, acinar cells undergo apoptosis, detach, and are carried away with salivary flow, leaving a space in the acinar basement membrane. This loss of acinar cells can be compensated by intercalated duct cells. These morphologically simple, undifferentiated progenitor cells undergo asymmetric divisions so that the progenitor cell pool is maintained at the same time that one of the daughter cells migrates to the acinus and receives site-specific signals for acinar differentiation to replace the lost acinar cell^{2,3}. Cloned intercalated duct cells have been shown to possess such a differentiation potential^{4,5}. In this acinar differentiation, process signaling between extracellular matrix (ECM) and cells is essential. Signaling between laminin α 1 chain and its cellular β 1 subunit containing integrin (INT) receptors is especially important⁶.

It has recently been shown that the major acinar cell INT able to mediate binding to acinar basement membrane-confined laminin-111 (LM-111) are α 1 β 1 and α 2 β 1 integrins, which both contain the ligand-binding I domain^{7,8}. Thus it seems that in addition to LM-332-INT α 6 β 4 interaction^{7,9}, one of the central interactions necessary for the maintenance of acini occurs between the laminin α 1-chain containing LM-111 in the acinar basement membrane and its α 1 β 1 and α 2 β 1 integrin receptors on the tubuloacinar epithelial cells. Interestingly, SS is characterized by low levels of acinar

From the Department of Medicine, Helsinki University Central Hospital; Department of Anatomy, University of Helsinki, Helsinki; ORTON Orthopaedic Hospital of the ORTON Foundation, Helsinki; COXA Hospital for Joint Replacement, Tampere, Finland; and the Institute of Aging, University of Arkansas for Medical Sciences, Little Rock, Arkansas, USA.

Supported by the Orion-Farmos Research Foundation, the Finnish Medical Society Duodecim, the Finnish Dental Society Apollonia, the Academy of Finland, EVO grants, the Sigrid Jusélius Foundation, the Finnish Cultural Foundation, the Emil Aaltonen Foundation, the Maire Lisko Foundation, the National PhD Graduate School of Musculoskeletal Disorders and Biomaterials, the Biomedicum Helsinki Foundation, and Finska Läkaresällskapet.

P. Porola, MSc, Department of Medicine, Helsinki University Central Hospital, Department of Anatomy, University of Helsinki; M. Laine, MD, DDS; I. Virtanen, MD, PhD, Department of Anatomy, University of Helsinki; R. Pöllänen, PhD, Department of Medicine, Helsinki University Central Hospital; B.D. Przybyla, PhD, Institute of Aging, University of Arkansas for Medical Sciences; Y.T. Konttinen, MD, PhD, Department of Medicine, Helsinki University Central Hospital, ORTON Orthopaedic Hospital of the ORTON Foundation, COXA Hospital for Joint Replacement.

Address correspondence to Prof. Y.T. Konttinen, Department of Medicine, Biomedicum 1, Helsinki, Helsinki University Central Hospital, PO Box 700, FIN-00029 HUS, Helsinki, Finland. E-mail: yrjo.konttinen@helsinki.fi
Accepted for publication January 12, 2010.

Personal non-commercial use only. The Journal of Rheumatology Copyright © 2010. All rights reserved.

compartment-specific $\alpha 1$ laminin¹⁰ and low levels of its integrin $\alpha 1\beta 1$ and $\alpha 2\beta 1$ receptors on acinar cells⁸. Additionally, in the early stages of SS, the ECM is actively remodeled, possibly as an attempt to diminish the acinar cell loss by enhancing the interaction between those cells and the acinar basement membrane⁹. In agreement with the intercalated duct progenitor cell hypothesis, advanced SS is characterized by diminished ECM remodeling, loss of acinar cells, and a reciprocal ductal cell hyperplasia^{9,11-13}.

Patients with SS have low concentrations of the prohormones DHEA and androstenedione, and the active androgens testosterone and dihydrotestosterone (DHT) in serum and saliva^{14,15}. Humans and other primates are unique in being able to synthesize active sex steroids both in gonads and locally in peripheral tissues. This is possible because adrenal glands secrete high amounts of prohormones, which are further processed into active androgens or estrogens by intracrine steroidogenic enzymes, according to local tissue needs¹⁶. Thus, concentrations of sex steroids in peripheral tissues can differ from the corresponding systemic levels. It has recently been shown that salivary glands possess organized enzymatic functions needed for local production of sex steroids for intracrine use. But in SS salivary glands the androgen-processing enzymes are deranged further, deteriorating the local androgen depletion¹⁷.

Signaling between extracellular matrix and cells through acinar-specific laminin $\alpha 1$ (or actually LM-111)-INT $\alpha 1\beta 1/\alpha 2\beta 1$ interactions is essential for salivary glands⁶. Results from our studies^{8,10} led to the hypothesis that these molecules, laminin $\alpha 1$ and/or its INT receptors $\alpha 1$ and $\alpha 2$, which are expressed in low numbers in androgen-deficient salivary glands in SS^{8,10}, are normally upregulated by sex steroids. In other words, androgens would help to maintain acini, the secretory end pieces, in healthy salivary glands, but fail to do this in SS. In our study we tested our hypothesis at the cellular level, using a well established *in vitro* human submandibular gland cell model, the human salivary gland (HSG) cell line¹⁸, and tried to extend this to the tissue level with labial salivary gland (LSG) obtained from healthy controls and patients with SS.

MATERIALS AND METHODS

Cell culture and stimulation. HSG cells¹⁸ were cultured without or with Matrigel (BD Biosciences, San Jose, CA, USA) in DMEM/F-12 Nut Mix medium (Gibco BRL, Grand Island, NY, USA) supplemented with 10% fetal calf serum, 2 mM glutamine, 100 U/ml penicillin, and 100 μ g/ml streptomycin at 37°C. Without Matrigel these cells maintain their intercalated duct phenotype, but on LM-111-containing Matrigel they differentiate to acinar cells¹⁹. The HSG cells, with either ductal or acinar cell phenotype, were cultured in a fetal calf serum (FCS)-free medium without or with 1, 10, and 100 μ M DHEA (Sigma-Aldrich, St. Louis, MO, USA), 1 μ M testosterone (Sigma-Aldrich), 1 and 10 nM and 1 μ M DHT (Sigma-Aldrich), or 100 nM 17 β -estradiol (Sigma-Aldrich) for 48 or 72 hours.

Tissue culture and stimulation. LSG biopsies were obtained from 4 healthy controls who were treated for mucocoele (3 women, 1 man, ages 17, 17, 21, and 57 yrs) and 3 patients with SS (all women, ages 44, 58, and 65 yrs).

Glands were split into pieces of about 2 mm³ and cultured overnight in Dulbecco modified Eagle (DME)-Ham's F-12 medium (Gibco) containing 10% FCS, L-glutamine, 10 \times antibiotics (1000 U/ml penicillin and 1000 μ g/ml streptomycin), and 2.5 μ g/ml amphotericin B (Gibco). The next day the media were replaced with basal DME-Ham's F-12 medium containing 10% FCS, L-glutamine, 1 \times antibiotics, and 2.5 μ g/ml amphotericin B, and after 1 more day with serum-free media for stimulation with 100 μ M DHEA for 72 hours.

RNA isolation and cDNA synthesis. Cells grown on Matrigel were detached from the Matrigel with dispase (BD Biosciences). Total RNA was isolated using the Trizol protocol (Invitrogen, San Diego, CA, USA) or with RNeasy Mini kit (Qiagen, Hilden, Germany) and messenger RNA (mRNA) was purified from total RNA using the Dynabeads mRNA Purification Kit (Dyna, Oslo, Norway). From tissue samples total RNA was isolated with High Pure RNA Tissue kit (Roche, Basel, Switzerland). cDNA was synthesized using SuperScript First Strand cDNA Synthesis System for quantitative reverse-transcription realtime polymerase chain reaction (RT-PCR; Invitrogen).

Quantitative RT-PCR. Quantitative RT-PCR was done using a Light CyclerTM PCR machine (Roche Molecular Biochemicals, Mannheim, Germany), SYBR Green I label, and purpose-designed primers (laminin $\alpha 1$ forward 5'-GCT CTG TGA CTG CAA ACC AA-3' and reverse 5'-TTT CTG GGT CGC AGG TAT TC-3'; integrin $\alpha 1$ forward 5'-TCC ACC GAA GAG GTA CTT GTT GCA-3' and reverse 5'-CCA AGC ATG ACC CAG TCC TGT GA-3'; and integrin $\alpha 2$ forward 5'-GGT GAG GAT GGA CTT TGC AT-3' and reverse 5'-GGC TTG GAA ACT GAG AGA CG-3'). Porphobilinogen deaminase (PBGD) and β -actin housekeeping genes were used for standardization of the results (in cell and tissue samples, respectively) by using 5'-ACA TGC CCT GGA GAA GAA TG -3' and 5'-AGA TGC GGG AAC TTT CTC TG-3' and 5'-TCA CCC ACA CTG TGC CCA TCT ACG A-3' and 5'-CAG CGG AAC CGC TCA TTG CCA ATG G-3' primers for PBGD and β -actin, respectively.

Indirect immunofluorescence of integrin receptors. For indirect immunofluorescence staining of the integrin-type laminin receptors, which have been described in human LSG⁸, the following monoclonal antibodies were used: TS2/7 against integrin $\alpha 1$ -subunit²⁰ and 10G11 against integrin $\alpha 2$ -subunit²¹. Cells were washed in 10 mM phosphate-buffered saline, 150 mM saline, pH 7.4, containing 0.1% Triton X-100. After incubation with primary antibodies, cells were washed in Triton X-100 containing phosphate buffer and the bound antibodies were visualized using FITC-conjugated secondary goat anti-mouse IgG antibody (Alexa Fluoro 488, Molecular Probes, Eugene, OR, USA). Propidium iodide diluted 1:1000 in phosphate buffer was used for nuclear counterstaining. After washes in Triton X-100 containing phosphate buffer, the specimens were embedded in fluorescent mounting medium (Dako, Glostrup, Denmark) and examined under an Olympus AX70 (Tokyo, Japan) microscope coupled with a CCD camera (Olympus DP71). Two different types of filters were used in order to show the FITC-positive staining with or without propidium iodide. Control immunostainings were performed using irrelevant primary monoclonal antibodies at the same concentration as and instead of the primary specific antibodies and by using conjugated secondary antibodies alone.

Statistical analysis. Statistical analysis was done using SPSS for Windows V.16.0 software (SPSS Inc., Chicago, IL, USA). Integrin $\alpha 1$ and $\alpha 2$ mRNA levels without and with hormone treatments were compared using the Mann-Whitney U test and the overall effects of stimulations with different androgen concentrations studied with the Kruskal-Wallis test. The level of significance throughout the study was set at 0.05.

RESULTS

Effect of Matrigel on expression of laminin $\alpha 1$ chain and integrin $\alpha 1$ and $\alpha 2$ -subunits in salivary gland cells. Culture on Matrigel causes differentiation of HSG cells of intercalated duct phenotype to acinar cells, which increased the

mRNA levels of laminin $\alpha 1$ constantly over 72 hours [copy numbers per 10^5 PBGD, 780 ± 668 ($n = 3$), 1262 ± 505 ($n = 3$), and 3233 ± 3512 ($n = 4$) for 24-hour, 48-hour, and 72-hour timepoints, respectively].

Also, levels of INT $\alpha 1$ ($p = 0.013$; Figure 1A) but not those of $\alpha 2$ -subunit mRNA ($p = 0.333$; Figure 1B) increased upon culture on Matrigel. The corresponding phenomenon was seen at the protein level in INT $\alpha 1$ -subunits (Figure 2).

Effect of sex steroids on expression of laminin $\alpha 1$ and INT $\alpha 1$ and $\alpha 2$ -subunits in salivary gland cells. Culture in the presence of sex steroids did not affect laminin $\alpha 1$ -chain mRNA levels in ductal or acinar HSG cells or LSG biopsies (data not shown).

Expression of the mRNA levels of INT $\alpha 1$ and $\alpha 2$ -subunits in ductal (cultured without Matrigel) and acinar (cultured on Matrigel) HSG cells was significantly increased after 72-hour stimulation by DHEA.




Figure 1. Effect of 72-hour culture on Matrigel on mRNA levels of integrin (INT) $\alpha 1$ (A) and $\alpha 2$ (B) chains in human submandibular gland HSG cells. Levels of INT $\alpha 1$ -chain mRNA (scale to 200,000) were much higher in ductal (without Matrigel) and acinar (with Matrigel) cells than levels of INT $\alpha 2$ -chains (scale to 20,000). *Statistically significant differences, $p < 0.05$.

In intercalated duct cells, 1, 10, and 100 μM DHEA increased INT $\alpha 1$ -subunits by 37% ($p = 0.312$), 46% ($p = 0.016$), and 449% ($p = 0.004$), respectively. The corresponding increases in INT $\alpha 2$ -subunits were 54% (0.153), 640% ($p = 0.003$), and 1100% ($p = 0.007$) (Figure 3).

In acinar cells 1, 10, and 100 μM DHEA induced changes of INT $\alpha 1$ -subunits by 90% (0.053), 102% ($p = 0.009$), and 113% ($p = 0.655$), respectively, and of INT $\alpha 2$ -subunits by 39% ($p = 0.551$), 473% ($p = 0.046$), and 3% ($p = 0.594$), respectively (Figure 3).

Stimulation with 1 μM testosterone increased INT $\alpha 1$ and $\alpha 2$ -subunits by 79% ($p = 0.009$) and 120% ($p = 0.020$), respectively, in intercalated duct cells. In acinar cells, testosterone changed INT $\alpha 1$ and $\alpha 2$ -subunits by 72% ($p = 0.026$) and 96% ($p = 0.086$), respectively (Figure 4).

Stimulation with 1 nM, 10 nM, or 1 μM DHT showed only a tendency by changing the expression of INT $\alpha 1$ -subunit mRNA levels/ 10^5 PBGD in ductal cells by 33%, 31%, and 55% ($66,620 \pm 35,175$, $n = 11$, vs $88,619 \pm 20,818$, $n = 3$; $87,172 \pm 19,015$, $n = 3$; and $102,971 \pm 7663$, $n = 3$, respectively). The corresponding changes of INT $\alpha 2$ -subunit were 41%, 3%, and 62% (5008 ± 2467 , $n = 8$, vs 7064 ± 1231 , $n = 3$; 5170 ± 470 , $n = 3$; and 8137 ± 1298 , $n = 3$). In acinar cells the changes in INT $\alpha 1$ -subunit were 16%, 94%, and 70% ($120,926 \pm 59,627$, $n = 8$, vs $140,696 \pm 99,367$, $n = 2$; $234,734 \pm 122,191$, $n = 3$; and $205,284 \pm 95,411$, $n = 3$) and for INT $\alpha 2$ -subunit 135%, 89%, and 90% (9422 ± 7167 , $n = 15$ vs $22,135 \pm 19,277$, $n = 2$; $17,835 \pm 11,051$, $n = 3$; and $17,932 \pm 6692$, $n = 3$). Despite consistent upregulation at all concentrations used, these increases were not statistically significant. Estradiol did not have significant effects on INT subunits (data not shown).

At the protein level, both ductal and acinar HSG cells cultured 48 and 72 hours in the presence of DHEA contained more cells intensively positive for INT $\alpha 1$ -subunit, compared to the cells cultured in the absence of DHEA (shown for acinar cells in Figure 2). According to results from qRT-PCR, INT $\alpha 2$ -subunit immunostaining was much weaker than that of the INT $\alpha 1$ -subunit (not shown). Staining controls were negative, confirming the specificity of the staining with the monoclonal antibodies used in our study.

Effect of DHEA stimulation on laminin $\alpha 1$ chain and INT $\alpha 1$ and $\alpha 2$ -subunits in salivary gland biopsies. Extending these observations to tissue samples, we observed that in agreement with the cell stimulations, the expression of laminin $\alpha 1$ -chain in LSG of healthy controls and patients with SS was unaffected by stimulation with androgens (data not shown). The expression of INT $\alpha 1$ and $\alpha 2$ -subunits was increased by DHEA stimulation in LSG from healthy controls, but in LSG from patients with SS such an increase was not seen (Figure 5). The effect of DHEA on the expression of INT $\alpha 1$ and $\alpha 2$ subunits showed a slight tendency to diminish with increasing age in both healthy and SS labial salivary glands (data not shown).




Figure 2. Immunofluorescence staining of integrin (INT) α 1-subunit in human submandibular gland HSG cells cultured on Matrigel without or with DHEA for 48 and 72 hours (B and D, respectively). Compared to non-DHEA-stimulated cells (A, 48 h; C, 72 h), a clear increase of INT α 1-subunit staining intensity is seen upon DHEA costimulation (DHEA and Matrigel). Some increase in the intensity of INT α 1-subunit staining is also seen in nonstimulated cells (A compared to C). Original magnification 200 \times .




Figure 3. Effect of DHEA stimulations on the expression of integrin (INT) α 1 (A and B) and α 2 (C and D) subunits in ductal and acinar human submandibular gland HSG cells (cultured without or with Matrigel, respectively). Stimulations were done using 1, 10, or 100 μ M DHEA for 72 hours (D1, D10, D100, respectively; D0: nonstimulated samples). Overall significance (Kruskal-Wallis test) is shown in the heading for each panel. *p < 0.05, **p < 0.01, individual statistically significant differences with nonstimulated samples.




Figure 4. Effects of testosterone stimulations on the expression of integrin (INT) $\alpha 1$ (A and B) and $\alpha 2$ (C and D) subunits in ductal and acinar human submandibular gland HSG cell line (cultured without or with Matrigel, respectively). Stimulations were done using $1 \mu\text{M}$ testosterone for 72 hours. * $p < 0.05$, ** $p < 0.01$.

DISCUSSION

Sjögren's syndrome is characterized by low levels of androgens both at the endocrine level in the systemic circulation and at the intracrine level locally in salivary glands^{14,15}. Earlier findings also suggested deficiencies in acinar-specific laminin molecules signaling from basement membrane to tubuloacinar epithelial cells, namely LM-111 signaling through the integrin receptors $\alpha 1\beta 1$ and $\alpha 2\beta 1$, in salivary glands in SS^{8,10}. Our hypothesis was that these characteristic features are interconnected. Results confirm our hypothesis and show that sex steroids increase the expression of $\alpha 1\beta 1$ and $\alpha 2\beta 1$ INT receptors for LM-111 in ductal and acinar cells. Interestingly, of the androgens used for stimulations, DHEA proved to be the most effective upregulator of both INT subunits and showed a dose-dependent behavior (except for INT $\alpha 2$ -subunit in acinar HSG cells). That DHT was not as effective could be for several reasons relating to dose, timing, passive cellular uptake by diffusion, intracrine balance between different sex steroids, or their nongenomic actions. Nevertheless, physiological concentrations of androgens elevated the levels of both INT. Further, these INT receptors were relatively low in intercalated duct cells, but increased about 2-fold upon differentiation to acinar cells on Matrigel, suggesting that contact with laminin $\alpha 1$ and in particular with DHEA and its metabolites increases these tubuloacinar cell-specific INT receptors⁸.

In some experiments, DHEA upregulated INT subunits in healthy salivary glands. However, in SS salivary glands this

androgen-INT link was interrupted, suggesting faulty intracrine DHEA processing^{15,17}. Since the composition and remodeling of ECM is affected by age^{22,23}, we also studied DHEA regulation of INT $\alpha 1$ and $\alpha 2$ in relation to age. The effect of DHEA treatment on the expression of these integrin subunits was perhaps slightly more remarkable in younger individuals and diminished in older tissue, but interindividual variation was also quite remarkable. Additionally, the upregulating effect of DHEA seen in healthy LSG was minor compared with effects seen at the cellular level, especially for INT $\alpha 1$, and this is possibly explained by the presence of INT $\alpha 1$ in cells other than salivary epithelial cells. This effect is also indicated by the finding that the mRNA levels of both INT subunits were higher in LSG in patients with SS than in healthy individuals. INT $\alpha 1$ and $\alpha 2$ -subunits are expressed in CD4-positive and CD8-positive T lymphocytes and lymphocyte infiltrates²⁴. The expression of INT $\alpha 1$ and $\alpha 2$ has been shown in infiltrating T cells in the chronic inflammatory environment and has been suggested as important for generation of inflammatory responses in tissues²⁵ in, for example, rheumatoid arthritis^{26,27}.

Current observations demonstrate for the first time a link between 2 features of SS, androgen depletion^{15,17} and extracellular matrix-INT signaling^{8,10}, by suggesting that androgens regulate molecules involved in matrix-cell signaling. These results highlight the importance of locally balanced intracrine processing and the effects of DHEA. DHEA was incapable of increasing the expression of INT in LSG of




Figure 5. The effect of 100 μ M DHEA stimulation for 72 hours on expression of INT α 1 (A) and α 2 (B) subunits in labial salivary glands of patients with SS (n = 3) and healthy controls (C, n = 4). Nonstimulated samples are marked C D0 (controls, no DHEA) or SS D0 (no DHEA), and DHEA-stimulated samples C D100 (100 μ M DHEA) or SS D100.

patients with SS. In addition to the LM-111-INT α 1 β 1/ α 2 β 1 connection, the connection between LM332 and INT α 6 β 4 is vital for acinar cells. These molecules form adhesion complexes and thus cooperate in attachment of acinar cells to ECM. In SS, disturbances in the expression and localization of these INT molecules have been reported²⁸ and may lead to detachment of acinar cells and further contribute to acinar cell loss. In our study we concentrated on the effects of androgens on acinar-specific INT subunits α 1 and α 2, which have been shown to be decreased in SS⁸, but it would be interesting to study the eventual androgen regulation of INT α 6 β 4 as well. If such a regulation were found, it would mean that androgen depletion in SS deteriorates the attachment, differentiation, and survival of salivary gland acinar cells.

We observed that androgens upregulate molecules essential for progenitor cell migration along, and matrix cell signaling from, LM-111 in the basement membrane to tubuloacinar epithelial cells in human salivary glands. Through this mechanism, low levels of DHEA and its metabolites in salivary glands in patients with SS could impair maintenance of acini.

REFERENCES

- Vitali CS, Bombardieri R, Jonsson R, Moutsopoulos HM, Alexander EL, Carsons SE, et al. Classification criteria for Sjögren's syndrome: a revised version of the European criteria proposed by the American-European Consensus Group. *Ann Rheum Dis* 2002;61:554-8.
- Denny PC, Denny PA. Dynamics of parenchymal cell division, differentiation, and apoptosis in the young adult female mouse submandibular gland. *Anat Rec* 1999;254:408-17.
- Redman RS. On approaches to the functional restoration of salivary glands damaged by radiation therapy for head and neck cancer, with a review of related aspects of salivary gland morphology and development. *Biotech Histochem* 2008;83:103-30.
- Sato M, Azuma M, Hayashi Y, Yoshida H, Yanagawa T, Yura Y. 5-Azacytidine induction of stable myoepithelial and acinar cells from a human salivary intercalated duct cell clone. *Cancer Res* 1987;47:4453-9.
- Man YG, Ball WD, Marchetti L, Hand AR. Contributions of intercalated duct cells to the normal parenchyma of submandibular glands of adult rats. *Anat Rec* 2001;263:202-14.
- Hoffman MP, Kibbey MC, Letterio JJ, Kleinman HK. Role of laminin-1 and TGF- β 3 in acinar differentiation of a human submandibular gland cell line (HSG). *J Cell Sci* 1996;109:2013-21.
- Patarroyo M, Tryggvason K, Virtanen I. Laminin isoforms in tumor invasion, angiogenesis and metastasis. *Cancer Biol* 2002;12:197-207.
- Laine M, Virtanen I, Porola P, Rotar Z, Rozman B, Poduval P, et al. Acinar epithelial cell laminin-receptors in labial salivary glands in Sjögren's syndrome. *Clin Exp Rheumatol* 2008;26:807-13.
- Kwon YJ, Pérez P, Aguilera S, Molina C, Leyton L, Alliende C, et al. Involvement of specific laminins and nidogens in the active remodeling of the basal lamina of labial salivary glands from patients with Sjögren's syndrome. *Arthritis Rheum* 2006;54:3465-75.
- Laine M, Virtanen I, Salo T, Kontinen YT. Segment-specific but pathologic laminin isoform profiles in human labial salivary glands of patients with Sjögren's syndrome. *Arthritis Rheum* 2004;50:3968-73.
- Daniel TE. Labial salivary gland biopsy in Sjögren's syndrome: assessment as a diagnostic criterion in 362 suspected cases. *Arthritis Rheum* 1984;27:147-56.
- Goicovich E, Molina C, Perez P, Aguilera S, Fernández J, Olea N, et al. Enhanced degradation of proteins of the basal lamina and stroma by matrix metalloproteinases from the salivary glands of Sjögren's syndrome patients: correlation with reduced structural integrity of acini and ducts. *Arthritis Rheum* 2003;48:2573-84.
- Poduval P, Sillat T, Virtanen I, Porola P, Kontinen YT. Abnormal basement membrane type IV collagen alpha chain composition in labial salivary glands in Sjögren's syndrome. *Arthritis Rheum* 2009;60:938-45.
- Laine M, Porola P, Udby L, Kjeldsen L, Cowland JB, Borregaard N, et al. Low salivary dehydroepiandrosterone and androgen-regulated cysteine-rich secretory protein 3 levels in Sjögren's syndrome. *Arthritis Rheum* 2007;56:2575-84.
- Porola P, Virkki L, Przybyla BD, Laine M, Patterson TA, Pihakari A, et al. Androgen deficiency and defective intracrine processing of DHEA in the salivary glands in Sjögren's syndrome. *J Rheumatol* 2008;35:2229-35.
- Labrie F, Luu-The V, Labrie C, Simard J. DHEA and its transformation into androgens and estrogens in peripheral target tissues: intracrinology. *Front Neuroendocrinol* 2001;22:185-212.
- Spaan M, Porola P, Laine M, Rozman B, Azuma M, Kontinen YT. Healthy human salivary glands contain a DHEA-sulphate processing intracrine machinery, which is deranged in primary Sjögren's syndrome. *J Cell Mol Med* 2009;13:1261-70.

18. Shirasuna K, Sato M, Miyazaki T. A neoplastic epithelial duct cell line established from an irradiated human salivary gland. *Cancer* 1981;48:745-52.
19. Royce LS, Kibbey MC, Mertz P, Kleinman HK, Baum BJ. Human neoplastic submandibular intercalated duct cells express an acinar phenotype when cultured on a basement membrane matrix. *Differentiation* 1993;52:247-55.
20. Hemler ME, Sanchez-Madrid ME, Flotte TJ, Krensky AM, Burakoff SJ, Bhan AK, et al. Glycoproteins of 210,000 and 130,000 m.w. on activated T cells: cell distribution and antigenic relation to components on resting cells and T cell lines. *J Immunol* 1984;132:3011-8.
21. Giltay JC, Brinkman HJ, Modderman PW, von dem Borne AE, van Mourik JA. Human vascular endothelial cells express a membrane protein complex immunohistochemically indistinguishable from the platelet VLA-2 (glycoprotein Ia-IIa) complex. *Blood* 1989;73:1235-41.
22. Albon J, Karwatowski WS, Easty DL, Sims TJ, Duance VC. Age related changes in the non-collagenous components of the extracellular matrix of the human lamina cribrosa. *Br J Ophthalmol* 2000;84:311-7.
23. Eikmans M, Baelde HJ, De Heer E, Bruijn JA. Effect of age and biopsy site on extracellular matrix mRNA and protein levels in human kidney biopsies. *Kidney Int* 2001;60:974-81.
24. Richter M, Ray SJ, Chapman TJ, Austin SJ, Rebhahn J, Mosmann TR, et al. Collagen distribution and expression of collagen-binding alpha 1 beta 1 (VLA-1) and alpha 2 beta 1 (VLA-2) integrins on CD4 and CD8 T cells during influenza infection. *J Immunol* 2007;178:4506-16.
25. Andreasen SØ, Thomsen AR, Kotliansky VE, Novobrantseva TI, Sprague AG, de Fougerolles AR, et al. Expression and functional importance of collagen-binding integrins, alpha 1 beta 1 and alpha 2 beta 1, on virus-activated T cells. *J Immunol* 2003;171:2804-11.
26. Hemler ME, Glass D, Coblyn JS, Jacobson JG. Very late activation antigens on rheumatoid synovial fluid T lymphocytes. Association with stages of T cell activation. *J Clin Invest* 1986;78:696-702.
27. Bank I, Roth D, Book M, Guterman A, Shnirer I, Block R, et al. Expression and functions of very late antigen 1 in inflammatory joint diseases. *J Clin Immunol* 1991;11:29-38.
28. Velozo J, Aguilera S, Alliende C, Ewert P, Molina C, Pérez P, et al. Severe alterations in expression and localisation of alpha 6 beta 4 integrin in salivary gland acini from patients with Sjogren syndrome. *Ann Rheum Dis* 2009;68:991-6.