

Clinical Relevance of Nitric Oxide Metabolites and Nitrative Stress in Thrombotic Primary Antiphospholipid Syndrome

PAUL R.J. AMES, JOANA R. BATUCA, ANTONIO CIAMPA, LUIGI IANNACCONE, and JOSE DELGADO ALVES

ABSTRACT. Objective. To assess the role of nitrite (NO_2^-), nitrate (NO_3^-), and nitrative stress in thrombotic primary antiphospholipid syndrome (PAPS).

Methods. We investigated 46 patients with PAPS: 21 asymptomatic but persistent carriers of antiphospholipid antibodies (PCaPL), 38 patients with inherited thrombophilia (IT), 33 patients with systemic lupus erythematosus (SLE), and 29 healthy controls (CTR). IgG anticardiolipin (aCL), IgG anti-beta₂-glycoprotein I (anti-β₂-GPI), IgG anti-high density lipoprotein (aHDL), IgG anti-apolipoprotein A-I (aApoA-I), crude nitrotyrosine (NT) (an indicator of nitrative stress), and high sensitivity C-reactive protein (CRP) were measured by immunoassays. Plasma nitrite (NO_2^-), nitrate (NO_3^-), and total antioxidant capacity (TAC) were measured by colorimetric spectroscopic assays.

Results. Average plasma NO_2^- was lower in PAPS, PCaPL, and IT ($p < 0.0001$); average NO_3^- was highest in SLE ($p < 0.0001$), whereas average NT was higher in PAPS and SLE ($p = 0.01$). In thrombotic PAPS, IgG aCL titer and number of vascular occlusions negatively predicted NO_2^- ($p = 0.03$ and $p = 0.001$, respectively), whereas arterial occlusions and smoking positively predicted NO_3^- ($p = 0.05$ and $p = 0.005$), and CRP positively predicted NT ($p = 0.004$). In the PCaPL group IgG aCL negatively predicted NO_3^- ($p = 0.03$). In the SLE group IgG aCL negatively predicted NO_2^- ($p = 0.03$) and NO_3^- ($p = 0.02$).

Conclusion. PAPS is characterized by decreased NO_2^- in relation to type and number of vascular occlusions and to aPL titers. Nitrative stress and low grade inflammation are linked phenomena in PAPS and may have implications for thrombosis and atherosclerosis. (J Rheumatol First Release Oct 1 2010; doi:10.3899/jrheum.100494)

Key Indexing Terms:

ANTIPHOSPHOLIPID SYNDROME
NITRIC OXIDE

NITRITE

THROMBOSIS
NITRATE

The primary antiphospholipid syndrome (PAPS) is characterized by venous and arterial thromboses, recurrent miscarriages, and premature atherosclerosis in persistent carriers of antibodies against β₂-glycoprotein I (anti-β₂-GPI) and other coagulation proteins in the absence of any other under-

lying immune disorder^{1,2}. From a biochemical standpoint PAPS is also characterized by an antioxidant/oxidant balance tilted towards the latter, partly due to decreased paraoxonase activity and enhanced oxidative stress^{3,4}. Indeed, IgG anticardiolipin (aCL) antibody titers positively correlated to plasma levels of F₂-isoprostanes, a marker of increased oxidative stress, and to decreased urinary excretion of nitric oxide ($\text{NO}\cdot$) metabolites in PAPS³. $\text{NO}\cdot$ is the main endothelial vasodilator agent, and interference with $\text{NO}\cdot$ biology induces vascular dysfunction, particularly in the early phases of atherosclerosis⁵. After physiological stimulation of constitutive endothelial nitric oxide synthase (eNOS)⁶ or inflammatory activation of inducible (iNOS) enzyme⁷, $\text{NO}\cdot$ is released at higher rates and behaves as a pathogenic mediator or a cytotoxic molecule.

In the latter case, most $\text{NO}\cdot$ mediated pathogenicity depends on formation of secondary intermediates such as peroxynitrite anion (ONOO^-) and nitrogen dioxide ($\text{NO}_2\cdot$), which are typically more reactive and toxic than $\text{NO}\cdot$ per se⁸. In the presence of oxidants such as superoxide radical ($\text{O}_2^{\cdot-}$) $\text{NO}\cdot$ gives rise to ONOO^- , a strong 1-electron and

From the Departments of Rheumatology and Haematology, Airedale General Hospital, Steeton, UK; Pharmacology Department, Faculty of Medical Sciences of Lisbon, Lisbon, Portugal; Chemical Pathology, Cardarelli Hospital, Naples, Italy; and Haemostasis Unit, Moscati Hospital, Avellino, Italy.

Supported in part by Senit Foundation (Greenock, Scotland, UK).

P.R.J. Ames, MD, MSc, Consultant Haematologist/Rheumatologist, Rheumatology and Haematology Departments, Airedale General Hospital; J.R. Batuca, BSc, PhD Student, Pharmacology Department, Faculty of Medical Sciences of Lisbon; A. Ciampa, MD, Senior Registrar in Haemostasis and Thrombosis, Haemostasis Unit, Moscati Hospital; L. Iannaccone, Senior Biomedical Scientist, Chemical Pathology, Cardarelli Hospital; J. Delgado Alves, MD, PhD, Consultant, Internal Medicine, Senior Lecturer in Clinical Pharmacology, Pharmacology Department, Faculty of Medical Sciences, University of Lisbon.

Address correspondence to Dr. P.R.J. Ames, Airedale General Hospital, Skipton Road, Steeton, Keighley, West Yorkshire, BD20 6TD, UK.

E-mail: paxmes@aol.com

Accepted for publication July 30, 2010.

Personal non-commercial use only. The Journal of Rheumatology Copyright © 2010. All rights reserved.

2-electron oxidant with such a short biological half-life (10–20 ms) that it cannot be measured directly but must be inferred by indirect methods^{9,10}. In fact ONOO⁻ interacts with CO₂ to give nitrosoperoxycarbonate (ONOOOCO₂⁻) that will nitrate tyrosine residues in proteins⁶: measurement of nitrated proteins therefore represents a fingerprint of the interaction of O₂^{•-} with NO[•]^{6,11}.

Possible involvement of NO[•] in APS has been explored in animal studies^{12,13} and in a few patient series whose numbers were too limited to provide a full understanding of its significance^{14,15}. We therefore hypothesized that NO[•] might play a role in the vascular pathogenesis of PAPS, and compared the behavior of NO[•] metabolites nitrite (NO₂⁻) and nitrate (NO₃⁻), total antioxidant capacity (TAC) (expressed as ONOO⁻ quenching), and nitrotyrosine (NT) in patients with thrombotic PAPS, in asymptomatic but persistent carriers of antiphospholipid antibodies (PCaPL), in patients with inherited thrombophilia (IT) with vascular occlusions, in patients with systemic lupus erythematosus (SLE), and in healthy subjects. Possible relationships between NO₂⁻, NO₃⁻, C-reactive protein (CRP), and several aPL were also investigated.

MATERIALS AND METHODS

Patients. Our study was devised as a cross-sectional case-quadruple control: PAPS patients with vascular occlusions represented cases; IT patients with vascular occlusions represented thrombotic controls PCaPL without vascular occlusions represented nonthrombotic aPL-positive controls; patients with SLE represented inflammatory controls; and healthy subjects represented normal controls. All participants were age- and sex-matched (where possible), except for SLE patients, who were all female. Consecutive patients with thrombotic PAPS, according to recent criteria¹, with IT and persistent aPL attending the Coagulation Unit of the Cardarelli Hospital (Naples, Italy) were invited to participate between January 2008 and July 2008. Our study was carried out according to the revised Declaration of Helsinki, with approval of the Ethics Board of the hospital and written consent of all participants. Exclusion criteria were acute or chronic hepatic, renal, and lung disease; diabetes; acute infection (within 6 weeks); post-thrombotic syndrome with or without venous ulcerations; positive urinary dipstick for nitrites on the day of sampling and treatment with statins or fibrates. PAPS and IT patients are seen on average every 3 to 4 weeks for oral anticoagulation monitoring and are instructed to self-report any illness during the intervening periods; their lipid profiles and kidney and liver function tests are checked annually. PCaPL subjects were diagnosed as such either because of the presence of prolonged clotting tests in routine assays, subsequently confirmed as lupus anticoagulants (LAC), or because of thrombocytopenia or other symptoms that prompted a search for aPL. Of the PAPS attendees (n = 50), 2 were excluded because they had gradually developed ankylosing spondylitis and SLE, one had developed kidney cancer, 2 were pregnant, one had suffered a recent recurrent event, one had post-thrombotic syndrome, and 2 were evasive regarding their smoking and contraceptive status. Of the IT (n = 46) attendees, 2 were excluded for post-thrombotic syndrome and venous ulcerations in lower limbs. Of the PCaPL attendees (n = 27) one was excluded for development of non-insulin-dependent diabetes; one for the development of SLE, hemolytic anemia, nephrotic syndrome, and pulmonary embolism after ovarian hyperstimulation; one for the development of chronic lymphoid leukemia; one for spontaneous onset of ischemic stroke; and 2 had moved to a different town. Of the remaining aPL subjects 4 had moderate thrombocytopenia (platelets < 100 × 10⁹/l) not requiring treatment.

Consecutive patients with SLE fulfilling the American Rheumatism Association (ACR) criteria¹⁶ were enrolled among those attending the Autoimmune Outpatient Clinic of the Curry Cabral Hospital, Lisbon (Portugal) between January 2008 and August 2008. Exclusion criteria included acute or chronic renal impairment that would significantly alter NO metabolites, liver cirrhosis, diabetes, acute infection (within 6 weeks), post-thrombotic syndrome with or without venous ulcerations, positive urinary culture following positive dipstick for nitrite (urinary excretion in SLE may be increased in the absence of infection), and treatment with statins or fibrates.

Of 52 patients with SLE, 16 were excluded on the basis of the above criteria. Of the remaining 36, one was found weeks later to have tuberculosis and 2 were pregnant; their samples were discarded. Therefore 33 SLE patients participated in the study: of these, 9% had visceral involvement without renal disease, 12% cardiac and lung involvement, 9% central nervous system involvement, 66% arthritis, 3% myositis, 9% alopecia, 3% hemolytic anemia, 21% thrombocytopenia, 9% neutropenia, 81% presence of anti-DNA antibodies, and 90% presence of antinuclear antibodies. The average SLE Disease Activity Index (SLEDAI) score was 4.85 ± 3.96 (median 3.5, range 0–16). Their medication intake was: prednisolone in 54%, (< 6 mg/day in 27%, 6–10 mg/day in 18%, > 10 mg/day in 9%) azathioprine in 24% (100 mg/day in 18%, 150 mg/day in 6%), hydroxychloroquine 200 mg/day in 60%, aspirin in 12%, warfarin in 9%. Twenty-nine healthy hospital staff served as normal controls: 15 from Cardarelli Hospital in Naples and 14 from the Curry Cabral Hospital in Lisbon. To minimize dietary influences on nitric oxide metabolite concentrations all participants were asked to refrain from foodstuffs containing high concentrations of nitrate/nitrite (such as lettuce, spinach, beetroot, radish, salamis, and pickled items) for 3 days before blood sampling, which was drawn between 8:00 and 10:00 AM. Blood samples were drawn by neat venepuncture into 5 ml citrate vacutainers, spun immediately at room temperature at 4000 rpm for 6 min; supernatant plasma was spun again at room temperature at 12,000 rpm for 4 min to obtain platelet-poor plasma: aliquots were frozen at –80°C and thawed on the day of testing. The study was therefore carried out on 46 thrombotic PAPS patients, 21 PCaPL subjects, 38 IT patients, 33 SLE patients, and 29 control subjects. Their demographics are shown in Table 1.

Determination of antiphospholipid antibodies. All participants had their aPL determined according to established criteria¹⁷; LAC screened by activated partial thromboplastin time (aPTT) and dilute Russell's viper venom time (DRVVT)¹⁷. A clotting time ratio between sample and control plasma > 1.2 for aPTT and > 1.18 for DRVVT indicated an abnormal result. After demonstrating the presence of an inhibitor using mixing studies, the platelet neutralization procedure confirmed the presence of a lupus inhibitor in aPTT and DRVVT. IgG aCL (Cambridge Life Sciences, Ely, UK) and IgG anti-β₂-GPI (Corgenix, Broomfield, CO, USA) were measured by ELISA according to manufacturer's instructions. Since the inception of the PAPS cohort (1994), after initial diagnosis with repeat testing of aPL after 6 weeks, IgG aCL was measured yearly, whereas IgG anti-β₂-GPI was measured yearly only since 2004.

Measurement of IgG anti-high density lipoprotein (aHDL) antibodies, IgG anti-apolipoprotein A-I (aApo A-I) antibodies, plasma nitrotyrosine, and high sensitivity C-reactive protein. aHDL and aApo A-I were measured by ELISA as described⁴; similarly ELISA was employed to measure nitrotyrosine (HyCult Biotechnology, Uden, The Netherlands) and high sensitivity CRP (Biosupply Ltd., Bradford, UK) according to the manufacturer's instructions. "CRP" stands for the high-sensitivity test throughout this article.

Measurement of plasma nitrate and nitrite. Nitric oxide metabolites nitrite (NO₂⁻) and nitrate (NO₃⁻) were determined using a modified Griess reaction, following the reduction of nitrate to nitrite using nitrate reductase and nicotinamide adenine dinucleotide phosphate (NADPH). Briefly, the assay was performed in a standard flat-bottomed 96-well microtiter plate half divided for simultaneous measurement of nitrite and nitrate concentration. To each well was added 50 μl/well of standard or diluted sample (1 in 4

Table 1. Demographics of participants in study groups.

	PAPS, n = 46	PCaPL, n = 21	IT, n = 38	SLE, n = 33	CTR, n = 29
Age, yrs (mean \pm SD)	43 \pm 12	40 \pm 9	41 \pm 14	40 \pm 12	41 \pm 7
M/F	18/28	6/15	13/25	0/33	15/14
Disease duration, yrs, median (range)	10 (1–18)	10 (1–16)	12 (1–22)	8 (1–13)	
LAC	44	18	0	6	0
IgG aCL < 40 (GPL)	15	14	38	22	29
IgG aCL 41–80 (GPL)	7	3	0	8	0
IgG aCL > 80 (GPL)	24	4	0	3	0
IgG aCL, median (range)	100 (5–500)	24 (2–309)	8.4 (4–20)	62 (12–112)	8 (2–18)
IgG anti- β_2 -GPI U/ml, median (range)	146 (2.6–184)	4 (2–90)	5 (2–12)	12 (3.6–102)	4 (2.5–8.2)
IgG aHDL, %, median (range)	70 (25–436)	112 (28–240)	70 (22–292)	98 (25.4–328)	61 (19–161)
IgG aApoA-I, %, median (range)	0.69 (0.17–6.3)	1.2 (0.17–7.4)	0.14 (0.02–1.2)	1.2 (0.17–6)	0.18 (0.06–0.57)
MTHFR +/-	13	6	5	4	5
PT20210	2	1	4	0	2
FVL	2	0	12	1	1
F-PS deficiency	32	4	8	NA	0
FVL + PT20210	0	0	7	0	0
FVL + PS deficiency	0	0	2	0	0
Thrombosis					
Arterial	14	0	11	4	0
Venous	26	0	27	1	0
Arterial + venous	6	0	0	0	0
No. of occlusions					
1	19	0	38	4	0
2	16	0	0	0	0
3	3	0	0	0	0
4	2	0	0	0	0
Smokers					
< 6 per day	6	3	4	2	4
7–15 per day	5	2	8	5	6
> 15 per day	4	0	1	1	2

PAPS: primary antiphospholipid syndrome; PCaPL: persistent carriers of antiphospholipid antibody with no underlying disorder and no thrombosis; IT: inherited thrombophilia; SLE: systemic lupus erythematosus; CTR: normal controls; LAC: lupus anticoagulant; aCL: anticardiolipin; β_2 GPI: beta-2-glycoprotein-1; aHDL: anti-high density lipoprotein; aApoA-I: anti-apolipoprotein A-I; MTHFR: homozygous methylenetetrahydrofolate reductase C667T thermolabile mutation; PT: heterozygous prothrombin; FVL: heterozygous factor V Leiden; F-PS: free protein S.

with phosphate buffer pH 7.4) in duplicate. The assay was blanked against phosphate buffer. In half plate, 4 μ l of nitrate reductase (Sigma-Aldrich) and 10 μ l of NADPH (Sigma-Aldrich) were added to each well, giving a final concentration of 6.3 U/l and 550 μ mol/l, respectively. The plate was incubated at room temperature for 2 h. Griess reaction was initiated by addition to each well of equal volumes of 2% sulfanilamide (Sigma-Aldrich) in H_3PO_4 5% and 0.2% N-(1-naphthyl)-ethylenediamine dihydrochloride (Sigma-Aldrich) in water, mixed just before use. After 10 min incubation at room temperature the absorbance of the reaction mixture was measured at 540 nm and the levels expressed as μ M.

Measurement of total antioxidant capacity of plasma. TAC of plasma was measured by peroxynitrite (ONOO^-) quenching: 100 μ l of phosphate buffer (50 mM, pH 7.4) containing Pholasin[®] (1.7 μ g/ml) was pipetted into a microcuvette. Plasma or buffer for control (5 μ l) was added. The reaction was initiated by adding 3-morpholino-sydnonimine HCl (SIN-1; 2 μ l of 2 mg/ml in water), and light emission was measured continuously at 5 min intervals until the maximum reading was obtained. Antioxidant capacity was expressed as the time at which maximum light was emitted. Lower values reflect decreased plasma TAC (peroxynitrite-related).

Statistical analysis. Variables were compared by ANOVA (post-hoc analysis) and ANCOVA with log transformation of variables that did not follow a normal distribution. The assumptions of univariate analysis within groups (not shown) were tested by multiple regression models. All statistical analyses were done using SPSS (SPSS, Chicago, IL, USA).

RESULTS

Comparison of variables in PAPS, PCaPL, IT, SLE, and healthy controls. Average plasma NO_2^- was lower in the PAPS, PCaPL, and IT groups (Figure 1A), whereas NO_3^- was higher in SLE (Figure 1B) and NT was higher in SLE and PAPS (Figure 1C). Mean plasma TAC was lowest in SLE (Figure 2A), where CRP was highest (Figure 2A and 2B). Average TAC was higher in males than in females in all non-SLE groups: in PAPS 11280 ± 3041 versus 9749 ± 2967 μ mol/l ($p = 0.02$); in IT 6392 ± 1399 versus 5325 ± 1720 μ mol/l ($p = 0.04$); and in healthy controls 7967 ± 991 vs 6194 ± 1265 μ mol/l ($p = 0.001$).

Age, sex, and smoking correlated to NO_2^- , NO_3^- , and TAC but had no confounding effect on resulting significance findings by ANCOVA. Age and IgG aCL related to NT and their confounding effect by ANCOVA reduced the comparative significance ($p < 0.02$).

Relationship among variables in PAPS. The effect of antibodies and that of other clinical and laboratory variables on plasma concentrations of NO_2^- , NO_3^- , and NT was tested by




Figure 1. Comparison (ANOVA) of nitrite (NO_2^-), nitrate (NO_3^-), and nitrotyrosine (NT) across study groups: (A) average concentration of NO_2^- (Bonferroni's multiple comparison test: PAPS vs SLE, $p < 0.001$; PAPS vs CTR, $p < 0.001$; PCaPL vs SLE, $p < 0.01$; aPL vs CTR, $p < 0.001$; IT vs SLE, $p < 0.01$; IT vs CTR, $p < 0.001$); (B) average concentration of NO_3^- (Bonferroni's multiple comparison test: PAPS vs SLE, $p < 0.001$; IT vs SLE, $p < 0.001$; SLE vs CTR, $p < 0.001$); (C) average concentration of NT (Bonferroni's multiple comparison test: PAPS vs IT, $p < 0.05$). PAPS: primary antiphospholipid syndrome; PCaPL: persistent carriers of antiphospholipid antibodies without thrombosis or miscarriages; IT: inherited thrombophilia; SLE: systemic lupus erythematosus; CTR: controls.




Figure 2. Comparison (ANOVA) of total antioxidant capacity (TAC) and high sensitivity C-reactive protein (CRP) across study groups: (A) average concentration of TAC (Bonferroni's multiple comparison test: PAPS vs IT, $p < 0.001$; PAPS vs SLE, $p < 0.001$; IT vs SLE, $p < 0.05$; IT vs CTR, $p < 0.05$; SLE vs CTR, $p < 0.001$); (B) average concentration of CRP (Bonferroni's multiple comparison: SLE vs PAPS, SLE vs aPL, SLE vs IT, SLE vs CTR, all $p < 0.001$). PAPS: primary antiphospholipid syndrome; PCaPL: persistent carriers of antiphospholipid antibody positive without thrombosis or miscarriages; IT: inherited thrombophilia; SLE: systemic lupus erythematosus; CTR: controls.

separate multiple regression models. In the model with NO_2^- as the dependent variable and IgG aCL, IgG anti- β_2 -GPI, IgG aHDL, and IgG aApoA-I antibodies as independent variables, IgG aCL resulted in the only negative predictor of NO_2^- ($p = 0.03$; Table 2). Average NO_2^- was lower in patients with a history of arterial thrombosis versus those with venous thrombosis (11.41 ± 7.6 vs 18.43 ± 11.06 $\mu\text{mol/l}$; $p = 0.03$) although in a separate model with NO_2^- as the dependent variable and age at first thrombotic event and thrombosis number and type as the independent variables, thrombosis number negatively predicted NO_2^- ($p = 0.001$; Table 2).

In the model with NO_3^- as the dependent variable and IgG aCL, IgG anti- β_2 -GPI, IgG aHDL, and IgG aApoA-I antibodies as independent variables, IgG aCL was the only negative predictor of NO_3^- ($p = 0.03$) (Table 2). In a different model, with NO_3^- as the dependent variable and age,

Table 2. Regression model predictors of nitric oxide metabolites and nitrotyrosine in primary antiphospholipid syndrome.

Independent Variables	Dependent Variables	Predictors	t	p
IgG aCL, IgG anti- β_2 -GPI, IgG aApoA-I	NO_2^-	IgG aCL	-1.87	0.03
Age at first thrombosis, thrombosis number, thrombosis type	NO_2^-	Thrombosis number	-3.24	0.001
IgG aCL, IgG anti- β_2 -GPI, IgG aHDL, IgG aApoA-I	NO_3^-	IgG aCL	-1.93	0.03
Age, sex, thrombosis type, smoking, TAC	NO_3^-	Arterial	1.67	0.05
		Smoking	2.66	0.005
NO_3^- , NO_2^- , TAC, smoking, CRP	NT	CRP	2.74	0.004

IgG aCL: anticardiolipin; IgG anti- β_2 -GPI: beta-2-glycoprotein-I; IgG aHDL, anti-high density lipoprotein; IgG aApoA-I, anti apolipoprotein A-I; NO_2^- : nitrite; NO_3^- : nitrate; TAC: total antioxidant capacity; NT: nitrotyrosine; CRP, C-reactive protein.

sex, thrombosis type, smoking, and TAC as the independent variables, arterial thrombosis and smoking independently predicted NO_3^- ($p = 0.05$ and $p = 0.005$, respectively).

In a further model with NT as the dependent variable and the antibodies as the independent variables none of the latter bore any relationship with NT; but in a similar model NT as the dependent variable and with NO_3^- , NO_2^- , TAC, smoking, and CRP as independent variables, CRP was the only independent predictor of NT ($p = 0.004$) (Table 2).

Relationship among variables in PcaPL. In the regression model with NO_2^- as the dependent variable and IgG aCL, IgG anti- β_2 -GPI, IgG aHDL, IgG aApoA-I, aPTT, and DRVVT as independent variables, IgG anti- β_2 -GPI showed only a negative trend with NO_2^- ($p = 0.07$) (Table 3).

In the model with NO_3^- as the dependent variable and IgG aCL, IgG anti- β_2 -GPI, IgG aHDL, IgG aApoA-I, aPTT, and DRVVT as independent variables, negative predictors were IgG aCL ($p = 0.03$) and DRVVT ($p = 0.03$), and a trend was seen for IgG anti- β_2 -GPI ($p = 0.06$; Table 3).

In a further model none of the antibodies bore any relationship with NT as the dependent variable but with NO_3^- , NO_2^- , TAC, and CRP set as independent variables, NO_2^- negatively predicted NT ($p = 0.05$; Table 3).

Relationship among variables in IT. In the regression model with NT as the dependent variable and NO_3^- , NO_2^- , TAC, smoking, and CRP as independent variables, only CRP independently predicted NT ($p = 0.0006$; Table 3).

Relationship among variables in SLE. In the regression model with NO_3^- as the dependent variable and IgG anti- β_2 -GPI, IgG aHDL, IgG aApoA-I, and IgG aCL as the independent variable, IgG aCL negatively predicted NO_3^- ($p = 0.03$); a similar result was obtained when NO_2^- was substituted for NO_3^- ($p = 0.02$; Table 3). In the model with NT as the dependent variable and NO_3^- , NO_2^- , TAC, smoking, and CRP as independent variables, NO_2^- negatively predicted NT ($p = 0.002$) and NO_3^- positively predicted NT ($p = 0.001$; Table 3). Finally, in the model with SLEDAI as the dependent variable and NO_3^- , NO_2^- , TAC, CRP, smoking,

Table 3. Regression model predictors of nitric oxide metabolites and nitrotyrosine in non-primary antiphospholipid antibody syndrome groups.

Independent Variables	Dependent Variables	Predictors	t	p
Persistent carriers of antiphospholipid antibodies				
IgG aCL, IgG anti- β -GPI, IgG aHDL, IgG aApoA-I, aPTT, DRVVT	NO_3^-	IgG aCL	-2.06	0.03
		DRVVT	-1.93	0.03
		IgG a β_2 GPI	-1.68	0.06
NO_3^- , NO_2^- , TAC, smoking, CRP	NT	NO_2^-	-1.74	0.05
Inherited thrombophilia				
NO_3^- , NO_2^- , TAC, smoking, CRP	NT	CRP	3.75	0.0006
Systemic lupus erythematosus				
IgG aCL, IgG anti- β_2 -GPI, IgG aHDL, IgG aApoA-I	NO_2^-	IgG aCL	-2.12	0.02
IgG aCL, IgG anti- β_2 -GPI, IgG aHDL, IgG aApoA-I	NO_3^-	IgG aCL	-1.84	0.03
NO_3^- , NO_2^- , TAC, smoking, CRP	NT	NO_2^-	-3.25	0.002
		NO_3^-	3.65	0.001
NO_3^- , NO_2^- , TAC, CRP, smoking, NT	SLEDAI	NT	2.55	0.009
		CRP	1.44	0.08
Normal controls				
Age, sex, TAC, CRP, smoking, NT	NO_3^-	Smoking	2.90	0.003
Age, sex, NO_3^- , NO_2^- , TAC, CRP, smoking	NT	Smoking	2.21	0.02
		NO_3^-	1.75	0.04

IgG aCL: anticardiolipin; IgG anti- β_2 -GPI: beta-2-glycoprotein-I; IgG aHDL, anti-high density lipoprotein; IgG aApoA-I, anti apolipoprotein A-I; aPTT: activated partial thromboplastin time; DRVVT: dilute Russell viper venom time; NO_2^- : nitrite; NO_3^- : nitrate; TAC: total antioxidant capacity; NT: nitrotyrosine; CRP, C-reactive protein; SLEDAI: Systemic Lupus Erythematosus Disease Activity Index..

and NT as the independent variables, NT predicted SLEDAI ($p = 0.009$) and a trend was seen for CRP ($p = 0.08$; Table 3).

Relationship among variables in the control group. No effect on NO_2^- was seen in a multiple regression model with NO_2^- as the dependent variable and age, sex, smoking, IgG aHDL, and IgG aApo-I as explanatory variables. In a similar model where NO_3^- was set as the dependent variable, smoking independently predicted NO_3^- ($p = 0.003$; Table 3). Similarly, when NT was set as the dependent variable with age, sex, smoking, TAC, NO_2^- , and NO_3^- as explanatory variables, smoking independently predicted NT ($p = 0.02$) alongside NO_3^- ($p = 0.04$; Table 3).

DISCUSSION

$\text{NO}\bullet$ is synthesized in the vasculature by 2 related nitric oxide synthases (NOS), constitutive eNOS and inducible NOS; both convert L-arginine to $\text{NO}\bullet$ and citrulline at different concentrations according to substrate availability¹⁸. The role of $\text{NO}\bullet$ in PAPS is unknown: one study found lower urinary NO_2^- in a small number of patients with PAPS in negative correlation with IgG aCL titer³. Of the $\text{NO}\bullet$ metabolites, it is widely accepted that in humans, only NO_2^- reflects changes in eNOS activity^{19,20} and endothelial dysfunction²¹ known to be impaired in APS²².

To evaluate the clinical significance of NO_2^- and NO_3^- with regard to thrombosis in PAPS we employed as comparator patients with IT who had vascular occlusions, patients with PCaPL who had no vascular occlusions, patients with SLE as an inflammatory disease control group, and healthy subjects. The low average concentration of NO_2^- found in PAPS and IT suggests that reduced NO_2^- may be involved in the vascular events of these patients, although causality cannot be established since vessel occlusion might have led to reduced NO_2^- : in fact, the number of vascular occlusions was a negative independent predictor of NO_2^- in PAPS. Nevertheless NO_2^- was also low in the PCaPL group who never had vessel occlusions, suggesting that impaired NO_2^- generation may precede and hence represent a predisposing factor for thrombosis.

Reduced $\text{NO}\bullet$ has a wider importance in the vascular biology of PAPS. $\text{NO}\bullet$ maintains vascular homeostasis against the vasopressor effects of endothelin-1, of isoprostanes derived from lipid peroxidation, and of thromboxane generated after platelet activation (as reviewed²³). Indeed, elevated plasma and/or urinary levels of the aforementioned molecules have all been described in PAPS^{15,24,25}; hence loss of the antiplatelet effect of $\text{NO}\bullet$ may be relevant to thrombosis²⁶, whereas loss of its vasodilator effect may be relevant both to thrombosis and to atherosclerosis, as recently confirmed in PAPS². In the latter study, carried out on almost the same cohort of PAPS patients, diastolic blood pressure in patients with arterial thrombosis was higher than in patients with venous thrombosis².

Of further interest, aPL negatively predicted NO_3^- and/or

NO_2^- in the PAPS, PCaPL, and SLE groups, adding further to the pathogenic potential of aPL. We demonstrated that monoclonal IgG aCL was associated with decreased concentrations of $\text{NO}\bullet$ metabolites in a mouse model¹². Inducible NOS may generate a 1000-fold higher concentration of $\text{NO}\bullet$ than eNOS, which is associated with vascular damage and cytotoxic effects, whereas $\text{NO}\bullet$ is generated by eNOS for short periods of time to maintain vascular homeostasis²⁷. Likely only the latter pathway is impaired in PAPS, whereas the former pathway may be more active in SLE, the group that showed a greater concentration of NO_3^- , although a large difference was not seen because our patients with SLE were mostly clinic attendees devoid of acute or chronic renal disease with low disease activity, hence inflammatory activity. Notwithstanding, our data in SLE would be consistent with possible iNOS activation, cytotoxic release of $\text{NO}\bullet$ ultimately leading to tyrosine nitration due to the inflammatory nature of the disease²⁷.

Among the antibodies that might have had an influence on $\text{NO}\bullet$ we included IgG aHDL and IgG aApoA-I because in previous work we demonstrated that their average plasma concentrations were elevated in SLE and PAPS, where they adversely affected the antioxidant system associated with HDL, favoring oxidation^{4,28}. In our present study, they failed to show any relation with $\text{NO}\bullet$ metabolites, indirectly confirming their specificity in blunting the antiatherogenic and antiinflammatory effects of HDL⁴.

To investigate nitrate stress in PAPS we measured crude plasma NT: this was higher in the SLE group, where NT related to disease activity and was predicted by NO_3^- , in keeping with findings from other inflammatory rheumatic disorders²⁹. On the other hand, having demonstrated that low grade inflammation characterizes PAPS³⁰, we found that CRP was an independent predictor of NT in the PAPS group, suggesting that nitrate stress and low grade inflammation may be related phenomena in these thrombotic patients. Interestingly, smoking predicted NO_3^- in PAPS, and it is known that active³¹ and passive smoking³² may induce oxidative stress.

Our study has several limitations: (1) its retrospective design prevented a full appreciation of the role of $\text{NO}\bullet$ in thrombosis as most PAPS patients were diagnosed after vascular occlusion; (2) our SLE group comprised female patients of whom only 5 had a history of thrombosis; however, we had opted for inclusion of the SLE group mostly to show the inflammatory behavior of $\text{NO}\bullet$ rather than to control for thrombosis, which was provided for by the IT group; (3) the method we employed for the measurement of $\text{NO}\bullet$ metabolites is not sensitive enough to detect nanomolar concentrations of NO_2^- and NO_3^- ³³, and we did not evaluate eNOS and/or iNOS gene polymorphisms that may have accounted for differences in measured metabolites^{34,35}, although our groups would have been too small to yield significant data.

In conclusion, our study, alongside our previous animal data¹², indicates a possible impairment of the vascular biology of NO• in PAPS, the consequences of which may be thrombosis and atherosclerosis. With regard to the former, we cannot define whether decreased NO₂⁻ is a cause or an effect of previous thromboses, but the low NO₂⁻ in PCaPL without vessel occlusions and the relationship between NO• metabolites and aPL in the PAPS, PCaPL, and SLE groups indicate that aPL may negatively influence some physiological activities of NO•. With regard to the latter, patients with PAPS exhibit a certain degree of nitrative stress that relates to low grade inflammation, also noted in other settings³⁶: given the finding of NT in vessels of atherosclerotic patients³⁷, this aspect needs to be further explored in PAPS. From a practical point of view, our study provides evidence that smoking should be avoided in patients with PAPS.

ACKNOWLEDGMENT

We are grateful to L. Lopez (Corgenix, Broomfield, CO, USA) for his help with the IgG anti-β₂-GPI assay.

REFERENCES

- Miyakis S, Lockshin MD, Atsumi T, Branch DW, Brey RL, Cervera R, et al. International consensus statement on an update of the classification criteria for definite antiphospholipid syndrome (APS). *J Thromb Haemost* 2006;4:295-306.
- Ames PRJ, Antinolfi I, Scenna G, Gaeta G, Margaglione M, Margarita A. Atherosclerosis in thrombotic primary antiphospholipid syndrome. *J Thromb Haemost* 2009;7:537-42.
- Ames PRJ, Nourooz-Zadeh J, Tommasino C, Alves J, Brancaccio V, Anggard EE. "Oxidative stress" in primary antiphospholipid syndrome. *Thromb Haemost* 1998;79:447-9.
- Delgado Alves J, Ames PRJ, Donohue S, Norouz-Zadeh J, Isenberg DA. Antibodies towards high density lipoprotein, apo-A-I and cardiolipin inversely correlate to paraoxonase activity in systemic lupus erythematosus and primary antiphospholipid syndrome. Possible atherogenic mechanisms. *Arthritis Rheum* 2002; 46:2686-94.
- Lubos E, Handy DE, Loscalzo J. Role of oxidative stress and nitric oxide in atherothrombosis. *Front Biosci* 2008;13:5323-244.
- Pacher P, Beckman JS, Liaudet L. Nitric oxide and peroxynitrite in health and disease. *Physiol Rev* 2007;87:315-24.
- Ponnuswamy P, Ostermeier E, Schröttle A, Chen J, Huang PL, Ertl G, et al. Oxidative stress and compartment of gene expression determine proatherosclerotic effects of inducible nitric oxide synthase. *Am J Pathol* 2009;174:2400-10.
- Thomas DD, Ridnour LA, Isenberg JS, Flores-Santana W, Switzer CH, Donzelli S, et al. The chemical biology of nitric oxide: Implications in cellular signalling. *Free Radic Biol Med* 2008;45:18-31.
- Denicola A, Batthyany C, Lissi E, Freeman BA, Rubbo H, Radi R. Diffusion of nitric oxide into low-density lipoprotein. *J Biol Chem* 2002;277:932-6.
- Radi R, Peluffo G, Alvarez MN, Naviliat M, Cayota A. Unraveling peroxynitrite formation in biological systems. *Free Radic Biol Med* 2001;30:463-88.
- Peluffo G, Radi R. Biochemistry of protein tyrosine nitration in cardiovascular pathology. *Cardiovasc Res* 2007;75:291-302.
- Delgado Alves J, Mason LJ, Ames PRJ, Chen PP, Rauch J, Levine JS, et al. Antiphospholipid antibodies are associated with enhanced oxidative stress, decreased plasma nitric oxide and paraoxonase activity in an experimental mouse model. *Rheumatology* 2005;44:1238-44.
- Belizna C, Lartigue A, Favre J, Gilbert D, Tron F, Lévesque H, et al. Antiphospholipid antibodies induce vascular functional changes in mice: a mechanism of vascular lesions in antiphospholipid syndrome? *Lupus* 2008;17:185-94.
- Porta C, Buggia I, Bonomi I, Caporali R, Scatola C, Montecucco C. Nitrite and nitrate plasma levels, as markers of nitric oxide synthesis, in antiphospholipid antibodies-related conditions and in thrombotic thrombocytopenic purpura. *Thromb Haemost* 1997;78:965-7.
- Ames PR, Tommasino C, Alves J, Morrow JD, Iannaccone L, Fossati G, et al. Antioxidant susceptibility of pathogenic pathways in subjects with antiphospholipid antibodies: a pilot study. *Lupus* 2000;9:688-95.
- Tan EM, Cohen AS, Fries JF, Masi AT, McShane DJ, Rothfield NF, et al. The 1982 revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum* 1982;25:1271-7.
- Greaves M, Cohen H, Machin SJ, Mackie I. Guidelines on the investigation and management of the antiphospholipid syndrome. *Br J Haematol* 2000;109:704-15.
- Tousoulis D, Boger RH, Antoniadou C, Siasos G, Stefanadi E, Stefanadis C. Mechanisms of disease: L-arginine in coronary atherosclerosis — a clinical perspective. *Nat Clin Pract Cardiovasc Med* 2007;4:274-83.
- Lauer T, Preik M, Rassaf T, Strauer BE, Deussen A, Feelisch M, et al. Plasma nitrite rather than nitrate reflects regional endothelial nitric oxide synthase activity but lacks intrinsic vasodilator action. *Proc Nat Acad Sci USA* 2001;98:12814-9.
- Kleimbongard P, Dejam A, Lauer T, Rassaf T, Schindler A, Picker O, et al. Plasma nitrite reflects constitutive nitric oxide synthase activity in mammals. *Free Radic Biol Med* 2003;35:790-6.
- Kleimbongard P, Dejam A, Lauer T, Jax T, Kerber S, Gharini P, et al. Plasma nitrite concentrations reflect the degree of endothelial dysfunction in humans. *Free Radic Biol Med* 2006;40:295-302.
- Stalic M, Poredos P, Peternel P, Tomsic M, Sebestjen M, Kveder T. Endothelial function is impaired in patients with primary antiphospholipid syndrome. *Thromb Res* 2006;118:455-61.
- Feletou M, Vanhoutte PM. Endothelial dysfunction: a multifaceted disorder (The Wiggers Award Lecture). *Am J Physiol Heart Circ Physiol* 2006;291:985-1002.
- Praticò D, Ferro D, Iuliano L, Rokach J, Conti F, Valesini G, et al. Ongoing prothrombotic state in patients with antiphospholipid antibodies: a role for increased lipid peroxidation. *Blood* 1999;93:3401-7.
- Atsumi T, Khamashta MA, Haworth RS, Brooks G, Amengual O, Ichikawa K, et al. Arterial disease and thrombosis in the antiphospholipid syndrome: a pathogenic role for endothelin 1. *Arthritis Rheum* 1998;41:800-7.
- Lowenstein CJ. Nitric oxide regulation of protein trafficking in the cardiovascular system. *Cardiovasc Res* 2007;15:75:240-6.
- Nathan C. Role of iNOS in human host defense. *Science* 2006;312:1874-5.
- Ames PR, Matsuura E, Batuca JR, Ciampa A, Lopez LL, Ferrara F, et al. High-density lipoprotein inversely relates to its specific autoantibody favoring oxidation in thrombotic primary antiphospholipid syndrome. *Lupus* 2010;19:711-6.
- Pham VV, Stichtenoth DO, Tsikas D. Nitrite correlates with 3-nitrotyrosine but not with the F(2)-isoprostane 15(S)-8-iso-PGF(2alpha) in urine of rheumatic patients. *Nitric Oxide* 2009;21:210-5.
- Ames PRJ, Antinolfi I, Ciampa A, Batuca J, Scenna G, Lopez LR, et al. Primary antiphospholipid syndrome: a low-grade auto inflammatory disease? *Rheumatology* 2008;47:1832-7.
- Basu S, Helmersson J, Jarosinska D, Sällsten G, Mazzolai B,

- Barregård L. Regulatory factors of basal F(2)-isoprostane formation: population, age, gender and smoking habits in humans. *Free Radic Res* 2009;43:85-91.
32. Ahmadzadehfar H, Oguogho A, Efthimiou Y, Kritz H, Sinzinger H. Passive cigarette smoking increases isoprostane formation. *Life Sci* 2006;78:894-7.
33. Tsikas D. Methods of quantitative analysis of the nitric oxide metabolites nitrite and nitrate in human biological fluids. *Free Radic Res* 2005;39:797-815.
34. Hingorani A. Polymorphisms in endothelial nitric oxide synthase and atherogenesis. John French Lecture. *Atherosclerosis* 2000;154:521-7.
35. Morris BJ, Glenn CL, Wilcken DE, Wang XL. Influence of an inducible nitric oxide synthase promoter variant on clinical variables in patients with coronary artery disease. *Clin Sci Lond* 2001;100:551-6.
36. Eleuteri E, Di Stefano A, Ricciardolo FL, Magno F, Gnemmi I, Colombo M, et al. Increased nitrotyrosine plasma levels in relation to systemic markers of inflammation and myeloperoxidase in chronic heart failure. *Int J Cardiol* 2009;135:386-90.
37. Sucu N, Unlü A, Tamer L, Aytacoglu B, Ercan B, Dikmengil M, et al. 3-Nitrotyrosine in atherosclerotic blood vessels. *Clin Chem Lab Med* 2003;41:23-5.

Ames PRJ, Batuca JR, Ciampa A, Iannaccone L, Delgado Alves J. Clinical relevance of nitric oxide metabolites and oxidative stress in thrombotic primary antiphospholipid syndrome. *J Rheumatol* 2010;37:2523-30. Table 2, under the heading "Predictors": "Arterial" should be "Arterial thrombosis". We regret the error.
doi:10.3899/jrheum.100494C1