

Platelet Reactivity and Disease Activity in Subjects with Psoriatic Arthritis

MATTEO NICOLA DARIO Di MINNO, SALVATORE IERVOLINO, ROSARIO PELUSO, RAFFAELE SCARPA, and GIOVANNI Di MINNO

ABSTRACT. *Objective.* Platelet aggregation plays a major role in vascular mortality. Individuals with psoriatic arthritis (PsA) are highly predisposed to vascular mortality. We evaluated the correlation between disease activity and platelet aggregation in individuals with PsA.

Methods. Individuals with PsA receiving tumor necrosis factor- α (TNF- α) blockers (n = 114) and healthy controls (n = 114) matched for age, sex, and cardiovascular risk factors were tested for light transmission aggregometry. None was receiving antiinflammatory drugs. Platelet aggregation (max-A%) was defined as maximal light transmittance achieved within 5 min after the addition of 0.1 or 0.2 mM arachidonic acid or 0.4 μ M adenosine diphosphate. A value of $\geq 50\%$ irreversible light transmittance (LT-50%) following platelet stimulation was used to define platelet hyperreactivity. Minimal disease activity (MDA) was evaluated in subjects with PsA.

Results. Regardless of the agent used, individuals with PsA showed a higher max-A% and achieved LT-50% more often than controls. Among individuals with PsA, those achieving MDA exhibited a max-A% similar to that of controls, both being significantly lower (p < 0.001) than max-A% of subjects with active disease. Subjects with active disease showed platelet hyperreactivity (LT-50%) more often than those achieving MDA (p < 0.001). For increasing quartiles of max-A%, C-reactive protein levels increased and prevalence of MDA decreased.

Conclusion. Compared with those achieving MDA, subjects with active PsA disease had abnormally high platelet reactivity. Whether this is relevant for the cardiovascular risk profile of subjects with PsA receiving TNF- α blockers requires further evaluation. (J Rheumatol First Release Dec 15 2011; doi:10.3899/jrheum.110741)

Key Indexing Terms:

PSORIATIC ARTHRITIS
TUMOR NECROSIS FACTOR- α BLOCKERS

PLATELET AGGREGATION
TREATMENT

Individuals with psoriatic arthritis (PsA) are predisposed to atherosclerotic vascular mortality¹. Platelets link atherosclerosis with inflammation². Little is known about disease activity and platelet aggregation in subjects with PsA receiving tumor necrosis factor- α (TNF- α) blockers.

MATERIALS AND METHODS

Among those referred to our outpatient clinic, 114 individuals with PsA (criteria of the CASPAR Study Group³) receiving TNF- α blockers for ≥ 1 year were enrolled in our study. Healthy individuals (n = 114) matched for age, sex, and cardiovascular risk factors⁴ served as controls. Exclusion criteria included a history of coronary artery disease, receiving active antiplatelet treatment [including nonsteroidal antiinflammatory drugs (NSAID)], serum

creatinine > 140 mmol/l, platelet count < 120,000/ $\times 10^9$ /l, and liver enzymes more than twice the upper limit of normal⁵.

The clinical subset (Moll and Wright criteria⁶) and the achievement of a sustained (> 1 year) minimal disease activity (MDA)⁷ were evaluated in each PsA subject.

The maximal light transmittance (Chrono-Log; Mascia Brunelli, Milan, Italy) achieved within 5 min after the addition of 0.1 and 0.2 mM arachidonic acid (AA) or of 0.4 μ M adenosine diphosphate (ADP) was defined as max-A%⁸. A value of $\geq 50\%$ irreversible light transmittance (LT-50%) defined the platelet hyperreactivity⁹.

Continuous data were expressed as means \pm SD, categorical variables as percentages. Continuous variables were compared by an independent sample T test and by ANOVA analysis. Correlations were assessed by Pearson's linear correlation coefficients (r). The chi-square test with Fisher's exact test was performed for categorical data.

RESULTS

Major characteristics of case and control subjects are reported in Table 1.

No subject with PsA was receiving traditional disease-modifying antirheumatic drugs. Adalimumab therapy was employed in 44 subjects (35.6%), etanercept and infliximab in 35 each (30.7%).

PsA subjects achieved LT-50% more often than controls following exposure to ADP 0.4 μ M (33.3% vs 14.0%, respectively; p = 0.001) or to AA 0.1 mM (33.3% vs 10.5%; p <

From the Department of Clinical and Experimental Medicine, the Reference Centre for Coagulation Disorders, and the Rheumatology Research Unit on Psoriatic Arthritis, Federico II University, Naples, Italy.

M.N.D. Di Minno, MD, Department of Clinical and Experimental Medicine, Reference Centre for Coagulation Disorders; S. Iervolino, MD; R. Peluso, MD, PhD; R. Scarpa, MD, Department of Clinical and Experimental Medicine, Rheumatology Research Unit on Psoriatic Arthritis; G. Di Minno, MD, Department of Clinical and Experimental Medicine, Reference Centre for Coagulation Disorders.

Address correspondence to Dr. M.N.D. Di Minno, Regional Reference Centre for Coagulation Disorders, Federico II University, Via S. Pansini 5, 80131, Naples, Italy. E-mail: dario.diminno@hotmail.it

Accepted for publication September 15, 2011.

Personal non-commercial use only. The Journal of Rheumatology Copyright © 2011. All rights reserved.

Table 1. Demographic and clinical features of the study population. Data are mean \pm SD unless otherwise specified.

Variable	PsA Subjects, n = 114	Controls, n = 114	p
Age, yrs	51.48 \pm 12.58	49.36 \pm 10.28	0.488
Male, n (%)	56 (49.1)	43 (37.7)	0.109
Metabolic syndrome, n (%)	38 (33.3)	33 (28.9)	0.567
IFG	11 (9.6)	10 (8.8)	1.000
Hypertension	46 (40.4)	40 (35.1)	0.495
Hypercholesterolemia with low HDL	59 (51.8)	63 (55.3)	0.690
Hypertriglyceridemia	30 (26.3)	29 (25.4)	1.000
Obesity	71 (62.3)	74 (64.9)	0.783
Smoking habit, n (%)	3 (28.9)	38 (33.3)	0.567
Disease duration, mo	98.11 \pm 54.12	—	—
Treatment duration, mo	48.11 \pm 16.41	—	—
Pattern of PsA, n (%)			
Distal joint disease	0 (0)	—	—
Oligoarthritis	26 (22.8)	—	—
Spondylitis	40 (35.1)	—	—
Polyarthritis	42 (36.8)	—	—
Mutilans	6 (5.3)	—	—

PsA: psoriatic arthritis; IFG: impaired fasting glucose; HDL: high-density lipoprotein.

0.001) or to AA 0.2 mM (68.4% vs 19.3%; $p < 0.001$). Max-A% was higher in PsA subjects than in controls as well (AA 0.1 mM: 54.50% \pm 16.10% vs 45.28% \pm 10.93%; $p < 0.001$; AA 0.2 mM: 68.14% \pm 18.45% vs 48.32% \pm 11.79%; $p < 0.001$; ADP 0.4 μ M: 51.15% \pm 20.68% vs 36.49% \pm 14.49%; $p < 0.001$).

Achievement of LT-50% in the 40 PsA subjects with MDA was comparable to that of controls and lower than that of subjects with active disease (Figure 1A). Similarly, max-A% values of those with MDA were comparable to controls and were lower than those of individuals with active disease (Figure 1B).

C-reactive protein (CRP) values were lower in subjects




Figure 1. Platelet aggregation in response to different aggregating agents in PsA. A. Achievement of LT-50% in the 40 PsA subjects with MDA was comparable to that of controls and lower than that of subjects with active disease. B. Max-A% values of those with MDA were comparable to controls and were lower than those of individuals with active disease. AA: arachidonic acid; ADP: adenosine diphosphate; LT-50%: irreversible light transmittance \geq 50%; MDA: minimal disease activity.

with MDA than in those with active disease (1.98 ± 1.8 vs 3.6 ± 3.9 ; $p = 0.007$). CRP values showed a direct correlation with max-A% (AA 0.1 mM: $r = 0.360$, $p < 0.001$; AA 0.2 mM: $r = 0.278$, $p = 0.003$; ADP 0.4 μ M: $r = 0.224$, $p = 0.018$) and with max-A% quartiles (AA 0.1 mM: $r = 0.446$, $p < 0.001$; AA 0.2 mM: $r = 0.216$, $p = 0.022$; ADP 0.4 μ M: $r = 0.209$, $p = 0.027$). The prevalence of MDA decreased for increasing quartiles of max-A% (Figure 2).

DISCUSSION

Our study shows that platelet function is increased in patients with PsA, especially in those with poorly controlled disease. The correlation of CRP with max-A% and the decreasing prevalence of MDA for increasing quartiles of max-A% argue for a link between inflammation and platelet reactivity. By interacting with specific platelet receptors, cytokines/chemokines involved in PsA¹⁰ cause intracellular calcium mobilization, nucleotide secretion, and platelet activation¹¹. Hyperreactivity to ADP has been reported in rheumatic diseases⁵. However, almost 50% of patients in that sample were receiving NSAID¹² and only 17% had PsA.

Platelet hyperreactivity was correlated with an elevated incidence of arterial thrombosis^{13,14}, and the effect of antiplatelet agents in the vascular risk profile of subjects with PsA requires investigation⁵. Our data show that inflammation influences platelet reactivity and that achievement of minimal disease activity may normalize platelet hyperreactivity.

ACKNOWLEDGMENT

The authors thank all members of the CaRRDs Study Group for collaboration and support in the literature search and the organization of the study; and Dr. Anna Lilia Ruocco and Dr. Fabrizio Giello, Department of Clinical and Experimental Medicine, Federico II University, for performing platelet aggregation tests.




Figure 2. Prevalence of MDA according to quartile of maximum aggregation in response to different agonists (AA 0.1 mM, AA 0.2 mM, ADP 0.4 μ M). AA: arachidonic acid; ADP: adenosine diphosphate; MDA: minimal disease activity.

REFERENCES

- Di Minno MN, Iervolino S, Peluso R, Scarpa R, Di Minno G; CaRRDs study group. Carotid intima-media thickness in psoriatic arthritis. Differences between tumor necrosis factor- α (TNF- α)-blockers and traditional disease modifying anti-rheumatic drugs. *Arterioscler Thromb Vasc Biol* 2011;31:705-12.
- Libby P, Ridker PM, Hansson GK; Leducq Transatlantic Network on Atherothrombosis. Inflammation in atherosclerosis: From pathophysiology to practice. *J Am Coll Cardiol* 2009;54:2129-38.
- Taylor W, Gladman D, Helliwell P, Marchesoni A, Mease P, Mielants H; CASPAR Study Group. Classification criteria for psoriatic arthritis: Development of new criteria from a large international study. *Arthritis Rheum* 2006;54:2665-73.
- Di Minno MN, Tufano A, Guida A, Di Capua M, De Gregorio AM, Cerbone AM, et al. Abnormally high prevalence of major components of the metabolic syndrome in subjects with early-onset idiopathic venous thromboembolism. *Thromb Res* 2011;127:193-7.
- MacMullan PA, Peace AJ, Madigan AM, Tedesco AF, Kenny D, McCarthy GM. Platelet hyper-reactivity in active inflammatory arthritis is unique to the adenosine diphosphate pathway: A novel finding and potential therapeutic target. *Rheumatology* 2010;49:240-5.
- Moll JM, Wright V. Psoriatic arthritis. *Semin Arthritis Rheum* 1973;3:55-78.
- Coates LC, Cook R, Lee KA, Chandran V, Gladman DD. Frequency, predictors, and prognosis of sustained minimal disease activity in an observational psoriatic arthritis cohort. *Arthritis Care Res* 2010;62:970-6.
- Brambilla M, Parolari A, Camera M, Colli S, Eligini S, Centenaro C, et al. Effect of two doses of aspirin on thromboxane biosynthesis and platelet function in patients undergoing coronary surgery. *Thromb Haemost* 2010;103:516-24.
- Yee DL, Sun CW, Bergeron AL, Dong JF, Bray PF. Aggregometry detects platelet hyperreactivity in healthy individuals. *Blood* 2005;106:2723-9.
- Loffredo S, Ayala F, Marone GC, Genovese A, Marone G. Immunopathogenesis of psoriasis and psoriatic arthritis and pharmacological perspectives. *Reumatismo* 2007;59 Suppl 1:28-39.
- Barrett NE, Holbrook L, Jones S, Kaiser WJ, Moraes LA, Rana R, et al. Future innovations in anti-platelet therapies. *Br J Pharmacol* 2008;154:918-39.
- Galliard-Grigioni KS, Reinhart WH. A randomized, controlled study on the influence of acetaminophen, diclofenac, or naproxen on aspirin-induced inhibition of platelet aggregation. *Eur J Pharmacol* 2009;609:96-9.
- Di Minno MN, Guida A, Camera M, Colli S, Di Minno G, Tremoli E. Overcoming limitations of current antiplatelet drugs. A concerted effort for more profitable strategies of intervention. *Ann Med* 2011;43:531-44.
- Bray PF. Platelet hyperreactivity: Predictive and intrinsic properties. *Hematol Oncol Clin North Am* 2007;21:633-45.