

Early Stage Spine Infarct Accurately Diagnosed by ^{99m}Tc -HMDP Bone Scintigraphy Performed on a Combined Single Photon Emission Computed Tomography/Computed Tomography System Correlation with Magnetic Resonance Imaging and Histopathological Findings

NICOLAS AIDE, MD; THIERRY FRANSON, MD, Nuclear Medicine Department; BRIGITTE MARIE, MD, Radiology Department; JACQUES CHASLE, MD, Pathology Department; FRANÇOIS LESAUNIER, MD, Radiation Oncology Department; STÉPHANE BARDET, MD, Nuclear Medicine Department, François Baclesse Comprehensive Cancer Center, Caen, France. Address reprint requests to Dr. N. Aide, Service de Médecine Nucléaire, Centre François Baclesse, Avenue du Général Harris, Caen, France. E-mail: n.aide@baclesse.fr. (J Rheumatol 2007;34:2121–2)

A 53-year-old man with no significant medical, drug, or trauma history was referred for bone scintigraphy following acute lumbar pain and suspicious magnetic resonance imaging (MRI) findings. MRI examination showed anomalies (Figure 1) that, although not specific, were consistent with neoplastic bone involvement. A ^{99m}Tc -HMDP single photon emission computed tomography (SPECT)/CT scan (Figure 2) showed a hypometabolic aspect of the whole corpus of L1, the posterior arch appearing normal. The CT part of the scan was normal, ruling out known causes of cold vertebrae such as osteolytic metastasis¹, plasmacytoma², and hemangioma³. The hypothesized diagnosis was early-stage bone infarct, with an increase in tracer uptake expected to be observed at a later stage, as in avascular necrosis of the femoral head⁴. However, due to the MRI results, a surgical biopsy was performed.

Pathological examination (Figure 3) was consistent with early bone necrosis. The lack of anomaly in the posterior part of the vertebrae in both MRI and bone scintigraphy results was consistent with knowledge on lumbar spine vascularization, the vertebral body and posterior arch comprising distinct entities⁵. Since our patient presented no risk factor for avascular bone necrosis, an idiopathic spine infarct was presumed⁶. After a 9-month followup period, persistent lumbar pain was noted, but no relapse was suspected.

In light of our report, one may assume that new hybrid SPECT/CT devices equipped with helical CT, which have already proven to be useful in the staging of cancer⁷, could also influence management of rheumatology patients.

REFERENCES

1. Kagan AR, Steckel RJ, Bassett LW. Diagnostic oncology case study: lytic spine lesion and cold bone scan. *AJR Am J Roentgenol* 1981;136:129-31.
2. Hartshorne MF, Cawthon MA, Bauman JM. Plasmacytoma of the lumbar spine by SPECT. *Clin Nucl Med* 1986;11:65-6.
3. Makhija M, Bofill ER. Hemangioma, a rare cause of photopenic lesion on skeletal imaging. *Clin Nucl Med* 1988;13:661-2.
4. Ryu JS, Kim JS, Moon DH, et al. Bone SPECT is more sensitive than MRI in the detection of early osteonecrosis of the femoral head after renal transplantation. *J Nucl Med* 2002;43:1006-11.
5. Chiras J, Morvan G, Merland J, Bories J. Blood supply to the thoracic (dorsal) and lumbar spine. *Surgical Radiologic Anatomy* 1982;4:23-31.
6. Lafforgue P. Pathophysiology and natural history of avascular necrosis of bone. *Joint Bone Spine* 2006;73:500-7.
7. Utsunomiya D, Shiraishi S, Imuta M, et al. Added value of SPECT/CT fusion in assessing suspected bone metastasis: comparison with scintigraphy alone and nonfused scintigraphy and CT. *Radiology* 2006;238:264-71.




Figure 1. MRI sagittal views centered on the lumbar spine. Signal intensity at L1 vertebral body is decreased on T1 weighted images (A), slightly increased on T2 weighted images (B), and markedly increased on T2 weighted images with fat saturation (C). No vertebral deformation or epidural anomaly is visible.




Figure 2. SPECT/CT sagittal (A, B, C) and transaxial (D, E, F) views. SPECT scans (A and D) show a hypometabolic aspect of the whole corpus of L1 (arrow). The posterior arch appears normal (arrowhead). The CT part of the SPECT/CT (B and E) scan is normal. Fused SPECT/CT (C and F) allows precise localization of scintigraphic anomalies.




Figure 3. Pathological examination of transpedicular biopsy shows trabecular bone with empty osteoplasts with neither osteoblastic nor osteoclastic activity. Trabeculars are normal in size. Peritrabecular tissue showed focal adiponecrosis. (A: original magnification $\times 4$; B: original magnification $\times 40$.)