

Ligands for Programmed Cell Death 1 Gene in Patients with Systemic Lupus Erythematosus

SHU-CHEN WANG, CHIA-HUI LIN, TSAN-TENG OU, CHENG-CHIN WU, WEN-CHAN TSAI, CHAUR-JONG HU, HONG-WEN LIU, and JENG-HSIEN YEN

ABSTRACT. Objective. To investigate the role of ligands for programmed cell death 1 (PD-L) in the pathogenesis of systemic lupus erythematosus (SLE).

Methods. One hundred sixty-four patients with SLE and 160 healthy controls were enrolled in our study. The *PD-L1* and *PD-L2* polymorphisms were determined by polymerase chain reaction (PCR)/direct sequencing or restriction fragment length polymorphism (RFLP)-PCR.

Results. The genotype distributions of *PD-L2* 47103 C/T polymorphisms in patients with SLE were significantly different from those of the controls ($p = 0.003$). The genotype frequency of *PD-L2* 47103 T/T, in comparison with 47103 C/C, was significantly increased in patients with SLE when compared with that of the controls (odds ratio 2.5, 95% confidence interval 1.4–4.4, $p = 0.001$). A similar finding could also be found in the allele frequency of *PD-L2* 47103 T (SLE vs control, OR 1.7, 95% CI 1.3–2.4, $p = 0.001$). There were no significant differences in the genotype and allele frequencies of *PD-L1* polymorphisms between the patients and controls.

Conclusion. *PD-L2* 47103 T may be associated with susceptibility to SLE in Taiwan. (First Release Mar 1 2007; J Rheumatol 2007;34:721–5)

Key Indexing Terms:

PROGRAMMED CELL DEATH 1 LIGAND
SYSTEMIC LUPUS ERYTHEMATOSUS

PROGRAMMED CELL DEATH 2 LIGAND
PROGRAMMED CELL DEATH 1

Systemic lupus erythematosus (SLE) is a prototype of autoimmune diseases, which frequently involves multiple organs. The detailed pathogenesis is still obscure. Many genes such as human leukocyte antigen, complement, FcR, mannose binding lectin, protein tyrosine phosphatase N22, cytokines, and chemokine genes have been described to be associated with the development of SLE^{1–5}.

Programmed cell death 1 (PD-1) is an immunoinhibitory receptor expressed by activated T cells, B cells, and myeloid cells⁶. The ligands for PD-1 (*PD-L1* and *PD-L2*, also known as B7-H1 and B7-DC) are type I transmembrane proteins

structurally related to the B7 family. They can be induced in monocytes, dendritic cells, endothelial cells, keratinocytes, and B cells^{6–10}. However, PD-L1 expression is different from that of PD-L2. PD-L1 is also expressed on activated T cells, placental trophoblasts, myocardial endothelium, and cortical thymic epithelial cells. In contrast, PD-L2 can also be presented on placental endothelium and medullary thymic epithelial cells¹⁰. The interactions of PD-1 with PD-L1 and PD-L2 result in the inhibition of T cell receptor-mediated lymphocyte proliferation and cytokine secretion, and they also inhibit CD28-mediated costimulation. The relative levels of inhibitory PD-L1 and costimulatory CD80/CD86 signals on antigen-presenting cells determine the extent of T cell activation and the threshold between tolerance and autoimmunity. Therefore, PD-L1 expression on nonlymphoid tissues and its interaction with PD-1 may determine the extent of immune responses at sites of inflammation⁶.

Recent studies using anti-PD-L1 monoclonal antibodies have suggested a role for PD-L1 in regulating autoimmune diseases. Blockade of the PD-L during experimental autoimmune encephalomyelitis (EAE) or diabetes exacerbates the diseases^{11,12}. PD-L1 blockade rapidly precipitated diabetes in prediabetic female non-obese diabetic (NOD) mice. PD-L2 blockade in animals also resulted in augmentation of EAE. These studies showed that PD-1–PD-L blockade was related to the development of autoimmune disease.

The polymorphisms in exons of *PD-L1* and *PD-L2* may result in amino acid substitution, structural changes, and expression of PD-L1 and PD-L2. The consequent interactions

From the Department of Laboratory Medicine and Division of Rheumatology, Department of Internal Medicine, Kaohsiung Medical University Hospital; College of Medicine and Graduate Institute of Medicine, College of Medicine, Kaohsiung Medical University, Kaohsiung; and the Department of Neurology, Taipei Medical University Hospital, Taipei, Taiwan.

S-C. Wang, MSc, Department of Laboratory Medicine; C-H. Lin, MSc; T-T. Ou, MD, Attending Physician; C-C. Wu, MD, Attending Physician, Division of Rheumatology, Department of Internal Medicine, Kaohsiung Medical University Hospital; W-C. Tsai, MD, PhD, Associate Professor; H-W. Liu, MD, Professor, Division of Rheumatology, Department of Internal Medicine and College of Medicine, Kaohsiung Medical University; C-J. Hu, MD, Assistant Professor, Department of Neurology, Taipei Medical University Hospital; J-H. Yen, MD, PhD, MSc, Professor, Division of Rheumatology, Department of Internal Medicine and Graduate Institute of Medicine, College of Medicine, Kaohsiung Medical University.

Address reprint requests to Dr. J-H. Yen, Division of Rheumatology, Department of Internal Medicine, Kaohsiung Medical University Hospital, No. 100 Zih-You 1st Road, Kaohsiung City 807, Taiwan.
E-mail: jehsy@kmu.edu.tw

Accepted for publication December 28, 2006.

Personal non-commercial use only. The Journal of Rheumatology Copyright © 2007. All rights reserved.

between PD-1 and its ligands may also be changed. Therefore, *PD-L1* and *PD-L2* polymorphisms may be related to the pathogenesis of autoimmune diseases. *PD-L1* polymorphisms may be associated with susceptibility to SLE¹³. Therefore, *PD-L1* and *PD-L2* polymorphisms may also be associated with susceptibility to SLE. Our purpose was to investigate the associations of *PD-L1* and *PD-L2* polymorphisms with susceptibility and clinical manifestations of SLE in Taiwan.

MATERIALS AND METHODS

One hundred sixty-four patients with SLE (148 female, 16 male) and 160 healthy controls (139 female, 21 male) were enrolled in our study. All patients and controls were Taiwanese. The diagnosis of SLE was according to the American College of Rheumatology (ACR) 1997 revised criteria for the classification of SLE¹⁴. Our study was approved by the Institutional Review Board of Kaohsiung Medical University Hospital.

There are 3 polymorphisms in the exons of *PD-L1* including *PD-L1* 1072 G/C (rs 12551333; exon 3, amino acid 49, Asp → His), 1113 G/A (rs 4278201; exon 3, amino acid 62, Lys → Lys, synonymous), and 6777 C/G (rs 17718883; exon 4, amino acid 146, Pro → Arg). The polymorphisms of *PD-L1* 1072 C/G and 1113 A/G were determined by polymerase chain reaction (PCR)/direct sequencing. The sequences of primers are shown in Table 1. PCR was performed under the following conditions: initial denaturation at 96°C for 3 min and 5 cycles of denaturation at 95°C for 1 min, annealing at 56°C for 1 min, and extension at 72°C for 1 min, and then 35 cycles of denaturation at 95°C for 1 min, annealing at 52°C for 1 min, and extension at 72°C for 1 min. A final extension phase was also performed at 72°C for 7 min. Then the nucleotide sequence was determined by the method of direct sequencing with the BigDye Terminator Cycle Sequencing Kit (Applied Biosystems). The polymorphisms of *PD-L1* 6777 C/G were determined by the restriction fragment length polymorphism (RFLP)-PCR method. The sequences of primers and restriction enzymes are shown in Table 1. The amplification conditions consisted of initial denaturation at 95°C for 3 min, followed by 5 cycles of denaturation at 94°C for 1 min, annealing at 55°C for 1 min, and extension at 72°C for 1 min, and 30 cycles of denaturation at 94°C for 1 min, annealing at 50°C for 1 min, and extension at 72°C for 1 min, and then a final extension phase at 72°C for 7 min. The restriction enzyme Bsr I was used to determine the 6777 C/G polymorphisms.

There are 3 non-synonymous polymorphisms in *PD-L2* including *PD-L2* 24293 G/C (rs 12339171; exon 3, amino acid 58, Ser → Thr), 47103 C/T (rs 7854303; exon 5, amino acid 229, Ser → Phe), and 47139 C/T (rs 7854413; exon 5, amino acid 241, Thr → Ile). These polymorphisms were also determined by RFLP-PCR. The sequences of primers and restriction enzymes are shown in Table 1. To determine the *PD-L2* 24293 G/C polymorphisms, PCR

was carried out under the following conditions: initial denaturation at 95°C for 3 min and 5 cycles of denaturation at 95°C for 1 min, annealing at 56°C for 1 min, and extension at 72°C for 1 min, and then 35 cycles of denaturation at 95°C for 1 min, annealing at 54°C for 1 min, and extension at 72°C for 1 min. A final extension phase was also performed at 72°C for 7 min. Then the PCR product was digested with Bsr I.

A mismatched nucleotide (underlined) was used in the downstream primer to determine 47103 C/T and 47139 C/T. The amplification conditions consisted of initial denaturation at 96°C for 3 min, followed by 5 cycles of denaturation at 95°C for 1 min, annealing at 56°C for 1 min, and extension at 72°C for 1 min, and 30 cycles of denaturation at 95°C for 1 min, annealing at 54°C for 1 min, and extension at 72°C for 1 min, and then a final extension phase at 72°C for 7 min. The restriction enzymes Bcc I and Tsp45 I were used to determine the 47103 C/T and 47139 C/T polymorphisms, respectively. The *PD-L2* 47103 C/T and 47139 C/T polymorphisms were also confirmed in some cases by direct sequencing with the BigDye Terminator Cycle Sequencing Kit. The sequences of primers were 5'- CCT GTT GGT CTA CCT CTT AG- 3' and 5'- TGA AAG CAG CAA GCC ATA GG- 3'. The amplification conditions consisted of initial denaturation at 96°C for 3 min, and 30 cycles of denaturation at 95°C for 1 min, annealing at 55°C for 1 min, and extension at 72°C for 1 min, and then a final extension phase at 72°C for 5 min.

The clinical manifestations of SLE were defined according to the 1997 revised ACR criteria. The severity of SLE was determined according to the SLICC/ACR damage index for SLE¹⁵.

The chi-square test (with Yates correction for 2 × 2 table) or Fisher's exact test was used for statistical analysis. Odds ratios (OR) and 95% confidence intervals (95% CI) were calculated using the SPSS program.

RESULTS

The distributions of *PD-L1* and *PD-L2* genotypes were compatible with Hardy-Weinberg equilibrium in the controls. The *PD-L1* 1072 C and *PD-L1* 1113 A alleles could not be detected in Taiwanese. Moreover, there was no significant difference in the genotype frequency of *PD-L1* 6777 C/G polymorphisms between the patients with SLE and controls (Table 2). Similar findings could also be found regarding their allele frequencies.

PD-L2 24293 C allele could not be found in Taiwanese patients and controls in our study. All of the patients and controls were *PD-L2* 24293 G. The study revealed that the genotype distributions of *PD-L2* 47103 C/T polymorphisms in the patients with SLE were significantly different from those of

Table 1. The sequences of primers and detection methods in determining *PD-L1* and *PD-L2* polymorphisms.

Polymorphisms	Primers	Detection Methods or Restriction Enzymes
<i>PD-L1</i>		
1072 G/C and 1113 G/A	5'-TGT GGT AGA GTA TGG TAG C-3' 5'-CTG TCT GTA GCT ACT ATG C-3'	Direct sequencing
6777 C/G	5'-TAC GTA GTT CTG TGC TCA G-3' 5'-GTT GAT TCT CAG TGT GCT G-3'	Bsr I
<i>PD-L2</i>		
24293 G/C	5'-AGC ATG GCA GCA ATG TGA C-3' 5'-CAC TCA CCT TTG ACT TTC AG-3'	Bsr I
47103 C/T	5'-GCT TCA CAT TTT CAT CCC <u>AT</u> -3' 5'-AGT GGC TCA TGT GCA GAC-3'	Bcc I
47139 C/T	5'-GCT TCA CAT TTT CAT CCC <u>AT</u> -3' 5'-AGT GGC TCA TGT GCA GAC-3'	Tsp45 I

Table 2. Frequencies of *PD-L1* polymorphisms in patients with SLE and controls. The *PD-L1* 1072 C and *PD-L1* 1113 A alleles could not be detected in Taiwanese. There were no significant differences in the genotype and allele frequencies of *PD-L1* 6777 polymorphisms between the patients with SLE and controls.

Polymorphisms of <i>PD-L1</i>	SLE, n = 164 (%)	Controls, n = 160 (%)
Genotype		
1072 G/G	164 (100)	160 (100)
1113 G/G	164 (100)	160 (100)
6777 C/C	124 (75.6)	127 (79.4)
C/G	39 (23.8)	31 (19.4)
G/G	1 (0.6)	2 (1.2)
Allele		
1072 G	328 (100)	320 (100)
1113 G	328 (100)	320 (100)
6777 C	287 (87.5)	285 (89.1)
G	41 (12.5)	35 (10.9)

the controls ($p = 0.003$; Table 3). The genotype frequency of *PD-L2* 47103 T/T, in comparison with 47103 C/C, was significantly higher in the patients with SLE than that of the controls (OR 2.5, 95% CI 1.4–4.4, $p = 0.001$). The allele frequency of *PD-L2* 47103 T was also significantly increased in patients with SLE compared with controls (OR 1.7, 95% CI 1.3–2.4, $p = 0.001$). However, there were no significant differences in the genotype and allele frequencies of *PD-L2* 47139 T/C polymorphisms between the patients with SLE and controls.

Our study also revealed that *PD-L1* and *PD-L2* polymorphisms were not associated with the clinical manifestations of SLE including nephritis, central nervous system involvement, cutaneous vasculitis, and various autoantibodies such as the antinuclear antibody, anti-dsDNA, anti-Sm, and anti-RNP. The *PD-L1* and *PD-L2* polymorphisms were also not associated with the SLICC/ACR damage index for SLE (data not shown).

DISCUSSION

Our study showed that *PD-L2* 47103 T was associated with susceptibility to SLE in Taiwan.

PD-L1 is constitutively expressed in T cells, B cells, macrophages, and dendritic cells, and is upregulated following activation of these cells. *PD-L1* is also expressed on non-lymphoid cells including cardiac endothelial cells, pancreatic islet cells, brain glial cells, and muscle cells. *PD-L2* expression is observed on activated macrophages and dendritic cells^{16,17}. Moreover, *PD-L1* and *PD-L2* are expressed in various tumor cells, such as ovary, esophagus, kidney, and brain tumors^{18–24}.

The *PD-1*–*PD-L* pathway regulates the immune response in both lymphoid and nonlymphoid organs. The interactions between *PD-1* and *PD-L* inhibit lymphocyte activation. Although positive effects of *PD-1*–*PD-L* interaction have been found^{8,9,25,26}, the negative effects are well documented. It is still unknown whether the positive effects are caused by the inhibition of negative signaling or by other stimulatory receptors.

Mice deficient in *PD-1* developed spontaneous autoimmune diseases, which suggested a negative costimulatory function. C57 BL/6-*Pdcd1*^{−/−} mice developed lupus-like glomerulonephritis and arthritis²⁷. BALB/c-*Pdcd1*^{−/−} mice developed dilated cardiomyopathy with production of autoantibodies against cardiac troponin I^{28–30}. In 2C-*Pdcd1*^{−/−} H-2^{bi/d} mice, the mice died of the graft-versus-host-like disease²⁷. Therefore, the *PD-1*–*PD-L* pathway may play a role in the induction and maintenance of peripheral tolerance. *PD-1*–*PD-L* interaction inhibits adverse immune response in 2 ways. *PD-L* on antigen-presenting cells inhibits T cell activation and induces peripheral tolerance. Moreover, *PD-L* on target cells inhibits the effector function of T cells to maintain tolerance.

PD-L on nonlymphoid organs prevents tissue destruction by suppressing the effector function of autoreactive lymphocytes. *PD-L1* on islet cells suppresses the effector function of diabetogenic T cells³¹. Moreover, antibody blockade of the

Table 3. Frequencies of *PD-L2* polymorphisms in patients with SLE and controls. The *PD-L2* 24293 C allele could not be found in this study.

Polymorphisms of <i>PD-L2</i>	SLE, n = 164 (%)	Controls, n = 160 (%)	p (for overall genotype frequency)	p (for individual genotype and allele frequency)	OR (95% CI)
Genotype					
24293 G/G	164 (100)	160 (100)	NS		
47103 C/C	63 (38.4)	77 (48.1)	0.003		1
C/T	46 (28.0)	56 (35.0)			
T/T	55 (33.5)	27 (16.9)		0.001	2.5 (1.4–4.4)
47139 T/T	151 (92.1)	144 (90.0)	NS		
C/T	13 (7.9)	15 (9.4)			
C/C	0 (0)	1 (0.6)			
Allele					
24293 G	328 (100)	320 (100)	NS		
47103 C	172 (52.4)	210 (65.6)			
T	156 (47.6)	110 (34.4)		0.001	1.7 (1.3–2.4)
47139 T	315 (96.0)	303 (94.7)	NS		
C	13 (4.0)	17 (5.3)			

PD-1–PD-L pathway in prediabetic NOD mice also induced type I diabetes¹². Wang, *et al* revealed that NOD-Pdcd1^{-/-} mice also developed type I diabetes³². It can therefore be concluded that PD-1–PD-L interaction may play a role in preventing the development of diabetes. PD-L1 on tumor cells suppresses the cytolytic activity of CD8+ T cells^{33,34}. Blocking PD-1–PD-L interaction will accelerate tumor eradication. A correlation between PD-L expression on tumor cells and poor prognosis has been found for human cancer patients^{20,23}. Therefore a PD-1–PD-L blockade may activate the immune system of tumor-bearing hosts to eradicate tumors.

PD-L1 on antigen-presenting cells results in inactivation of T cells. Resting dendritic cells induce inactivation or anergy of T cells and CD8+ T cell tolerance through the PD-1–PD-L pathway³⁵. Virus- or parasite-infected cells will induce PD-L1 expression on dendritic cells, and then induce T cell anergy, which will result in immune paralysis against viruses or parasites^{36,37}. However, activated dendritic cells express lower levels of PD-L than costimulatory and MHC molecules. The activating signals overcome the inhibitory signal of PD-1, and T cell activation will result.

PD-1–PD-L interaction inhibits adverse immune responses to prevent autoimmunity. The *PD-1* or *PD-L* polymorphisms may interfere with the interaction between PD-1 and PD-L, and then diminish the prevention of autoimmune response. Some studies revealed that *PD-1* polymorphisms were associated with several immune-mediated diseases including rheumatoid arthritis, SLE, and type I diabetes^{13,38-41}. Therefore *PD-L* polymorphisms may also be related to the development of autoimmune diseases. A report in regard to the association of *PD-L* polymorphisms with SLE is still unavailable. Our study showed that the polymorphisms in the exons of *PD-L1* were not associated with susceptibility to SLE. However, the *PD-L2* 47103 T was related to the development of SLE. *PD-L2* 47103 C/T are non-synonymous polymorphisms. *PD-L2* 47103 C encodes an amino acid serine in the transmembrane region. However, *PD-L2* 47103 T encodes amino acid phenylalanine, which contains an aromatic ring. The expression of protein molecule on the cell surface may be regulated by its transmembrane domain⁴². Transmembrane domain also plays a role in dimerization or oligomerization for surface molecules^{43,44}. Similar to a mutation in CD80, the amino acid substitution of PD-L2 may result in PD-L2 molecule expression on cell surface in small clusters⁴⁵. The small clusters of PD-L2 may be insufficient to initiate PD-1-mediated inhibitory signals, which indicates that the inhibiting effect of the PD-1–PD-L pathway on autoimmunity will be decreased.

The frequencies of *PD-L2* 47103 C/T and 47139 T/C polymorphisms in Taiwanese were different from those of the Han Chinese and Japanese (NCBI Web resource: www.ncbi.nlm.nih.gov/SNP/). This discrepancy may be due to different genetic backgrounds. The associations of *PD-1*

polymorphisms with lupus nephritis have been demonstrated in Sweden and Denmark^{46,47}, but not in the United States. However, a similar finding could not be found in *PD-L2* polymorphisms, which encode a ligand of PD-1. This may be due to different target genes and ethnic difference. Our study also revealed that the *PD-L1* and *PD-L2* polymorphisms were not associated with the severity of SLE, which was evaluated by the SLICC/ACR damage index.

PD-L2 47103 T may be associated with susceptibility to SLE in Taiwan. However, *PD-L1* polymorphisms are not related to the development of SLE.

REFERENCES

1. Tsao BP. Update on human systemic lupus erythematosus genetics. *Curr Opin Rheumatol* 2004;16:513-21.
2. Shen N, Tsao BP. Current advances in the human lupus genetics. *Curr Rheumatol Rep* 2004;6:391-8.
3. Prokunina L, Alarcon-Riquelme M. The genetic basis of systemic lupus erythematosus — knowledge of today and thoughts for tomorrow. *Hum Mol Genet* 2004;13 Spec No 1:R143-8.
4. Moser KL, Neas BR, Salmon JE, et al. Genome scan of human systemic lupus erythematosus: evidence for linkage on chromosome 1q in African-American pedigrees. *Proc Natl Acad Sci USA* 1998;95:14869-74.
5. Croker JA, Kimberly RP. Genetics of susceptibility and severity in systemic lupus erythematosus. *Curr Opin Rheumatol* 2005;17:529-37.
6. Freeman GJ, Long AJ, Iwai Y, et al. Engagement of the PD-1 immunoinhibitory receptor by a novel B7 family member leads to negative regulation of lymphocyte activation. *J Exp Med* 2000;192:1027-34.
7. Latchman Y, Wood CR, Chernova T, et al. PD-L2 is a second ligand for PD-1 and inhibits T cell activation. *Nat Immunol* 2001;2:261-8.
8. Dong H, Zhu G, Tamada K, Chen L. B7-H1, a third member of the B7 family, co-stimulates T-cell proliferation and interleukin-10 secretion. *Nat Med* 1999;5:1365-9.
9. Tseng SY, Otsuji M, Gorski K, et al. B7-DC, a new dendritic cell molecule with potent costimulatory properties for T cells. *J Exp Med* 2001;193:839-46.
10. Brown JA, Dorfman DM, Ma FR, et al. Blockade of programmed death-1 ligands on dendritic cells enhances T cell activation and cytokine production. *J Immunol* 2003;170:1257-66.
11. Salama AD, Chitnis T, Imitola J, et al. Critical role of the programmed death-1 (PD-1) pathway in regulation of experimental autoimmune encephalomyelitis. *J Exp Med* 2003;198:71-8.
12. Ansari MJ, Salama AD, Chitnis T, et al. The programmed death-1 (PD-1) pathway regulates autoimmune diabetes in nonobese diabetic (NOD) mice. *J Exp Med* 2003;198:63-9.
13. Prokunina L, Castillejo-Lopez C, Oberg F, et al. A regulatory polymorphism in PDCD1 is associated with susceptibility to systemic lupus erythematosus in humans. *Nat Genet* 2002;32:666-9.
14. Hochberg MC. Updating the American College of Rheumatology revised criteria for the classification of systemic lupus erythematosus [letter]. *Arthritis Rheum* 1997;40:1725.
15. Gladman D, Ginzler E, Goldsmith C, et al. The development and initial validation of the Systemic Lupus International Collaborating Clinics/American College of Rheumatology damage index for systemic lupus erythematosus. *Arthritis Rheum* 1996;39:363-9.
16. Yamazaki T, Akiba H, Iwai H, et al. Expression of programmed death 1 ligands by murine T cells and APC. *J Immunol* 2002;169:5538-45.

17. Ishida M, Iwai Y, Tanaka Y, et al. Differential expression of PD-L1 and PD-L2, ligands for an inhibitory receptor PD-1, in the cells of lymphohematopoietic tissues. *Immunol Lett* 2002;84:57-62.
18. Curiel TJ, Wei S, Dong H, et al. Blockade of B7-H1 improves myeloid dendritic cell-mediated antitumor immunity. *Nat Med* 2003;9:562-7.
19. Dong H, Strome SE, Salomao DR, et al. Tumor-associated B7-H1 promotes T-cell apoptosis: a potential mechanism of immune evasion. *Nat Med* 2002;8:793-800.
20. Ohigashi Y, Sho M, Yamada Y, et al. Clinical significance of programmed death-1 ligand-1 and programmed death-1 ligand-2 expression in human esophageal cancer. *Clin Cancer Res* 2005;11:2947-53.
21. Konishi J, Yamazaki K, Azuma M, Kinoshita I, Dosaka-Akita H, Nishimura M. B7-H1 expression on non-small cell lung cancer cells and its relationship with tumor-infiltrating lymphocytes and their PD-1 expression. *Clin Cancer Res* 2004;10:5094-100.
22. Strome SE, Dong H, Tamura H, et al. B7-H1 blockade augments adoptive T-cell immunotherapy for squamous cell carcinoma. *Cancer Res* 2003;63:6501-5.
23. Thompson RH, Gillett MD, Cheville JC, et al. Costimulatory B7-H1 in renal cell carcinoma patients: Indicator of tumor aggressiveness and potential therapeutic target. *Proc Natl Acad Sci USA* 2004;101:17174-9.
24. Wintterle S, Schreiner B, Mitsdoerffer M, et al. Expression of the B7-related molecule B7-H1 by glioma cells: a potential mechanism of immune paralysis. *Cancer Res* 2003;63:7462-7.
25. Subudhi SK, Zhou P, Yerian LM, et al. Local expression of B7-H1 promotes organ-specific autoimmunity and transplant rejection. *J Clin Invest* 2004;113:694-700.
26. Shin T, Yoshimura K, Shin T, et al. In vivo costimulatory role of B7-DC in tuning T helper cell 1 and cytotoxic T lymphocyte responses. *J Exp Med* 2005;201:1531-41.
27. Nishimura H, Nose M, Hiai H, Minato N, Honjo T. Development of lupus-like autoimmune diseases by disruption of the PD-1 gene encoding an ITIM motif-carrying immunoreceptor. *Immunity* 1999;11:141-51.
28. Nishimura H, Okazaki T, Tanaka Y, et al. Autoimmune dilated cardiomyopathy in PD-1 receptor-deficient mice. *Science* 2001;291:319-22.
29. Okazaki T, Tanaka Y, Nishio R, et al. Autoantibodies against cardiac troponin I are responsible for dilated cardiomyopathy in PD-1-deficient mice. *Nat Med* 2003;9:1477-83.
30. Okazaki T, Honjo T. Pathogenic roles of cardiac autoantibodies in dilated cardiomyopathy. *Trends Mol Med* 2005;11:322-6.
31. Okazaki T, Honjo T. The PD-1-PD-L pathway in immunological tolerance. *Trends Immunol* 2006;27:195-201. Epub 2006 Feb 24.
32. Wang J, Yoshida T, Nakaki F, Hiai H, Okazaki T, Honjo T. Establishment of NOD-Pdcd1^{-/-} mice as an efficient animal model of type I diabetes. *Proc Natl Acad Sci USA* 2005;102:11823-8.
33. Iwai Y, Ishida M, Tanaka Y, Okazaki T, Honjo T, Minato N. Involvement of PD-L1 on tumor cells in the escape from host immune system and tumor immunotherapy by PD-L1 blockade. *Proc Natl Acad Sci USA* 2002;99:12293-7.
34. Hirano F, Kaneko K, Tamura H, et al. Blockade of B7-H1 and PD-1 by monoclonal antibodies potentiates cancer therapeutic immunity. *Cancer Res* 2005;65:1089-96.
35. Probst HC, McCoy K, Okazaki T, Honjo T, van den Broek M. Resting dendritic cells induce peripheral CD8⁺ T cell tolerance through PD-1 and CTLA-4. *Nat Immunol* 2005;6:280-6.
36. Kirchberger S, Majdic O, Steinberger P, et al. Human rhinoviruses inhibit the accessory function of dendritic cells by inducing sialoadhesion and B7-H1 expression. *J Immunol* 2005;175:1145-52.
37. Smith P, Walsh CM, Mangan NE, et al. Schistosoma mansoni worms induce anergy of T cells via selective up-regulation of programmed death ligand 1 on macrophages. *J Immunol* 2004;173:1240-8.
38. Nielsen C, Hansen D, Husby S, Jacobsen BB, Lillevang ST. Association of a putative regulatory polymorphism in the PD-1 gene with susceptibility to type 1 diabetes. *Tissue Antigens* 2003;62:492-7.
39. Ferreira-Vidal I, Gomez-Reino JJ, Barros F, et al. Association of PDCD1 with susceptibility to systemic lupus erythematosus: evidence of population-specific effects. *Arthritis Rheum* 2004;50:2590-7.
40. Selenko-Gebauer N, Majdic O, Szekeres A, et al. B7-H1 (programmed death-1 ligand) on dendritic cells is involved in the induction and maintenance of T cell anergy. *J Immunol* 2003;170:3637-44.
41. Lin SC, Yen JH, Tsai JJ, et al. Association of a programmed death 1 gene polymorphism with the development of rheumatoid arthritis, but not systemic lupus erythematosus. *Arthritis Rheum* 2004;50:770-5.
42. Chou WC, Liao KW, Lo YC, Jiang SY, Yeh MY, Roffler SR. Expression of chimeric monomer and dimer proteins on the plasma membrane of mammalian cells. *Biotechnol Bioeng* 1999;65:160-9.
43. Choi S, Lee E, Kwon S, et al. Transmembrane domain-induced oligomerization is crucial for the functions of syndecan-2 and syndecan-4. *J Biol Chem* 2005;280:42573-9.
44. Bourguignon LY, Singer SJ. Transmembrane interactions and the mechanism of capping of surface receptors by their specific ligands. *Proc Natl Acad Sci USA* 1977;74:5031-5.
45. Doty RT, Clark EA. Two regions in the CD80 cytoplasmic tail regulate CD80 redistribution and T cell costimulation. *J Immunol* 1998;161:2700-7.
46. Prokunina L, Gunnarsson I, Sturfelt G, et al. The systemic lupus erythematosus-associated PDCD1 polymorphism PD1.3A in lupus nephritis. *Arthritis Rheum* 2004;50:327-8.
47. Nielsen C, Laustrop H, Voss A, Junker P, Husby S, Lillevang ST. A putative regulatory polymorphism in PD-1 is associated with nephropathy in a population-based cohort of systemic lupus erythematosus patients. *Lupus* 2004;13:510-6.