

Bone Scanning of Limited Value for Diagnosis of Symptomatic Oligofocal and Multifocal Osteonecrosis

MICHAEL A. MONT, SLIF D. ULRICH, THORSTEN M. SEYLER, JONATHAN M. SMITH, DAVID R. MARKER, MIKE S. McGRATH, DAVID S. HUNGERFORD, and LYNNE C. JONES

ABSTRACT. *Objective.* Bone scintigraphy has been advocated as a useful diagnostic tool for patients with suspected osteonecrosis or in screening for multifocal disease. We evaluated the sensitivity of bone scanning relative to magnetic resonance imaging (MRI) in the diagnosis of osteonecrosis.

Methods. Forty-eight patients presented with suspected osteonecrosis of the shoulder, hip, knee, or ankle. All patients underwent simultaneous (< 3 months apart) bone scans and MRI studies as part of diagnostic investigations. Histological confirmation of osteonecrosis was obtained for all suspected lesions. The diagnostic result for each imaging modality was then assessed and compared.

Results. All 163 (100%) histologically confirmed lesions were identified by MRI, while only 91 lesions (56%) were identified by bone scan. There was complete congruency of bone scans with MR images in only 38% of patients (18/48). Bone scanning identified 72% of lesions (47/65) in oligofocal patients (≤ 2 joints involved) compared with 45% of the lesions (44/98) in multifocal patients (≥ 3 joints involved). Sensitivity of lesions was highest for the knee and hip and lower for the shoulder and ankle. Larger and later-stage lesions had a higher bone scan sensitivity.

Conclusion. Our results demonstrated the low sensitivity of bone scintigraphy for diagnosing symptomatic osteonecrosis. It is least sensitive for early-stage lesions where it might be most useful to diagnose the disease. Our study also confirms that this test is less sensitive for joints other than the hip and is also not useful as a screening tool. Our study does not support the use of bone scans as a diagnostic or screening tool for osteonecrosis. (First Release June 1 2008; J Rheumatol 2008; 35:1629–34)

Key Indexing Terms:

OSTEONECROSIS BONE SCAN MAGNETIC RESONANCE IMAGING
AVASCULAR NECROSIS OUTCOME DIAGNOSIS

Osteonecrosis is a devastating disease that typically afflicts a young patient population. The disease can affect various joints such as the knee, shoulder, and ankle, although it most commonly involves the hip. Radiographs are often initially used to diagnose osteonecrosis. If disease is apparent radiographically, then often no other imaging or screening tests are necessary. However, disease often may not be detectable in early stages because radiographic abnormalities appear only after prolonged changes in ischemic tissue. As a result, other diagnostic methods including bone marrow pressure

measurements, intramedullary venography, and biopsies have been utilized for detecting early ischemic changes in bone¹. As imaging technology developed, investigators recorded varying degrees of success at detecting osteonecrosis with multiple imaging techniques, including bone scanning, computed tomography (CT), single-photon emission CT, and magnetic resonance imaging (MRI)²⁻⁵. These additional tests are needed to investigate early changes of the contralateral hip or to investigate other symptomatic joints. If these joints can be diagnosed early, there is the best chance for treatment aimed at saving the bone from collapse and requiring joint replacement.

Bone scanning has been advocated as a useful diagnostic tool for patients with suspected osteonecrosis of the femoral head. Before the advent and utilization of MRI, scanning was believed to be a more sensitive diagnostic test than standard radiographs for early detection of the disease⁶⁻¹⁴. MRI has been shown in multiple studies to be more sensitive than bone scanning for detection of osteonecrosis, some authors reporting 99% or higher sensitivity for this disease^{2,15-18}. Despite this, many practitioners still use bone scans as an initial screening modality for detection of symptomatic or asymptomatic disease. Additionally, various authors have advocated the use of bone scanning for early detection of

From the Rubin Institute for Advanced Orthopedics, Center for Joint Preservation and Reconstruction, Sinai Hospital of Baltimore; Good Samaritan Hospital, Orthopaedics Division, Johns Hopkins Medical Institution, Baltimore, Maryland, USA.

M.A. Mont, MD, Director; J.M. Smith, BS, Graduate Student; D.R. Marker, BS, Graduate Student; M.S. McGrath, MD, Fellow; T.M. Seyler, MD, Fellow; S.D. Ulrich, MD, Fellow, Center for Joint Preservation and Reconstruction, Sinai Hospital of Baltimore; L.C. Jones, PhD, Professor; D.R. Hungerford, MD, Professor, Good Samaritan Hospital, Orthopaedics Division, Johns Hopkins Medical Institution.

Dr. Mont and Dr. Ulrich contributed equally to this work.

Address reprint requests to Dr. M.A. Mont, Rubin Institute of Advanced Orthopedics, Center for Joint Preservation and Reconstruction, Sinai Hospital of Baltimore, 2401 West Belvedere Avenue, Baltimore, MD 21215. E-mail: Sulrich@lifebridgehealth.org

Accepted for publication March 12, 2008.

Personal non-commercial use only. The Journal of Rheumatology Copyright © 2008. All rights reserved.

osteonecrosis of other joints or to screen for multifocal disease^{3,9,11,19,20}.

In clinical practice, we found that bone scintigraphy often missed lesions that were diagnosed on radiographs and MR images. This led us to investigate the utility of bone scanning in diagnosis or screening of osteonecrosis. Specifically, we compared the diagnostic sensitivity of bone scanning with that of MRI in patients presenting with suspected atraumatic osteonecrosis in various joints who also had confirmed osteonecrosis on bone histological evaluation after surgical procedures.

MATERIALS AND METHODS

Study design and inclusion criteria. Between January 1, 1994, and November 23, 2005, 48 patients presenting to 2 institutions who underwent simultaneous (i.e., within 3 months) radiographs, bone scans, and MRI studies as part of diagnostic investigation for possible symptomatic osteonecrosis of the hip, knee, shoulder, or ankle were included in the study. There were 33 women and 15 men, mean age 39 years (range 20–76 yrs). All patients had histological confirmation of osteonecrosis from tissue obtained by various surgical procedures (core decompression, bone grafting, joint replacement) within 3 months of the imaging studies as well. We retrospectively assessed the diagnostic result for bone scanning and compared it to MRI results.

Patients were selected for diagnostic imaging and inclusion in the study based on a number of inclusion and exclusion criteria. The patient had to be at least 18 years of age and have had deep joint pain associated with either 2 or more known risk factors for osteonecrosis or prior radiographic evidence of osteonecrosis in at least one other joint. Patients who had trauma-associated lesions were excluded. Patients were also excluded if all diagnostic imaging was not completed within a 3-month period from the date of the first test procedure. Patients with endstage osteoarthritis (Ficat and Arlet; stage IV) with femoral head collapse and acetabular involvement were not included. Patients were later stratified into 2 cohorts: oligofocal osteonecrosis (≤ 2 joints involved) and multifocal osteonecrosis (≥ 3 joints involved).

From the outset, we did not use positive MRI scans as the key to the diagnosis. Seventeen of the patients had MRI scans (to delineate locations of lesions) after the diagnosis had been made with radiographs, bone scans, and clinical measures. Since we found that all lesions had a positive MRI (no false-positives), we retrospectively reanalyzed the cases to determine if there was an MRI bias that would clarify them as cases of osteonecrosis (positive radiographs, clinical factors, bone scans, histological confirmation). Therefore, for the initial selection of patients and further analysis, we do not believe there was a selection bias for only MRI-positive patients.

A clinical and radiographic review was performed to assess whether any variables were associated with positive bone scans. Hospital outpatient records were analyzed with respect to demographic data, including patient's age at presentation, sex, associated diseases, and other factors known to increase the risk of osteonecrosis such as alcohol abuse (> 400 ml of 100% absolute ethanol per week), tobacco use, and use of prednisone (or equivalent other corticosteroid medication) in doses exceeding 2 g^{3,21-25}. Associated diagnoses included systemic lupus erythematosus (SLE) in 10 patients, inflammatory bowel disease in 3, breast carcinoma in 2, connective tissue disorders in 2, renal disease in 3, and one patient each with idiopathic thrombocytopenic purpura, arteriovenous malformation, Hodgkin's lymphoma, T-cell acute lymphoblastic leukemia, and asthma. Nine patients had a history of alcohol abuse and 8 patients smoked more than 1 pack of cigarettes per day.

Radiographic and histological assessment. All MR imaging was performed using T1-weighted spin-echo images [echo time (TE) 15–20 ms; repetition time (TR) 150–200 ms] and T2-weighted spin-echo images (TE 80–90 ms;

TR 2000–2500 ms) obtained in the coronal, sagittal, and axial planes. MR images were interpreted by a radiologist, with equivocal cases interpreted by another radiologist specializing in bone diseases. Images were considered positive for osteonecrosis if areas of low signal intensity in the weight-bearing portion of the affected bone were noted.

For bone scans, all patients were injected intravenously with 20 to 30 mCi ^{99m}technetium methylenediphosphonate. Immediate flow images as well as static images of the skeleton were obtained. Anterior and posterior planar whole-body delayed images were obtained 3 h after injection. Findings by bone scan were considered compatible with a diagnosis of osteonecrosis when increased or decreased activity was observed in the humeral or femoral head, distal femur, proximal or distal tibia, or talus.

Histological confirmation of osteonecrosis was obtained for each suspected lesion, and this was considered the "gold standard" for comparison purposes. Histological material was obtained from bones during one of the following procedures: core decompression (n = 122), total hip arthroplasty (n = 13), limited femoral resurfacing (n = 14), various bone grafting procedures (n = 6), total knee arthroplasty (n = 3), shoulder hemiarthroplasty (n = 3), or ankle arthrodesis (n = 2). Specimens were fixed in 10% neutral buffered formalin and decalcified in a solution of formic acid and sodium citrate. Sections were cut at 4 μ m thickness and were stained with routine hematoxylin and eosin. All specimens were classified by a bone pathologist as Type III or IV osteonecrosis by the classification system of Arlet and Durroux²⁶. This represents medullary and trabecular necrosis without repair (Type III) and bone formation in apposition to dead trabeculae (Type IV).

Anteroposterior and lateral plain radiographs obtained at presentation were used to stage all symptomatic-positive joints by the Ficat and Arlet system, which was originally described for use in the hip but can be applied to any joint (Table 1)^{1,27}. All positive lesions were stratified by lesion size by the radiographic method of Kerboul and co-workers²⁸ or by MRI volumetric analysis as described²⁹. Lesions were classified as small ($< 15\%$), medium (15% to 30%), and large ($> 30\%$). Ficat and Arlet staging and size of lesion analysis were performed to evaluate if there was an association of the size of the lesion with positive bone scans.

Statistical methods. The data were compiled utilizing an Access 7.0 database (Microsoft, Redmond, WA, USA). Descriptive statistics were calculated. Histological diagnosis of osteonecrosis was used as the gold standard for assessing sensitivity of the imaging studies. Sensitivity was calculated with 95% confidence intervals and defined as the number of true-positives divided by the sum of the number of true-positives plus the number of false-negatives. Chi-square analysis with Yates' correction was performed to determine if differences between frequencies for different groups were statistically significant. All analyses were performed using Program for Epidemiologic Analysis (PEPI) software, version 2.03 (USD Inc., Stone Mountain, GA, USA).

RESULTS

We identified 48 patients who met the inclusion and exclusion criteria. Overall, 163 lesions were identified by MRI and histology, while only 91 lesions were identified by bone scan ($p = 2.56 \times 10^{-21}$). The sensitivity of the bone scan was 55.8%. None of the lesions were identified by bone scanning

Table 1. Ficat and Arlet staging system.

| Stage | Radiographic Findings |
|-------|---|
| I | None (only evident on MRI) |
| II | Diffuse sclerosis, cysts (visualized on radiographs) |
| III | Subchondral fracture (crescent sign; with or without head collapse) |
| IV | Femoral head collapse, acetabular involvement, and joint destruction (osteoarthritis) |

that were not observed on radiographs or MRI. All positive bone scans revealed increased flow and delayed activity, and no lesion was diagnosed on the basis of photopenic areas. There was complete consistency of bone scans with MR images in only 38% of patients (18/48).

Bone scanning identified a greater proportion of lesions (47/65, 72%) in oligofocal patients as compared to the multifocal patients (44/98, 45%; $p = 0.001$). In only 13 of 28 oligofocal patients and in only one of 19 multifocal patients were all of the lesions identified by bone scan. Stratification by joints involved revealed the highest yields for the knee (37/58 lesions, 64%) followed by the hip (37/61 lesions, 61%), with lower yields for the ankle (7/14 lesions), and shoulder (10/30 lesions, 33%) ($p = 0.038$; Figure 1).

A higher Ficat-Arlet stage led to a higher percentage of positive lesions. Overall, radiographic Stage I lesions had 19% positive bone scans (5/27), Stage II lesions had 56% positive scans (53/95), and Stage III lesions had 80% positive bone scans (33/41) ($p = 3 \times 10^{-6}$; Figure 2). Combining radiographically evident lesions (Stage II and III), of 136 cases, only 86 (63%) were positive on bone scans. Larger lesions also led to a higher percentage (41/58 positive scans, 71%) of positive bone scans versus medium-size (42/79 positive scans, 53%) and small (8/26 positive scans, 31%) lesions ($p = 0.002$; Figure 3).

There was no statistical difference in percentage of positive bone scans when lesions were subcategorized by patient sex, use of corticosteroids, diagnosis of SLE, or alcohol and tobacco use.

DISCUSSION

The use of radionuclides in diagnosing femoral head osteonecrosis was introduced in the 1950s by Tucker and Boyd and associates using phosphorus-32^{30,31}. Later investigation used strontium-85 in the late 1960s and early

1970s¹⁰. ^{99m}Tc was introduced in 1971 and continues to be used for standard 3-phase bone scans. These can show both deficient uptake in the femoral head (usually in posttraumatic cases of osteonecrosis) and increased uptake in the later stages of osteonecrosis¹. Before the advent of MRI, studies compared bone scans to plain radiographs and found that bone scanning was more sensitive for early lesions. Conklin and co-workers compared the sensitivities of the 2 modalities in diagnosing osteonecrosis in patients with SLE. They found sensitivities of 89% (24/27) for bone scanning and 41% (11/27) for standard radiographs (the gold standard)⁹. A reason for the higher sensitivity in their study compared to ours is that their patients had SLE, which presents at later stages of disease and differed from the many earlier-stage (I and II) lesions as in our study.

Recently, studies have shifted to comparison of the diagnostic capabilities of MRI with those of bone scanning and other techniques. MRI was found to be more sensitive than bone scans and CT in scanning for early lesions by Mitchell, *et al* in a controlled statistical study¹⁸. Similarly, in a comparison by Markisz, *et al*, MRI had an overall sensitivity of 100% compared with 81% for bone scanning (37/37 and 30/37 hips, respectively)¹⁷. MRI was found to be better than bone scanning for the diagnosis of early osteonecrosis in 25 patients with suspected lesions by Bassett, *et al*, who reported that no false-negative MR images were found, but diphosphonate scans were negative in 9 hips with normal radiographic and abnormal MR images¹⁵. Hauzeur, *et al*² studied MRI, radiographs, and bone scans in 25 patients with suspicion for osteonecrosis in 49 hips. Thirty-three hips were confirmed positive by histological examination. Among these, every MRI test was always positive, while only 77% (24/31) of available bone scans showed any signs of osteonecrosis. Additionally, 22 (67%) radiographs and 18/29 (62%) available CT scans were positive². Thus, these




Figure 1. Bone scan findings classified by joints. Percentage of positive bone scans was lower for shoulders and ankles than for hips and knees. Nevertheless, each group had low sensitivity compared to MRI scanning (100%), ranging from 33% to 64%.




Figure 2. Positive bone scan results classified by lesion size. Larger lesions in general led to higher percentages of positive bone scans compared to medium-size or small lesions.




Figure 3. Positive bone scan findings stratified by the Ficat and Arlet stage. Higher stages of disease led to higher percentages of positive bone scans.

2 studies were similar to ours, in which MRI scans were 100% sensitive for osteonecrosis, with lower sensitivities for bone scanning. Other studies have found similar low-sensitivity bone scintigraphy results for the hip^{16,32} or knee²⁰.

Some studies have reported bone scanning results more comparable to those of MRI. In a study by Steinberg of biopsy-proven hip osteonecrosis, the sensitivity, specificity, and accuracy of MRI were calculated to be 96%, 71%, and 91%, respectively, which were still superior to technetium scans (86%, 79%, and 85%), although neither test was completely accurate⁴. These results were not found in our study, in which MRI was 100% sensitive, with bone scanning detecting only 56% (91/163) of confirmed lesions. This may be because many tests in the Steinberg study involved MRI

techniques and scanners from the 1980s. In a prospective study, Stulberg and co-workers⁵ compared the multiple diagnostic modalities and calculated their respective sensitivity, specificity, and predictive values (positive and negative; Table 2). They concluded that since no imaging modality evaluated was completely accurate and since bone scanning was less expensive than MRI, the most cost-efficient initial method would be bone scans. Stulberg suggested performing MRI studies on all patients with symptomatic hips who had negative bone scans. However, since our data demonstrated that bone scanning did not detect 44% of the 163 lesions identified by MRI and histology, we disagree with this algorithm as a means of reducing the cost of a diagnostic investigation.

We believe that in our study many lesions were diag-

Table 2. Diagnostic modality comparison⁵.

| Modality | Sensitivity | Specificity | Predictive Value (positive/negative) |
|-----------------------|-------------|-------------|---|
| Bone scan | 83 | 83 | 96/48 |
| MRI | 87 | 83 | 96/55 |
| SPECT | 87 | 83 | 96/55 |
| Interosseous pressure | 80 | 60 | 95/25 |
| Biopsy | 88 | 100 | 100/25 |

MRI: magnetic resonance imaging; SPECT: single photon emission computed tomography.

nosed in early stages of the disease (especially for joints other than the hip), which may be why the sensitivity is much lower than in previous reports where lesions were diagnosed at later stages. This is important because early diagnosis is imperative, as multiple studies have shown that femoral head preservation treatment in the early stages has the most efficacy.

Some limitations of our study are that it is a retrospective review of a small population of patients. Additionally, although all imaging tests were completed within a 3-month period for each patient, durations of time between the bone scans and MRI studies varied. These limitations can be corrected in the future with a prospective study on a larger patient cohort that minimizes variability in the diagnostic investigations. Despite these limitations, we believe our conclusions concerning bone scans are accurate.

Bone scintigraphy might be justified in regions where MRI scans are not available. It could possibly be used as a screening tool for the entire body, with caution (many joints missed), but we prefer to obtain MR images of the multiple symptomatic joints to rule out this disease. Claustrophobic patients might consider this test if "open-air" MRI are not available. Bone scintigraphy might be useful in situations where metal hardware or fixation would make MRI interpretation difficult. We cannot comment on the role of bone scanning in the diagnosis of asymptomatic patients.

We observed a lower sensitivity of bone scintigraphy than MRI in diagnosing symptomatic osteonecrosis and do not support its use for diagnosis of this condition. It was least sensitive for early stage I lesions, where it might be most useful to diagnose the disease for early treatment. The utility was less effective for joints other than the hip and we do believe it is useful as a screening tool.

ACKNOWLEDGMENT

The authors thank Colleen Kazmarek for assistance with preparation of the manuscript.

REFERENCES

- Ficat R. Idiopathic bone necrosis of the femoral head. Early diagnosis and treatment. *J Bone Joint Surg Br* 1985;67:3-9.
- Hauzeur JP, Pasteels JL, Schoutens A, et al. The diagnostic value of magnetic resonance imaging in non-traumatic osteonecrosis of the femoral head. *J Bone Joint Surg Am* 1989;71:641-9.
- Koo KH, Kim R, Cho SH, Song HR, Lee G, Ko GH. Angiography, scintigraphy, intraosseous pressure, and histologic findings in high-risk osteonecrotic femoral heads with negative magnetic resonance images. *Clin Orthop Rel Res* 1994;308:127-38.
- Steinberg ME. Early diagnosis of avascular necrosis of the femoral head. *AAOS Instructional Course Lectures* 1988;37:51-7.
- Stulberg BN, Bauer TW, Belhobek GH, Levine M, Davis A. A diagnostic algorithm for osteonecrosis of the femoral head. *Clin Orthop Rel Res* 1989;249:176-82.
- Alavi A, McCloskey JR, Steinberg ME. Early detection of avascular necrosis of the femoral head by 99m technetium diphosphonate bone scan: a preliminary report. *Clin Orthop Rel Res* 1977;127:137-41.
- Arden GP, Veall N. The use of radioactive phosphorus in early detection of avascular necrosis in the femoral head in fractured neck of femur. *Proc R Soc Med* 1953;46:344-6.
- Cameron RB. Strontium-85 scintimetry in nontraumatic necrosis of the femoral head. *Clin Orthop Rel Res* 1969;65:243-61.
- Conklin JJ, Alderson PO, Zizic TM, et al. Comparison of bone scan and radiograph sensitivity in the detection of steroid-induced ischemic necrosis of bone. *Radiology* 1983;147:221-6.
- Crutchlow WP. Sr85 scintimetry of the hip in osteoarthritis and osteonecrosis. *Am J Roentgenol Radium Ther Nucl Med* 1970;109:803-12.
- D'Ambrosia RD, Shoji H, Riggins RS, Stadalnik RC, DeNardo GL. Scintigraphy in the diagnosis of osteonecrosis. *Clin Orthop Rel Res* 1978;130:139-43.
- Greiff J, Lannig S, Hoilund-Carlsen PF, Karle AK, Uhrenholdt A. Early detection by 99mTc-Sn-pyrophosphate scintigraphy of femoral head necrosis following medial femoral neck fractures. *Acta Orthop Scand* 1980;51:119-25.
- Meyers MH, Telfer N, Moore TM. Determination of the vascularity of the femoral head with technetium 99m-sulphur-colloid. *J Bone Joint Surg Am* 1977;59:658-64.
- Turner JH. Post-traumatic avascular necrosis of the femoral head predicted by preoperative technetium-99m antimony-colloid scan. An experimental and clinical study. *J Bone Joint Surg Am* 1983;65:786-96.
- Bassett LW, Gold RH, Reicher M, Bennett LR, Tooke SM. Magnetic resonance imaging in the early diagnosis of ischemic necrosis of the femoral head. Preliminary results. *Clin Orthop Rel Res* 1987;214:237-48.
- Camp JF, Colwell CW Jr. Core decompression of the femoral head for osteonecrosis. *J Bone Joint Surg Am* 1986;68:1313-9.
- Markisz JA, Knowles RJ, Altchek DW, Schneider R, Whalen JP, Cahill PT. Segmental patterns of avascular necrosis of the femoral heads: early detection with MR imaging. *Radiology* 1987;162:717-20.
- Mitchell MD, Kundel HL, Steinberg ME, Kressel HY, Alavi A, Axel L. Avascular necrosis of the hip: comparison of MR, CT, and scintigraphy. *AJR Am J Roentgenol* 1986;147:67-71.
- Gregg PJ, Walder DN. Scintigraphy versus radiography in the early diagnosis of experimental bone necrosis, with special reference to caisson disease of bone. *J Bone Joint Surg Br* 1980;62B:214-21.
- Sakai T, Sugano N, Nishii T, Haraguchi K, Yoshikawa H, Ohzono K. Bone scintigraphy for osteonecrosis of the knee in patients with non-traumatic osteonecrosis of the femoral head: comparison with magnetic resonance imaging. *Ann Rheum Dis* 2001;60:14-20.
- Hirota Y, Hirohata T, Fukuda K, et al. Association of alcohol intake, cigarette smoking, and occupational status with the risk of idiopathic osteonecrosis of the femoral head. *Am J Epidemiol* 1993;137:530-8.
- Mok CC, Lau CS, Wong RW. Risk factors for avascular bone necrosis in systemic lupus erythematosus. *Br J Rheumatol* 1998;37:895-900.

23. Hong N, Du XK. Avascular necrosis of bone in severe acute respiratory syndrome. *Clin Radiol* 2004;59:602-8.
24. Koo KH, Kim R, Kim YS, et al. Risk period for developing osteonecrosis of the femoral head in patients on steroid treatment. *Clin Rheumatol* 2002;21:299-303.
25. Matsuo K, Hirohata T, Sugioka Y, Ikeda M, Fukuda A. Influence of alcohol intake, cigarette smoking, and occupational status on idiopathic osteonecrosis of the femoral head. *Clin Orthop Rel Res* 1988;234:115-23.
26. Arlet J, Durroux R, Fauchier C, Thiechart M. Histopathology of the nontraumatic necrosis of the femoral head: Topographic and evolutive aspect. In: Arlet J, Ficat RP, Hungerford DS, editors. *Bone circulation*. Baltimore: Williams and Wilkins; 1984.
27. Ficat RP, Arlet J. Ischemia and necrosis of bone. In: Arlet J, Ficat RP, Hungerford DS, editors. *Bone circulation*. Baltimore: Williams and Wilkins; 1980.
28. Kerboul M, Thomine J, Postel M, Merle d'Aubigne R. The conservative surgical treatment of idiopathic aseptic necrosis of the femoral head. *J Bone Joint Surg Br* 1974;56:291-6.
29. Steinberg ME, Hayken GD, Steinberg DR. A quantitative system for staging avascular necrosis. *J Bone Joint Surg Br* 1995;77:34-41.
30. Boyd HB, Zilversmit DB, Calandruccio RA. The use of radioactive phosphorus (P32) to determine the viability of the head of the femur. *J Bone Joint Surg Am* 1955;37:260-9.
31. Tucker FR. The use of radioactive phosphorus in the diagnosis of avascular necrosis of the femoral head. *J Bone Joint Surg* 1950;32:100-7.
32. Scheiber C, Meyer ME, Dumitresco B, et al. The pitfalls of planar three-phase bone scintigraphy in nontraumatic hip avascular osteonecrosis. *Clin Nucl Med* 1999;24:488-94.