Let's Talk About Sex!

To the Editor:

The report by Silva and colleagues¹ and the accompanying editorial² have not only informed us about pregnancy outcome in juvenile systemic lupus erythematosus (SLE) but also reminded us of the importance of addressing sexual health within pediatric rheumatology. Schwarz and Manzi similarly advocated increasing attention to sexual health in adult rheumatology clinics³. However, addressing such issues within the context of adolescence demands different knowledge and expertise than that required in adult practice.

In a recent audit of 19 adolescents with SLE seen in a pediatric rheumatology clinic at our institution (median age 15 yrs, range 10–18), documentation of sexual health being addressed was found in only 4 patients⁴. In a similar audit of adolescents with juvenile idiopathic arthritis (mean age 19.2 yrs), documentation of sexual health was found in only 11.4% of case notes⁵, despite young people calling for sexual health information and advice within the context of their rheumatology care⁶. This is particularly pertinent in view of the use of methotrexate and biologics in pediatric rheumatology and the fact that at least a third of young people are likely to be sexually active prior to transfer to adult care⁷. The benefit of transitional care for adolescents that addresses such issues is highlighted by significant improvement in documentation of sexual health (albeit still suboptimal) following implementation of a transitional care program⁵.

In order to address sexual health within pediatric settings, developmentally appropriate methods of assessment and health promotion are required. Few adolescents will initiate discussion of such issues, so routine use of screening tools such as HEADSS (Home environment, Education/employment, peer-related Activities, Drugs, Sexuality, Suicide/depression)⁸ is useful within pediatric rheumatology clinics. Sexual health is also not all about preventing pregnancy, despite this often being the only aspect of sexual health discussed — 100% of health professionals reported discussion about infertility with male adolescents with cystic fibrosis, but only 38% discussed importance of condoms, thus ignoring the risk of sexually transmitted infections and unsafe sex⁹. A potential barrier to addressing sexual health issues in juvenile SLE is the lack of opportunity for young people to be seen independently of their parents and the triage of other competing medical problems. Service development for juvenile SLE must take this into consideration and advocate for confidential services for young people, with appropriate appointment duration and continuity of health professionals. Such rheumatology services must also ensure effective signposting to young-person-friendly sexual health clinics to ensure access to contraception, not only knowledge, as highlighted by Dr. Kaufman². Finally, professionals with adolescent health expertise are integral to such services, although not universally available in pediatric and

adult rheumatology¹⁰. The development of recent e-learning training resources in the UK (e.g., the Royal College of Paediatrics and Child Health; www.rcpch.ac.uk/ahp) is a welcome and exciting initiative from which rheumatology can only gain.

CLARE BIDWELL, MBChB; ISABEL BOLT, MD, Paediatric and Adolescent Rheumatology, Birmingham Children's Hospital; JANET E. McDONAGH, MBBS, MD, FRCP, Institute of Child Health, University of Birmingham, Birmingham, United Kingdom.

REFERENCES

1. Silva CAA, Hilario MO, Febronio MV, et al. Pregnancy outcome in juvenile systemic lupus erythematosus: A Brazilian multicenter cohort study. *J Rheumatol* 2008;35:1414–8.
2. Kaufman M. Pregnant adolescents and youth with systemic lupus erythematosus: can new data inform our approach to young women with SLE? [editorial]. *J Rheumatol* 2008;35:1240–1.
3. Schwarz EB, Manzi S. Risk of unintended pregnancy among women with systemic lupus erythematosus. *Arthritis Care Res* 2008;59:863–6.
4. Bidwell C, McDonagh JE. Cardiovascular risk factors in juvenile SLE: A missed adolescent health opportunity in rheumatology clinics? *Arch Dis Child* 2008;93 Suppl: A67.
5. Robertson LP, McDonagh JE, Southwood TR, Shaw KL. Growing up and moving on. A multicentre UK audit of the transfer of adolescents with juvenile idiopathic arthritis from paediatric to adult centred care. *Ann Rheum Dis* 2006;65:74–80.
6. Shaw KL, Southwood TR, McDonagh JE, on behalf of the British Paediatric Rheumatology Group. User perspectives of transitional care for adolescents with juvenile idiopathic arthritis. *Rheumatology Oxford* 2004;43:770–8.
7. Packham JC, Hall MA. Long-term follow-up of 246 adults with juvenile idiopathic arthritis: social function, relationships and sexual activity. *Rheumatology Oxford* 2002;41:1440–3.
8. Goldenring JM, Rosen DS. Getting into adolescent heads: an essential update. *Contemporary Pediatrics* 2004;21:64–90.
9. Sawyer SM, Tully M-AM, Colin AA. Reproductive and sexual health in males with cystic fibrosis: a case for health professional education and training. *J Adolescent Health* 2001;28:36–40.
10. McDonagh JE, Southwood TR, Shaw KL. Unmet education and training needs of rheumatology health professionals in adolescent health and transitional care. *Rheumatology Oxford* 2004;43:737–43.

J Rheumatol 2009;36:6; doi:10.3899/jrheum080956