

Visual Manifestations in Giant Cell Arteritis: Trend over 5 Decades in a Population-based Cohort

Abha G. Singh, Tanaz A. Kermani, Cynthia S. Crowson, Cornelia M. Weyand, Eric L. Matteson, and Kenneth J. Warrington

ABSTRACT. Objective. To evaluate clinical characteristics, treatment, and outcomes of patients with visual changes from giant cell arteritis (GCA) and to examine trends over the last 5 decades.

Methods. We reviewed the medical records of a population-based cohort of patients with GCA diagnosed between 1950 and 2004. The clinical, ophthalmological, and laboratory features of patients with visual manifestations attributable to GCA were compared to patients without visual complications. Trends over time were examined using logistic regression modeling adjusted for age and sex.

Results. In a cohort of 204 cases of GCA (mean age 76.0 ± 8.2 yrs, 80% female), visual changes from GCA were observed in 47 patients (23%), and 4.4% suffered complete vision loss. A higher proportion of patients with visual manifestations reported jaw claudication than did patients without visual changes (55% vs 38%, $p = 0.04$). Over a period of 55 years, we observed a significant decline in the incidence of visual symptoms due to GCA. There was a lower incidence of ischemic optic neuropathy in the 1980–2004 cohort vs 1950–1979 (6% vs 15%, $p = 0.03$). Patients diagnosed in later decades were more likely to recover from visual symptoms (HR 1.34, 95% CI 1.06–1.71). Chances of recovery were poor in patients with anterior ischemic optic neuropathy or complete vision loss.

Conclusion. Incidence of visual symptoms has declined over the past 5 decades, and chances of recovery from visual symptoms have improved. However, complete loss of vision is essentially irreversible. Jaw claudication is associated with higher likelihood of development of visual symptoms. (First Release Dec 15 2014; J Rheumatol 2015;42:309–15; doi:10.3899/jrheum.140188)

Key Indexing Terms:

GIANT CELL ARTERITIS
VASCULITIS

VISION LOSS
STEROIDS

VISUAL MANIFESTATIONS
BLINDNESS

Giant cell arteritis (GCA) is a granulomatous large vessel vasculitis and is the most common vasculitis in Western countries in the elderly age group¹. Since Horton, *et al*'s first description² of temporal arteritis in the United States, this form of systemic vasculitis has become more widely recognized, as has its potential for serious sequelae such as blindness and death³.

The annual incidence of GCA in a population-based

From the Division of Rheumatology, Mayo Clinic, Rochester, Minnesota; Division of Rheumatology, University of California, Los Angeles; Division of Immunology and Rheumatology, Stanford University School of Medicine, Stanford, California, USA.

Supported by the resources of the Rochester Epidemiology Project, which is supported by the US National Institute on Aging of the National Institutes of Health (NIH) under Award Number R01AG034676. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

A.G. Singh, MBBS, Division of Rheumatology, Mayo Clinic; T.A. Kermani, MD, MS, Division of Rheumatology, University of California, Los Angeles; C.S. Crowson, MS, Division of Rheumatology, Mayo Clinic; C.M. Weyand, MD, PhD, Division of Immunology and Rheumatology, Stanford University School of Medicine; E.L. Matteson, MD, MPH; K.J. Warrington, MD, Division of Rheumatology, Mayo Clinic.

Address correspondence to Dr. K.J. Warrington, MD, Mayo Clinic, 200 1st St. SW, Rochester, Minnesota, USA.

E-mail: warrington.kenneth@mayo.edu

Accepted for publication October 15, 2014.

cohort from Olmsted County, Minnesota, USA, has been estimated around 18.9 per 100,000 people over the age of 50 years^{4,5}. The highest incidence of GCA has been observed in whites of Northern European descent (up to 30/100,000)⁶. Women are affected 2 to 3 times more often than men^{1,7}.

GCA affects medium and large-sized vessels, with predisposition for cranial arteries¹. Cranial ischemic complications (particularly permanent visual loss due to vasculitic involvement of the posterior ciliary arteries) are the most feared aspects of GCA⁸. Incidence of visual complications was particularly high (35%–60%) before the introduction of corticosteroid use for GCA^{9,10,11}.

Early initiation of empiric corticosteroids at onset of symptoms in patients with suspected GCA may decrease the risk of permanent vision loss¹²; however, it is unclear whether this strategy has modified the course of visual complications over time. We performed this population-based, historical cohort analysis evaluating visual complications in patients with GCA and examined trends over the last 5 decades.

MATERIALS AND METHODS

Study design. This is a retrospective population-based cohort study performed using the resources of the Rochester Epidemiology Project (REP) medical records linkage system in Olmsted County¹³. Popula-

tion-based epidemiologic research can be conducted in Olmsted County because medical care is virtually self-contained within the community, and complete (inpatient and outpatient) medical records for county residents are available for review. After approval by the Institutional Review Board, we used the resources of the REP to identify all Rochester, Minnesota, residents > 18 years of age who fulfilled the American College of Rheumatology (ACR) 1990 criteria for GCA between January 1, 1950, and December 31, 2004.

Data collection. Patient records were reviewed and details were abstracted regarding demographics, clinical features at the time of GCA diagnosis, results of ophthalmologic evaluation, and laboratory variables at the time of diagnosis and on subsequent visits. Two groups of patients with GCA were identified from the cohort: patients with visual manifestations attributed to GCA, and patients without visual manifestations or with vision changes deemed to be unrelated to GCA (cataract, age-related macular degeneration, etc.). Ophthalmologic examination records, when available, were reviewed to ascertain that the vision manifestations were related to GCA. When ophthalmologic examination findings were not available, data were obtained from review of the treating physician records. Data on the entire treatment course and response to therapy were also collected. Patients were followed until migration, death, or the end of the study (December 31, 2009).

We also analyzed the association between vision changes in GCA and risk of cardiovascular events and all-cause mortality.

Statistical methods. Descriptive statistics (mean and SD, or median and interquartile range, or percentages, as appropriate) were used to describe the cohort. Baseline variables were compared in patients with GCA who developed visual manifestations to the remainder of the cohort, using chi-squared tests for categorical variables and Wilcoxon rank-sum tests for continuous variables. The cohort was divided into 2 time periods (those diagnosed with GCA in 1950–1979, and those diagnosed in 1980–2004), and comparison between time periods was performed using chi-squared and Wilcoxon rank-sum tests.

Trends over time were analyzed using smoothing splines in logistic regression models adjusted for age and sex. Factors associated with time to resolution of visual symptoms were examined using Cox proportional hazard models. P values < 0.05 were considered statistically significant. All analyses were performed using SAS statistical software, version 9.3.

RESULTS

Study cohort. This study cohort included 204 patients with GCA (mean age 76.0 ± 8.2 yrs, 80% female), all of whom met the ACR 1990 criteria for classification of GCA. Temporal artery biopsy was positive in 92% of the 192 patients in whom it was performed. At diagnosis, 47 patients (23%) had visual manifestations attributable to GCA. Demographic characteristics and clinical features in those with visual manifestations are compared to those without visual manifestations in Table 1.

Visual manifestations. Among the 47 patients with visual manifestations, the most common visual symptoms were blurred vision (31 patients, 66%) and diplopia (11 patients, 23%; Table 2). Other visual manifestations included amaurosis fugax (7 patients, 15%) and partial visual field loss (9 patients, 19%). Nine patients (19%) had complete loss of vision, which was unilateral in 7 patients and bilateral in 2 patients. Ischemic optic neuropathy (ION) was the predominant ophthalmologic diagnosis (17 patients, 36%). Other diagnoses included central retinal artery occlusion (2 patients, 4%) and nonspecific ophthalmologic

findings such as venous congestion and retinal hemorrhages (Table 3).

Treatment. The cumulative dose of corticosteroids administered on the first day of suspected GCA was significantly higher in patients presenting with vision changes, as compared to those with no vision changes (97.6 mg vs 58.3 mg, respectively; $p < 0.001$). Steroids were also initiated earlier in this group when compared to those with no vision-related symptoms. Indeed, corticosteroids were started an average of 2 days prior to confirmed diagnosis of GCA in patients presenting with vision changes. However, in assessing time from first symptom suspicious for GCA to initiation of corticosteroids between the 2 groups, no significant difference was noted (Table 1). There was no difference in the cumulative dose of steroids at 1, 2, or 5 years after GCA diagnosis between those with and without vision loss (Table 4). We also did not observe an increased risk of steroid complications among those with and without vision loss (including diabetes mellitus, cataracts, hip fractures, symptomatic vertebral fractures, Colles fracture of the wrist, avascular necrosis, bacteremia, pneumonitis, other infections, gastrointestinal bleeding, myopathy, hypertension, and hyperlipidemia; $p > 0.45$ for each steroid-related complication comparing patients with vision loss to those without vision loss, adjusting for age, sex, and calendar year of GCA diagnosis).

Ten patients in the GCA cohort had been treated with other immunosuppressive agents: methotrexate (5 patients), cyclophosphamide (2 patients), and azathioprine (5 patients); none of them had visual manifestations attributable to GCA.

Prognosis. Among patients without complete loss of vision, 75% had full resolution of their symptoms within 3 months. Recovery from visual symptoms, however, was less likely in patients with complete vision loss (HR 0.20, 95% CI 0.06–0.63, $p < 0.01$) when compared with blurred vision alone. Four of the 9 patients with complete loss of vision reported a slight improvement in vision after initiation of treatment, but visual acuity testing failed to confirm an improvement except in 1 patient whose vision in the affected eye improved from absence of light perception to appreciation of hand movement after treatment. One patient, diagnosed in 1954, developed vision loss after the start of corticosteroids. In this patient, corticosteroids were started 7 days prior to the vision loss, at a dose of 60 mg/day. Patients with ophthalmologic diagnosis of anterior ION were less likely to have recovery of symptoms compared to other ophthalmologic diagnoses (HR 0.27, 95% CI 0.12–0.61, $p = 0.002$).

In our cohort, during followup, 154 patients with GCA died and 26 developed acute coronary syndrome. There was no evidence of an association between vision loss and all-cause mortality (HR 1.14, 95% CI 0.77–1.68, $p = 0.53$) or acute coronary syndrome (HR 0.56, 95% CI 0.16–1.87,

Table 1. Clinical and laboratory features of patients with and without visual manifestations in a cohort of 204 patients with giant cell arteritis (GCA). Values are n (%) or mean \pm SD, except where indicated.

Characteristics	Patients With Visual Manifestations, n = 47	Patients Without Visual Manifestations, n = 157	p
Demographics			
Age, years	77.4 \pm 9.2	75.6 \pm 7.8	0.16
Sex, females	36 (77)	127 (81)	0.52
Ever smoker	17 (42)	67 (46)	0.62
Clinical features			
Fever > 100°F	6 (13)	33 (22)	0.20
Duration of symptoms, days, median (IQR)	26 (7.8–44)	24 (10–53)	0.40
Significant weight loss*	6 (13)	38 (24)	0.09
Fatigue	9 (19)	51 (33)	0.07
Headache	29 (62)	115 (74)	0.10
Jaw claudication	26 (55)	59 (38)	0.036
Scalp tenderness	15 (33)	62 (42)	0.29
Associated polymyalgia rheumatica	10 (21)	40 (26)	0.53
Neurologic symptoms	3 (6)	2 (1)	0.050
Temporal tenderness on examination	10 (28)	42 (31)	0.68
Carotid/subclavian bruit on examination	1 (2)	6 (4)	0.54
CAD before GCA diagnosis	13 (28)	28 (18)	0.14
Aspirin use before GCA diagnosis	10 (24)	5 (16)	0.22
Corticosteroid use			
Start dose CS, initial, mg	59.6 \pm 20	52 \pm 14	0.030
Cumulative CS dose on day 1, mg	97.7 \pm 275	58.3 \pm 145	0.002
Cumulative CS within 1st week, mg	466.3 \pm 534	398 \pm 437	0.010
Time to initiation of steroids (relative to GCA diagnosis), days	–2.0 \pm 2.7	2.6 \pm 44	0.010
Laboratory features			
Hemoglobin, g/dl	11.9 \pm 1.3	11.7 \pm 1.4	0.16
ESR, mm/h	77 \pm 28	80 \pm 31	0.45
White blood cell count, $\times 10^3/\mu\text{l}$	5.4 \pm 3.6	5.9 \pm 4.4	0.63
Hyperlipidemia before GCA diagnosis	18 (38)	91 (58)	0.018

P values < 0.05 shown in bold. * Defined as > 10% body weight in 6 months. IQR: interquartile range; CAD: coronary artery disease; CS: corticosteroids; ESR: erythrocyte sedimentation rate.

Table 2. Visual manifestations among 204 patients with giant cell arteritis in Olmsted County, Minnesota, USA, diagnosed between 1950 and 2004 (n = 47). Some patients had more than 1 symptom.

Visual Symptom	No. Patients (%)
Amaurosis fugax	7 (15)
Diplopia	11 (23)
Blurred vision	31 (66)
Field loss	9 (19)
Complete loss	
Unilateral	7 (15)
Bilateral	2 (4)

p = 0.34), after adjustment for age, sex, and calendar year of GCA diagnosis.

Time trends. Over the 55-year study period, a significant decline in the incidence of visual manifestations from GCA was observed (Figure 1). Incidence of blurred vision decreased from 25% in patients diagnosed in 1950–1979 to

Table 3. Ophthalmologic examination findings in 47 patients with visual symptoms from giant cell arteritis in Olmsted County, Minnesota, USA, between 1950 and 2004.

Diagnosis	No. Patients (%)
Anterior ischemic optic neuropathy	
Unilateral	12 (25)
Bilateral	5 (11)
Central retinal artery occlusion	2 (4)
Other unrelated findings*	5 (11)
No ophthalmologic findings	16 (34)
Eye examination not performed	7 (15)

*Included venous congestion, retinal hemorrhages, choroidal sclerosis, arteriolar narrowing, external rectus weakness, and oculomotor nerve palsy.

11% in 1980–2004 (p = 0.015). Fewer patients diagnosed with GCA in 1980–2004 had complete vision loss (2%) compared to those diagnosed from 1950–1979 (9.8%, p = 0.014). Incidence of ION was also less in patients diagnosed

Table 4. Cumulative corticosteroid exposure in patients with GCA, with and without visual manifestations.

	Patients With Visual Manifestations, n = 47	Patients Without Visual Manifestations, n = 157	p
One-yr cumulative corticosteroid dose			
Mean (SD; g)	5.4 (2.7)	5.5 (2.5)	0.93
Median (IQR)	5.4 (3.4–6.9)	5.4 (3.7–7.1)	
2-yr cumulative corticosteroid dose			
Mean (SD; g)	7.0 (3.7)	7.3 (3.8)	0.69
Median (IQR)	7.1 (4.0–9.3)	7.2 (4.2–10.0)	
5-yr cumulative corticosteroid dose			
Mean (SD; g)	9.9 (6.3)	9.3 (6.3)	0.63
Median (IQR)	8.3 (4.8–14.3)	8.1 (4.3–12.7)	

GCA: giant cell arteritis; IQR: interquartile range.




Figure 1. The incidence of visual manifestations declined over time ($p = 0.02$; adjusted for age and sex) among 204 patients with giant cell arteritis (GCA) in Olmsted County, Minnesota, USA, between 1950 and 2004. Point estimate and 95% CI for each 5-year time period are displayed.

with GCA from 1980–2004 (8/143, 6%) compared to those between 1950 and 1979 (9/61, 15%, $p = 0.03$).

Patients diagnosed with GCA in later decades were more likely to have recovery from visual symptoms [(HR 1.34, 95% CI 1.06–1.71) per 10-yr increase in calendar year, $p = 0.016$]. In the cohort diagnosed with GCA in 1980–2004, 80% of patients with visual symptoms had resolution within 3 months of diagnosis, compared to 68% in patients diagnosed in earlier years (1950–1979).

DISCUSSION

In 1950, Shick, *et al* from the Mayo Clinic reported the first

use of corticosteroids in GCA¹⁴. Much experience has been gained since then in the treatment of GCA with corticosteroids, but the potential for vision loss, the most feared complication of this form of vasculitis, is not well explored.

This is the first report, to our knowledge, describing incidence and trends in visual manifestations in patients with GCA over a long period (spanning 55 years) beginning shortly after the introduction of corticosteroids for treatment of GCA¹⁴ and extending to the current era. We found a decline in visual complications from GCA over the decades. The overall incidence of blurred vision decreased from 25% (1950–1979) to 11% (1980–2004), and only 2% of patients

diagnosed with GCA in recent decades had complete vision loss, as compared to 9.8% of patients diagnosed from 1950–1979. Moreover, patients diagnosed with GCA in later decades were more likely to recover from visual symptoms; in the cohort diagnosed with GCA in 1980–2004, 80% of patients with visual symptoms had resolution within 3 months of diagnosis, compared to 68% in patients diagnosed in earlier years (1950–1979). This may be due to increased awareness of the disease and its potential for causing permanent blindness, earlier diagnosis, and timely initiation of treatment, reflected in the early and higher-dose corticosteroid use in patients with visual symptoms in this cohort. Recovery from some visual complications has become more likely in recent decades.

patient develops visual loss. It has also been suggested that acute-phase reactants such as haptoglobin may promote neovascularization in patients with higher inflammatory response, hence the lower incidence of ischemic complications in this group¹⁹. Hyperlipidemia was noted to be more frequent in patients without vision changes in our cohort (Table 1); this finding is of uncertain clinical significance. While hyperlipidemia can be associated with aortic aneurysm and/or dissection in patients with large vessel involvement²³, prior studies have shown no association between use of lipid-lowering agents and cranial complications from GCA^{24,25}. We did not find any association between patients with visual changes and risk of acute coronary syndrome or all-cause mortality.

Table 5. Summary of previous reports showing a declining trend in visual manifestations in patients with GCA over the years.

*No. patients with vision changes due to GCA/total no. patients with GCA in that cohort. NR: not reported; GCA: giant cell arteritis.

endothelin receptor blockers may have a potential role in acute management of vision changes in GCA.

Not only has the incidence of visual manifestations declined, but we also noted a higher rate of recovery from visual symptoms in more recent decades, an interesting and novel finding. In the period 1980–2004, 80% of visual symptoms resolved within 3 months compared to 68% in patients diagnosed in earlier years (1950–1979). Chances of recovery from complete loss of vision, however, remain grim. Of the 9 patients with complete loss of vision in at least 1 eye, all were treated with systemic glucocorticoids. Four reported a slight improvement, but this could not be confirmed on objective ophthalmologic testing except for marginal improvement in vision in 1 patient. Two patients diagnosed with GCA, in 1953 and 1974, had complete vision loss in both eyes. Notably, patients with a diagnosis of anterior ION were significantly less likely to recover compared to those with any other ophthalmologic diagnoses. Pathogenically, vision loss is a result of tissue ischemia due to luminal occlusion of inflamed blood vessel. The underlying pathomechanism is the formation of lumen-occlusive intimal hyperplasia, a neotissue that remains in place even in treated vasculitis. Data presented here confirm that vision loss is permanent once tissue ischemia has occurred. This finding may be of prognostic significance in counseling patients who present with changes in vision secondary to GCA and highlights the importance of an urgent ophthalmologic examination in suspected GCA.

The duration of our study and important new findings regarding factors associated with recovery in patients with GCA add significantly to our knowledge about this disease. According to the 2000 US census data, 90.3% of the Olmsted County population is white. GCA predominantly affects individuals of Northern European descent, but the results of our study would be generalizable to other white patients with GCA in the United States. Population-based design and use of the REP, which allowed review of the entire individual medical record covering all inpatient and outpatient care from all of the local healthcare providers, add significantly to the comprehensiveness and reliability of our findings.

The primary limitation of our study is its retrospective design. In spite of comprehensive medical records after presentation, baseline visual acuity prior to onset of visual symptoms was not available in all cases and hence, change in vision could not be objectively studied for many patients. The majority of patients with vision changes had a complete evaluation by an ophthalmologist (41/47). However, this evaluation varied over the course of this 5-decade longitudinal study. The retrospective nature of our study also limited us in using strict objective criteria for improvement in vision after treatment initiation. We defined vision improvement based on a combination of patient self-report

and physician assessment, as documented in medical records. We acknowledge that such subjective evaluation is not ideal, and propose that future prospective cohort studies on this topic use strictly defined objective criteria for vision changes. Additionally, while every attempt was made to confirm that vision changes were acute and clinically related to GCA, the possibility of misclassification bias cannot be completely excluded.

It is reassuring to find that the incidence of visual manifestations in GCA has significantly declined over the decades, but there remains a 4% incidence of complete sight loss and 8% incidence of anterior ION in GCA, both of which signify permanence of vision loss with essentially no chance of recovery. Patients with vision changes are likely to receive earlier and higher doses of corticosteroids, which remains the most important treatment strategy. Although there has been progress over the years, clinicians must work toward avoiding loss of vision by patient education, prompt recognition of this disease, and early institution of treatment. The potential psychosocioeconomic effect of vision loss in patients with GCA merits further review.

REFERENCES

1. Salvarani C, Cantini F, Hunder GG. Polymyalgia rheumatica and giant-cell arteritis. *Lancet* 2008;372:234–45.
2. Horton BT, Magath TB, Brown GE. Undescribed form of arteritis of temporal vessels. *Proc Staff Meet Mayo Clin* 1932;7:700–1.
3. Paulley JW, Hughes JP. Giant-cell arteritis, or arteritis of the aged. *Br Med J* 1960;2:1562–7.
4. Kermani TA, Schafer VS, Crowson CS, Hunder GG, Gabriel SE, Matteson EL, et al. Increase in age at onset of giant cell arteritis: a population-based study. *Ann Rheum Dis* 2010;69:780–1.
5. Salvarani C, Crowson CS, O’Fallon WM, Hunder GG, Gabriel SE. Reappraisal of the epidemiology of giant cell arteritis in Olmsted County, Minnesota, over a fifty-year period. *Arthritis Rheum* 2004;51:264–8.
6. Gonzalez-Gay MA, Garcia-Porrua C, Llorca J, Hajeer AH, Brañas F, Dababneh A, et al. Visual manifestations of giant cell arteritis. Trends and clinical spectrum in 161 patients. *Medicine* 2000;79:283–92.
7. Machado EB, Michet CJ, Ballard DJ, Hunder GG, Beard CM, Chu CP, et al. Trends in incidence and clinical presentation of temporal arteritis in Olmsted County, Minnesota, 1950–1985. *Arthritis Rheum* 1988;31:745–9.
8. Figus M, Talarico R, Posarelli C, d’Ascanio A, Elefante E, Bombardieri S. Ocular involvement in giant cell arteritis. *Clin Exp Rheumatol* 2013;31:S96.
9. Birkhead NC, Wagener HP, Shick RM. Treatment of temporal arteritis with adrenal corticosteroids; results in fifty-five cases in which lesion was proved at biopsy. *J Am Med Assoc* 1957; 163:821–7.
10. Bruce GM. Temporal arteritis as a cause of blindness; review of literature and report of a case. *Trans Am Ophthalmol Soc* 1949;47:300–16.
11. Cooke WT, Cloake PC, Govan AD, Colbeck JC. Temporal arteritis; a generalized vascular disease. *Q J Med* 1946;15:47–75.
12. Hayreh SS, Zimmerman B, Kardon RH. Visual improvement with corticosteroid therapy in giant cell arteritis. Report of a large study and review of literature. *Acta Ophthalmol Scand* 2002;80:355–67.
13. Melton LJ 3rd. History of the Rochester Epidemiology Project. *Mayo Clin Proc* 1996;71:266–74.

14. Shick RM, Baggenstoss AH, Fuller BF, Polley HF. Effects of cortisone and ACTH on periarteritis nodosa and cranial arteritis. *Proc Staff Meet Mayo Clin* 1950;25:492-4.
15. Calamia KT, Hunder GG. Giant cell arteritis (temporal arteritis) presenting as fever of undetermined origin. *Arthritis Rheum* 1981;24:1414-8.
16. Caselli RJ, Hunder GG, Whisnant JP. Neurologic disease in biopsy-proven giant cell (temporal) arteritis. *Neurology* 1988;38:352-9.
17. Gonzalez-Gay MA, Miranda-Filloy JA, Lopez-Diaz MJ, Perez-Alvarez R, Gonzalez-Juanatey C, Sanchez-Andrade A, et al. Giant cell arteritis in northwestern Spain: a 25-year epidemiologic study. *Medicine* 2007;86:61-8.
18. Nesher G, Berkun Y, Mates M, Baras M, Nesher R, Rubinow A, et al. Risk factors for cranial ischemic complications in giant cell arteritis. *Medicine* 2004;83:114-22.
19. Cid MC, Font C, Oristrell J, de la Sierra A, Coll-Vinent B, López-Soto A, et al. Association between strong inflammatory response and low risk of developing visual loss and other cranial ischemic complications in giant cell (temporal) arteritis. *Arthritis Rheum* 1998;41:26-32.
20. Liozon E, Herrmann F, Ly K, Robert PY, Loustaud V, Soria P, et al. Risk factors for visual loss in giant cell (temporal) arteritis: a prospective study of 174 patients. *Am J Med* 2001;111:211-7.
21. Salvarani C, Cimino L, Macchioni P, Consonni D, Cantini F, Bajocchi G, et al. Risk factors for visual loss in an Italian population-based cohort of patients with giant cell arteritis. *Arthritis Rheum* 2005;53:293-7.
22. Kermani TA, Schmidt J, Crowson CS, Ytterberg SR, Hunder GG, Matteson EL, et al. Utility of erythrocyte sedimentation rate and C-reactive protein for the diagnosis of giant cell arteritis. *Semin Arthritis Rheum* 2012;41:866-71.
23. Nuenninghoff DM, Hunder GG, Christianson TJ, McClelland RL, Matteson EL. Incidence and predictors of large-artery complication (aortic aneurysm, aortic dissection, and/or large-artery stenosis) in patients with giant cell arteritis: a population-based study over 50 years. *Arthritis Rheum* 2003;48:3522-31.
24. Schmidt J, Kermani TA, Muratore F, Crowson CS, Matteson EL, Warrington KJ. Statin use in giant cell arteritis: a retrospective study. *J Rheumatol* 2013;40:910-5.
25. Narvaez J, Bernad B, Nolla JM, Valverde J. Statin therapy does not seem to benefit giant cell arteritis. *Semin Arthritis Rheum* 2007;36:322-7.
26. Aiello PD, Trautmann JC, McPhee TJ, Kunselman AR, Hunder GG. Visual prognosis in giant cell arteritis. *Ophthalmology* 1993;100:550-5.
27. Bengtsson BA, Malmvall BE. The epidemiology of giant cell arteritis including temporal arteritis and polymyalgia rheumatica. Incidences of different clinical presentations and eye complications. *Arthritis Rheum* 1981;24:899-904.
28. Font C, Cid MC, Coll-Vinent B, López-Soto A, Grau JM. Clinical features in patients with permanent visual loss due to biopsy-proven giant cell arteritis. *Br J Rheumatol* 1997;36:251-4.
29. Kupersmith MJ, Langer R, Mitnick H, Spiera R, Spiera H, Richmond M, et al. Visual performance in giant cell arteritis (temporal arteritis) after 1 year of therapy. *Br J Ophthalmol* 1999;83:796-801.
30. Chan CC, Paine M, O'Day J. Steroid management in giant cell arteritis. *Br J Ophthalmol* 2001;85:1061-4.
31. Hayreh SS, Zimmerman B. Visual deterioration in giant cell arteritis patients while on high doses of corticosteroid therapy. *Ophthalmology* 2003;110:1204-15.
32. Liu GT, Glaser JS, Schatz NJ, Smith JL. Visual morbidity in giant cell arteritis. Clinical characteristics and prognosis for vision. *Ophthalmology* 1994;101:1779-85.
33. Hayreh SS, Bioussé V. Treatment of acute visual loss in giant cell arteritis: should we prescribe high-dose intravenous steroids or just oral steroids? *J Neuroophthalmol* 2012;32:278-87.
34. Lee MS, Smith SD, Galor A, Hoffman GS. Antiplatelet and anticoagulant therapy in patients with giant cell arteritis. *Arthritis Rheum* 2006;54:3306-9.
35. Lozano E, Segarra M, Corbera-Bellalta M, García-Martínez A, Espígol-Frigolé G, Plà-Campo A, et al. Increased expression of the endothelin system in arterial lesions from patients with giant-cell arteritis: association between elevated plasma endothelin levels and the development of ischaemic events. *Ann Rheum Dis* 2010;69:434-42.
36. Nesher G, Rubinow A, Sonnenblick M. Trends in the clinical presentation of temporal arteritis in Israel: reflection of increased physician awareness. *Clin Rheumatol* 1996;15:483-5.