True Vitamin D Deficiency with Secondary Hyperparathyroidism

To the Editor:

We read with interest the article by Koeckhoven, *et al*¹ suggesting that upper leg strength is associated with 25-hydroxy Vitamin D [25(OH)D] levels in patients with knee osteoarthritis (OA). The association seems to exist when adjustment is made for several confounding factors. However, adjusting for body mass index attenuated this association. One important limitation of their study is the inadequate assessment of true Vitamin D deficiency. Although < 20 ng/ml is a widely accepted threshold for Vitamin D deficiency, many subjects with low 25(OH)D levels do not have clinical symptoms of Vitamin D deficiency, such as muscle weakness or widespread pain. High parathyroid hormone (PTH) level is an important indicator of true Vitamin D deficiency, which occurs in only 10% to 33% of people with Vitamin D insufficiency². Further, as we can see clearly in the study by Visser, *et al*³, a 25(OH)D < 10 ng/ml is associated with a significant loss of grip strength during the followup period, whereas this was not the case for subjects with 25(OH)D levels between 10 ng/ml and 20 ng/ml. The study by Visser, *et al*³ also indicated that higher PTH level was associated with loss of muscle strength. Another recent study by Jin, *et al*⁴ in patients with knee OA and 25(OH)D levels < 24 ng/ml suggested that monthly 50,000 IU of Vitamin D3 treatment did not improve knee pain or preserve tibial cartilage volume over 2 years. Of note, PTH levels were not reported in the study of Jin, *et al*⁴ as well. Thus, these studies indicate that 25(OH)D levels would be better assessed along with PTH levels.

We suggest that studies analyzing the association between Vitamin D deficiency and skeletal and extraskeletal disorders should better define subjects with true Vitamin D deficiency.

AYSE NUR TUFAN, MD, Department of Rheumatology, Haseki Training and Research Hospital; OZLEM SOYLUK, MD, Assistant Professor, Department of Endocrinology, Istanbul Faculty of Medicine, Istanbul University; FATİH TUFAN, MD, Associate Professor Department of Geriatrics, Istanbul Faculty of Medicine, Istanbul University, Istanbul, Turkey. Address correspondence to Dr. F. Tufan, Istanbul University, Istanbul Faculty of Medicine, Department of Geriatrics, Fatih, P.B. 34093, Istanbul, Turkey. E-mail: drfatih@istanbul.edu.tr

REFERENCES

1. Koeckhoven E, van der Leeden M, Roorda LD, van Schoor NM, Lips P, de Zwart A, et al. The association between serum 25-hydroxy vitamin D level and upper leg strength in patients with knee osteoarthritis: results of the Amsterdam Osteoarthritis Cohort. *J Rheumatol* 2016;43:1400-5.
2. Hansen KE, Johnson RE, Chambers KR, Johnson MG, Lemon CC, Vo TN, et al. Treatment of vitamin D insufficiency in postmenopausal women: a randomized clinical trial. *JAMA Intern Med* 2015;175:1612-21.
3. Visser M, Deeg DJ, Lips P; Longitudinal Aging Study Amsterdam. Low vitamin D and high parathyroid hormone levels as determinants of loss of muscle strength and muscle mass (sarcopenia): the Longitudinal Aging Study Amsterdam. *J Clin Endocrinol Metab* 2003;88:5766-72.
4. Jin X, Jones G, Cicuttini F, Wluka A, Zhu Z, Han W, et al. Effect of vitamin D supplementation on tibial cartilage volume and knee pain among patients with symptomatic knee osteoarthritis: a randomized clinical trial. *JAMA* 2016;315:1005-13.

J Rheumatol 2016;43:11; doi:10.3899/jrheum.160600