Dr. Elsaman, *et al* reply

To the Editor:

The theory that pain in osteoarthritis (OA) is due to effusion is not answered in our study¹. All the patients included in our study initially had effusion on clinical and ultrasound examination¹. To prove the theory that pain in OA had been due to effusion, we would have had to include patients with painful osteoarthritic knee and examine all of them by ultrasound to detect the cause of pain.

The ratio of painful osteoarthritic knee effusion ranges from 44% to 79%^{2,3}. However, 16% of osteoarthritic knee effusion is painless⁴. It can be stated that many painful osteoarthritic knees had effusion, but others may have had bursitis or associated activity of crystal-induced arthritis or severe thinning of the knee cartilage or more rarely, degenerative meniscal tear.

According to the results of our study we can say that most of the effusion is painful, but we cannot confirm that every painful osteoarthritic knee is due to effusion. Pain in OA is multifactorial, and effusion plays a crucial role.

The theory that effusion is due to mechanical and not inflammatory cause in OA has been supported in our study because reduction of effusion thickness using spironolactone was maintained after stoppage of treatment for 2 weeks. Further, the addition of power Doppler examination to the painful osteoarthritic knee could clarify more the inflammatory or mechanical nature of effusion. If the cause of effusion is inflammatory, we expect marked power Doppler activity.

AHMED ELSAMAN, MD, consultant and lecturer in rheumatology and rehabilitation, Sohag University; AHMED RADWAN, MD, consultant and

lecturer in rheumatology and rehabilitation, Sohag University; WALAA MOHAMMED, MD, consultant and lecturer in pharmacology, Sohag University, Sohag, Egypt; SARAH OHRNDORF, MD, specialist in Internal Medicine/Rheumatology, Department of Rheumatology and Clinical Immunology, Charité Universitätsmedizin Berlin, Berlin, Germany. Address correspondence to Dr. S. Ohrndorf, MD, Department of Rheumatology and Clinical Immunology, Charité Universitätsmedizin Berlin, Charitéplatz 1, Berlin, Germany. E-mail: sarah.ohrndorf@charite.de

REFERENCES

1. Elsaman AM, Radwan AR, Mohammed WI, Ohrndorf S. Low-dose spironolactone: treatment for osteoarthritis-related knee effusion. A prospective clinical and sonographic-based study. *J Rheumatol* 2016;43:1114-20.
2. D'Agostino MA, Conaghan P, Le Bars M, Baron G, Grassi W, Martin-Mola E, et al. EULAR report on the use of ultrasonography in painful knee osteoarthritis. Part 1: prevalence of inflammation in osteoarthritis. *Ann Rheum Dis* 2005;64:1703-9.
3. de Miguel Mendieta E, Ibanez TC, Jaeger JU, Hernán GB, Mola EM. Clinical and ultrasonographic findings related to knee pain in osteoarthritis. *Osteoarthritis Cartilage* 2006;14:540-4.
4. Hill CL, Gale DG, Chaisson CE, Skinner K, Kazis L, Gale ME, et al. Knee effusions, popliteal cysts, and synovial thickening: association with knee pain in osteoarthritis. *J Rheumatol* 2001;28:1330-7.

J Rheumatol 2016;43:12; doi:10.3899/jrheum.161169