

Genetic Variants That Are Associated with Neuropsychiatric Systemic Lupus Erythematosus

Roger C. Ho, Huiyi Ong, Chandra Thiaghu, Yanxia Lu, Cyrus S. Ho, and Melvyn W. Zhang

ABSTRACT. *Objective.* While genetic risks have been implicated in systemic lupus erythematosus (SLE), the involvement of various genotypes in neuropsychiatric SLE (NPSLE) remains uncertain. The present metaanalysis aimed to combine data from different studies and evaluate the association between each genotype and the risk of developing NPSLE.

Methods. Studies were searched and retrieved from online databases (PubMed, EMBASE, BIOSIS, and ScienceDirect). Case-control studies were chosen if they reported genotype frequencies of the γ Fc region (FC γ R) receptors II-A, III-A, and III-B; tumor necrosis factor- α (TNF- α); mannan-binding lectin (MBL); integrin alpha M (ITGAM); interleukin (IL) 1, IL-1 β , and IL-6; IL-10 promoter; and vitamin D genes. The OR were used to assess the strength of this association between patients with NPSLE and SLE.

Results. A total of 33 studies were considered in this metaanalysis. The results suggest that these genotypes demonstrated a significant association with NPSLE: the homozygous FC γ R IIIa 158 FF genotype (OR 1.89, $p = 0.03$ for FF vs VV + FV), heterozygous FC γ R IIIb NA1/2 genotype (OR 2.14, $p = 0.03$ for NA1/2 vs NA1/1; OR 1.81, $p = 0.04$ for NA1/2 vs NA1/1 + NA2/2), and homozygous ITGAM rs1143679 HH genotype (OR 3.39, $p = 0.04$ for HH vs RH; OR 3.11, $p = 0.048$ for HH vs RR + RH). Polymorphisms of the TNF- α , MBL2, IL-1, IL-1 β , IL-6, IL-10 promoter, and vitamin D receptor genes did not show a statistically significant association with the risk of developing NPSLE ($p > 0.05$).

Conclusion. This metaanalysis indicates that polymorphisms in the pathways of immune complex clearance, such as the Fc γ RIIIa, Fc γ RIIIb, and ITGAM genotypes, are potential susceptibility genes for NPSLE. (First Release January 15 2016; J Rheumatol 2016;43:541–51; doi:10.3899/jrheum.150884)

Key Indexing Terms:

NEUROPSYCHIATRIC SYSTEMIC LUPUS ERYTHEMATOSUS
GENETIC VARIANT Fc γ RIIIa Fc γ RIIIb INTEGRIN ALPHA M

Systemic lupus erythematosus (SLE) is a prototypic autoimmune disease that is characterized by a higher prevalence in women than in men (female:male 9:1), loss of immunological tolerance to self-nuclear antigens, and abnormal B cell and T cell response¹. Neuropsychiatric SLE (NPSLE) is

one of the major and most damaging presentations² of SLE. The American College of Rheumatology (ACR) research committee devised a nomenclature providing case definitions for 19 NP syndromes in SLE³. The ACR criteria include a wide range of neurological syndromes affecting the central (e.g., headache, meningitis, and seizure), peripheral (e.g., neuropathy), and autonomic nervous systems, as well as psychiatric syndromes (e.g., anxiety, cognitive dysfunction, and depression).

The heritability of SLE is suggested to be 66%, but current understandings of genetic variants explain 10–20% of SLE heritability⁴. The concordance rate of SLE in monozygotic twins (24–56%) is higher than the rate of dizygotic twins (2–4%)⁵. Similar to other psychiatric disorders such as schizophrenia, bipolar disorder, and Alzheimer's disease, SLE is a disease of polygenic inheritance. The evidence of genetic susceptibility for SLE is mainly from genome-wide association studies, which have identified 40 disease-associated genes that are linked to SLE⁴. Mok and Lau⁶ estimated that at least 4 susceptibility genes are required for the development of SLE. These susceptibility genes are classified as HLA genes (e.g., *DQA1* and *DQB1*) and non-HLA genes [e.g., γ Fc region (Fc γ) RIIA, interleukin (IL) 6, IL-10

From the Department of Psychological Medicine, Yong Loo Lin School of Medicine, National University of Singapore; Institute of Mental Health, Singapore; Department of Clinical Psychology and Psychiatry/School of Public Health, Zhejiang University College of Medicine, Hangzhou, China.

R.C. Ho, FRCP, Assistant Professor, Department of Psychological Medicine, Yong Loo Lin School of Medicine, National University of Singapore; H. Ong, MBBS, Medical Officer, Department of Psychological Medicine, Yong Loo Lin School of Medicine, National University of Singapore; C. Thiaghu, MBBS, Medical Officer, Department of Psychological Medicine, Yong Loo Lin School of Medicine, National University of Singapore; Y. Lu, PhD, Assistant Professor, Department of Clinical Psychology and Psychiatry/School of Public Health, Zhejiang University College of Medicine; C.S. Ho, MRCPsych, Specialist Registrar, Department of Psychological Medicine, Yong Loo Lin School of Medicine, National University of Singapore; M.W. Zhang, MRCPsych, Specialist Registrar, Institute of Mental Health.

Address correspondence to Dr. Y. Lu, Assistant Professor, Department of Clinical Psychology and Psychiatry/School of Public Health, Zhejiang University College of Medicine, Zijingang Campus, 866 Yuhangtang Road, Hangzhou, Zhejiang 310058, China. E-mail: A0068932@u.nus.edu

Accepted for publication October 15, 2015.

promoter, mannose-binding lectin (MBL), and plasminogen activator inhibitor 1 (PAI-1)]⁷. Genetic polymorphisms in the *IL-10* promoter, *MBL*, and *PAI-1* genes may be associated with lupus nephritis in Chinese patients with SLE^{6,8,9,10}. Chong, *et al*¹¹ proposed that *IL-10* causes the hyperactivity of B cells in SLE, and *IL-10* is implicated in the pathogenesis of NPSLE¹².

Accumulating studies suggest that susceptibility genes contribute to the NP manifestations in SLE. Koga, *et al*¹³ combined the risk alleles of 7 genes (*BLK*, *HLA-DRB1*, *FCγRIIb*, *IRF5*, *STAT4*, *TNFAIP3*, and *TNFSF13*) and found that patients with SLE carrying more than 10 risk alleles demonstrated a greater risk of neurological symptoms compared with those carrying fewer than 10 risk alleles. Guerra, *et al*⁵ reported that the *integrin alpha M* (*ITGAM*) and *FcγR* genes contributed to apoptosis, which may lead to programmed cell death in neurons and result in NP symptoms in patients with SLE. Yang, *et al*¹⁴ found that *ITGAM* was associated with severe manifestations of SLE. May, *et al*¹⁵ proposed that tumor necrosis factor-α (TNF-α) causes local damage in the brain and results in demyelination.

Metaanalyses are useful because they combine multiple studies on the same alleles of genes to enhance the statistical power of the analysis and derive more reliable results of the genetic effects^{16,17}. Previous metaanalyses studying the relationship between gene polymorphisms and SLE found that the *MBL* variants are risk factors for SLE, including exon 1 codon, 54 B, promoter -550L, and promoter -221X¹⁸, TNF-α -308 A/G polymorphism¹⁹, DD (deletion/deletion) genotype of intron 16 angiotensin-converting enzyme (ACE) gene²⁰, cytotoxic T-lymphocyte antigen 4 (CTLA-4) 1722T/C polymorphism²¹, TNF ligand superfamily member 4 (TNFSF 4) rs2205960 polymorphism¹⁷, *IL-1A* -889 C/T polymorphism²², and *IL-6* -174 G/C and -572 G/C polymorphisms²³. Supplementary Table 1 (available online at jrheum.org) summarizes the pathological roles of specific genotypes in SLE.

SLE is a multisystem disease with diverse and protean clinical features, and genetic studies focusing on specific endophenotypes may result in reduced genetic heterogeneity²⁴. The association between genetic polymorphism and susceptibility to NPSLE was not explored in previous metaanalyses. We propose that a metaanalysis of case-control association studies may identify candidate genes for NPSLE susceptibility. The primary aim of our metaanalysis was to assess the association between different genotypes and the risk of developing NPSLE. To achieve this aim, we combined results from previous studies and assessed the strength of the association between particular genotypes and the OR to develop NPSLE. The second aim of our study was to identify genes that have an influential effect on the NP manifestation of NPSLE.

MATERIALS AND METHODS

Search strategy. Studies of genotypes in SLE and NPSLE were systematically searched in the following databases from inception to February 2014:

PubMed, EMBASE, BIOSIS, and ScienceDirect. The search terms that were used were interleukin-1, *IL-1*; interleukin-2, *IL-2*; interleukin-6, *IL-6*; interleukin-8, *IL-8*; interleukin-10, *IL-10*; interleukin-18, *IL-18*; interferon regulatory factor 5, *IRF5*; tyrosine kinase 2, *TYK2*; Fc receptor-like 3, *FCRL3*; Fc gamma receptor IIb, *Fcγ receptor IIb*, *FcγRIIb*; Fc gamma receptor IIa, *Fcγ receptor IIa*, *FcγRIIa*; Fc gamma receptor IIIa, *Fcγ receptor IIIa*, *FcγRIIIa*; Fc gamma receptor IIIb, *Fcγ receptor IIIb*, *FcγRIIIb*; TRAF1-C5; *TNFSF4*; toll-like receptor 9, *TRL-9*; tumour necrosis factor-α, *TNF-α*; lymphotoxin alpha, *LTA*; endothelial nitric oxide synthase, *eNOS*; *PXK*; *BANK1*; *TNFAIP3*; *ITGAM*; *STAT4*; programmed cell-death 1, *PDCD1*; mannose-binding lectin, *MBL*; *RANTES*, angiotensin converting enzyme; *ACE*; *FAS*; *CTLA-4*; *C4*; *C2*; *C1q*; human leukocyte antigen, *HLA*; T cell receptor, *TCR*; *CR1*; *C3b*; *C4b*; Immunoglobulin Gm; *Km*; Poly (ADP-ribose) polymerase, *PARP*; heat shock protein 70, *hsp70*; vitamin D; *XRCC1*; *TREX-1*; and a combination of key words for SLE: “systemic, erythematosus, lupus, SLE, and neuropsychiatric”. We also performed a manual review of reference lists from relevant studies. The search was limited to articles with English abstracts and to articles that reported data from humans.

Inclusion criteria and exclusion criteria. Studies were included if the following criteria were met: (1) studies that used the standard criteria of the ACR to diagnose SLE and NPSLE, (2) case-control studies that reported the allele frequencies of a particular gene in patients with SLE who did and did not experience NP symptoms, (3) prospective studies that reported the allele frequencies of a particular genotype in patients with SLE who did and did not develop NP symptoms at the end of the followup period, (4) the presence of NP symptoms was assessed clinically or using standardized instruments (i.e., subjective reports of NP symptoms were not acceptable for inclusion), and (5) studies that provided adequate information to calculate the effect size and the pooled R. Studies were excluded if one of the following criteria were met: (1) no report of the allele frequencies of a particular gene, or (2) animal studies.

Data extraction. We conducted a metaanalysis in line with the Meta-Analysis of Observational Studies in Epidemiology guidelines on the synthesis of observational data²⁵. A selection of relevant publications was conducted independently by 2 researchers (OHY and CT), and any disagreements were resolved through discussions with RCH. The articles were deidentified [blinding of the title, author(s), year of publication, and journal name] before data extraction. The following information were extracted from each article and crosschecked by the second researcher: average age of the participants, the proportion of men, other moderators, and the relevant statistics describing the relationship between a particular genotype and the risk of developing NPSLE. Summary statistics, such as allele frequencies and the number of patients with SLE who were positive and negative for NP symptoms, were extracted or calculated from the original data.

Statistical analysis. All of the statistical analyses were performed with Comprehensive Meta-Analysis Version 2.0. A comparison was performed between the patients with NPSLE and SLE. The OR and 95% CI for developing NPSLE were calculated for genotype contrast. Tests of heterogeneity were conducted using the Q statistic, which is distributed as a chi-square variate under the assumption of the homogeneity of effect sizes. The between-study heterogeneity was assessed with the *I*² statistic and p value. Random-effects model was used when there was significant between-study heterogeneity. Fixed-effects model was chosen to calculate the average OR across the relevant studies when heterogeneity was not statistically significant because the random-effects model assumes a markedly conservative null-hypothesis model in genetic metaanalysis²⁶. Finally, Egger's regression test was performed to test for evidence of publication bias.

RESULTS

Among an initial 1938 potentially relevant articles, 52 articles met our inclusion criteria, of which 19 did not report relevant data to calculate the effect size. Finally, we included 33 articles with 905 patients with NPSLE and 4928 patients with

Table 1. Characteristics of the included studies for this metaanalysis.

Studies	Identified Genes	Hardy-Weinberg Equilibrium	Country of Origin	Ethnicity of Participants	No. Patients with NPSLE	No. Patients with SLE	Method
Tsai, <i>et al</i> ²⁷	<i>IL1RN, IL1β -511, TNF-α -308</i>	Control yes, patients no	Taiwan	100% Chinese	35	87	PCR
Tahmasebi, <i>et al</i> ²⁸	<i>IL1RN</i>	Yes	Iran	100% Iranian	16	205	PCR
Huang, <i>et al</i> ²⁹	<i>IL1β -511</i>	No	Taiwan	100% Chinese	9	52	PCR
Santos, <i>et al</i> ³⁰	<i>IL6 -174</i>	Yes	Portugal	100% white	10	115	PCR
Schotte, <i>et al</i> ³¹	<i>IL6 -174</i>	Yes	Germany	100% white	62	211	PCR
Hristova, <i>et al</i> ³²	<i>IL6 -174</i>	Yes	Bulgaria	100% white	10	52	PCR
Asano, <i>et al</i> ³³	<i>IL6 -174</i>	Yes	Brazil	85% African, 15% European	54	80	PCR
Rood, <i>et al</i> ¹²	<i>IL-10 -1082 GA, -818 CT, -592 CA</i>	Yes	Netherlands	100% Dutch	42	50	PCR
Hirankarn, <i>et al</i> ³⁴	<i>IL-10 -1082 GA, -818 CT, -592 CA</i>	Yes	Thai	100% Thai	10	230	PCR
Toller-Kawahisa, <i>et al</i> ³⁵	<i>ITGAM (rs1143679)</i>	Yes	Brazil	100% Brazilian	51	157	PCR
Warchol, <i>et al</i> ³⁶	<i>ITGAM (rs1143679)</i>	Yes	Poland	100% white	32	154	PCR
Monticelo, <i>et al</i> ³⁷	<i>MBL-2</i>	Yes	Brazil	76.1% European, 23.9% African	38	327	PCR
Tsai, <i>et al</i> ³⁸	<i>MBL-2</i>	Yes	Taiwan	100% Chinese	32	150	PCR
Piao, <i>et al</i> ³⁹	<i>MBL-2</i>	Yes	North America	83.8% white, 16.2% other races	21	130	PCR
Garred, <i>et al</i> ⁴⁰	<i>MBL-2</i>	Yes	Denmark	100% white	4	91	PCR
Garred, <i>et al</i> ⁴¹	<i>MBL-2</i>	Yes	Denmark	98% white	9	99	PCR
Sandrin-Garcia, <i>et al</i> ⁴²	<i>MBL-2, MBL-2 promoter</i>	Yes	Brazil	85% European, 15% African	27	134	PCR
Jakab, <i>et al</i> ⁴³	<i>MBL-2, MBL-2 promoter</i>	Yes	Hungary	100% white	116	252	PCR
Rood, <i>et al</i> ⁴⁴	<i>TNF-α -308</i>	Yes	Netherlands	100% white	43	99	PCR
Lin, <i>et al</i> ⁴⁵	<i>TNF-α -308</i>	Yes	Taiwan	100% Chinese	17	161	PCR
Santos, <i>et al</i> ³⁰	<i>TNF-α -308, LTA 252</i>	Yes	Portugal	100% white	10	115	PCR
Ahmed, <i>et al</i> ⁴⁶	<i>TNF-α -308, LTA 252</i>	Yes	Egypt	100% Egyptian	16	100	PCR
May, <i>et al</i> ¹⁵	<i>TNF-α -308</i>	Yes	Australia	99% white, 1% other	17	64	PCR
Angelo, <i>et al</i> ⁴⁷	<i>TNF-α -308</i>	Control yes, patients no	Brazil	100% Brazilian	26	98	PCR
Dijstelbloem, <i>et al</i> ⁴⁸	<i>FcγRIIa</i>	Yes	Netherlands	100% white	20	230	PCR
Chen, <i>et al</i> ⁴⁹	<i>FcγRIIa, FcγRIIIa, FcγRIIIb</i>	No	Taiwan	100% Chinese	36	302	PCR
Hong, <i>et al</i> ⁵⁰	<i>FcγRIIIa, FcγRIIIb</i>	Yes	Korea	100% Korean	19	183	PCR
Hatta, <i>et al</i> ⁵¹	<i>FcγRIIa, FcγRIIIa, FcγRIIIb</i>	Yes	Japan	100% Japanese	22	81	PCR
Warchol, <i>et al</i> ⁵²	<i>XRCC1</i>	Yes	Poland	100% white	53	265	PCR
Lin, <i>et al</i> ⁵³	<i>XRCC1</i>	Yes	Taiwan	100% Chinese	14	164	PCR
Huang, <i>et al</i> ⁵⁴	<i>Vitamin D receptor Fok I</i>	Yes	Taiwan	100% Chinese	9	52	PCR
Sakulpipatsin, <i>et al</i> ⁵⁵	<i>Vitamin D receptor BsmI</i>	Yes	Thailand	100% Thai	18	101	PCR
Luo, <i>et al</i> ⁵⁶	<i>Vitamin D receptor BsmI</i>	Yes	China	100% Chinese	7	337	PCR

SLE: systemic lupus erythematosus; NPSLE: neuropsychiatric SLE; IL: interleukin; ITGAM: integrin alpha M; MBL: mannan-binding lectin; TNF-α: tumor necrosis factor-α.

SLE without NP symptoms in our analysis (Figure 1). Table 1 summarizes the characteristics of the included studies for our metaanalysis.

A summary of the metaanalysis findings on the relationship between all of the gene polymorphisms and NPSLE is given in Table 2.

FCγR IIIa gene polymorphism and NPSLE association. When the FF homozygotes were contrasted with VV and FV combined, there was a significant association between the homozygous FF genotype and the risk of NPSLE (OR 1.89, 95% CI 1.08–3.30, $p = 0.03$; Figure 2A), with no evidence of significant between-study heterogeneity ($I^2 = 0$, $p = 0.61$).

FCγR IIIb gene polymorphism and NPSLE association. When NA1/2 was contrasted with NA1/1, a significant association between the heterozygous NA1/2 genotype and the risk of NPSLE was found (OR 2.14, 95% CI 1.07–4.30, $p = 0.03$; Figure 2B), with no evidence of significant between-study heterogeneity ($I^2 = 52.38$, $p = 0.15$). When

NA1/2 was contrasted with NA1/1 and NA2/2 combined, a significant association between the heterozygous NA1/2 genotype and the risk of NPSLE was found (OR 1.81, 95% CI 1.02–3.20, $p = 0.04$; Figure 2C), with no evidence of significant between-study heterogeneity ($I^2 = 29.10$, $p = 0.24$).

ITGAM gene polymorphism and NPSLE association. When HH was contrasted with RH, a significant association between the homozygous HH genotype and the risk of NPSLE was found (OR 3.39, 95% CI 1.04–11.03, $p = 0.04$; Figure 3A), with no evidence of significant between-study heterogeneity ($I^2 = 0$, $p = 0.98$). When HH was contrasted with RR and HH combined, a significant association between the homozygous HH genotype and risk of NPSLE was found (OR 3.11, 95% CI 1.01–9.59, $p = 0.048$; Figure 3B), with no evidence of significant between-study heterogeneity ($I^2 = 0$, $p = 0.77$).

Negative findings in the gene polymorphisms and NPSLE




Figure 1. Flowchart describing the process of study selection.

association. We did not find any association between NPSLE and MBL2 rs7096206, MBL2 Exon1, LTA 252, IL-1 β -511, IL-1 RN, IL-6 -174, IL-10 promoter haplotypes (-1082 GA, -819CT, -592CA), and vitamin D receptor BsmI gene polymorphisms ($p > 0.05$).

Heterogeneity and publication bias. There was no statistically significant heterogeneity between studies. The I^2 value varied from 0% to 73.90%, and the p value varied from 0.05 to 0.95. There was evidence of publication bias in 2 analyses (TNF- α A-308G AA vs AG, Egger's regression coefficient = -12.12, $p = 0.03$; IL-6 -174GC GG+GC vs CC, Egger's regression coefficient = 0.93, $p = 0.0068$). Statistically, it was not possible to assess the publication bias for analyses that involved fewer than 3 studies.

DISCUSSION

To our knowledge, ours is the first metaanalysis investigating the association between various genetic polymorphisms and NPSLE. Our metaanalysis demonstrated that the homozygous Fc γ RIIIa 158FF genotype, the heterozygous Fc γ IIIb NA1/2 genotype, and the homozygous ITGAM HH genotype may be associated with the susceptibility to NPSLE. The

overall homozygosity for the HH genotype of the *ITGAM* rs1143679 gene demonstrated the highest risk for NPSLE compared with the heterozygous RH genotype, with a 3.4-fold greater risk for the development of NPSLE. Our metaanalysis highlights several important unresolved issues in the genetics of NPSLE. Future studies of NPSLE may focus on the above genes, which are related to pathways of the immune complex clearance. Our metaanalysis demonstrated no significant association of the TNF- α , MBL-2 rs7096206, MBL-Exon 1, LTA 252 rs909253, IL-6 -174GC, IL1 β -511CT, and IL-1 RN polymorphisms with NPSLE. In our metaanalysis, the between-study heterogeneity was low and nonsignificant. There was no publication bias in the significant findings (i.e., the Fc γ RIIIa, Fc γ RIIIb, and ITGAM rs1143679 genotypes).


FC γ RIII. Fc γ receptors are found in the IgG membrane and form an important link between the cellular and humoral elements of the immune system⁵⁷. Kimberly, *et al*⁵⁸ detected a defective Fc γ receptor-mediated uptake of IgG ligand-coated erythrocytes by macrophages in patients with SLE and postulated that the Fc γ receptor-mediated clearance defect is caused by underlying genetic etiology. Genetic

Table 2. Results of the metaanalysis for genetic variants and NPSLE association.


Locus	Variant Alleles	No. Studies	Polymorphism	Genetic Models	Reference Allele	No. Subjects, NPSLE/SLE	OR	Test of Association 95% CI	p	Test of Heterogeneity I ²	p	Publication Bias Egger's Regression Coefficient	p
FCγR	Ila	2	RR vs HH + RH	Dominant	R	56/476	0.74	0.35–1.56	0.42	31.58	0.23	NA	NA
	IIIa 158	2	FF vs VV + FV	Dominant	F	56/476	1.89	1.08–3.30	0.03*	0.00	0.61	NA	NA
	IIIb	2	NA 2/2 vs NA1/1	Codominant	NA2	22/233	1.69	0.69–4.10	0.25	0.00	0.39	NA	NA
		2	NA 1/2 vs NA 1/1	Codominant		45/353	2.14	1.07–4.30	0.03*	52.38	0.15	NA	NA
		2	NA 1/2 vs NA 2/2	Codominant		43/274	1.34	0.63–2.85	0.45	0.00	0.78	NA	NA
		3	NA1/1+NA1/2 vs NA 2/2	Dominant		75/640	1.13	0.63–2.04	0.68	0.00	0.80	0.63	0.84
		2	NA1/1 vs NA2/2 + NA1/2	Recessive		55/430	1.99	1.02–3.90	0.05	48.98	0.16	NA	NA
		2	NA1/2 vs NA 1/1 + NA 2/2	Over-dominant		55/430	1.81	1.02–3.20	0.04*	29.10	0.24	NA	NA
TNF-α	A-308	3	AA vs GG	Codominant	A	52/186	0.86	0.17–4.34	0.86	36.56	0.21	-10.03	0.14
		3	AA vs AG	Codominant		18/65	0.86	0.16–4.66	0.86	1.54	0.36	-12.12	0.03*
		3	GG vs AG	Codominant		68/235	1.17	0.59–2.30	0.65	0.00	0.84	-4.09	0.13
		7	AA + AG vs GG	Recessive		154/570	0.80	0.54–1.19	0.26	0.00	0.70	3.55	0.06
		3	AA vs GG + AG	Dominant		69/243	0.89	0.18–4.45	0.89	32.07	0.23	-9.96	0.12
		3	AA + GG vs AG	Over-dominant		69/243	1.10	0.56–2.15	0.79	0.00	0.91	-2.65	0.33
MBL2	rs7096206	2	XX vs YY	Codominant	Y	100/186	1.41	0.58–3.43	0.45	0.00	0.37	NA	NA
		2	XX vs XY	Codominant		48/74	1.63	0.64–4.16	0.31	0.00	0.99	NA	NA
		2	YY vs XY	Codominant		123/241	1.09	0.66–1.78	0.74	71.90	0.06	NA	NA
		2	XX + XY vs YY	Dominant		134/252	1.00	0.63–1.60	0.99	73.90	0.05	NA	NA
		2	XX vs XY + YY	Recessive		134/252	1.50	0.62–3.59	0.37	0.00	0.49	NA	NA
		2	XX + YY vs XY	Over-dominant		134/252	1.23	0.75–2.01	0.41	75.63	0.04	NA	NA
MBL2	Exon 1	7	AA vs OO	Codominant	O	135/501	0.65	0.36–1.17	0.15	0.00	0.95	0.53	0.29
		7	AA vs AO	Codominant		194/757	1.03	0.71–1.43	0.99	0.66	0.42	-0.53	0.63
		7	OO vs AO	Codominant		101/378	1.45	0.80–2.63	0.21	0.00	0.89	-0.62	0.30
		7	AA + AO vs OO	Dominant		215/818	0.67	0.38–1.17	0.16	0.00	0.96	0.57	0.21
		8	AA vs AO + OO	Recessive		247/939	0.87	0.64–1.18	0.36	8.50	0.36	-0.13	0.90
		7	AA + OO vs AO	Over-dominant		215/818	1.04	0.74–1.45	0.82	0.00	0.43	-0.66	0.53
ITGAM	rs1143679	2	HH vs RR	Codominant	R	56/147	2.94	0.94–9.23	0.06	0.00	0.65	NA	NA
		2	RR vs RH	Codominant		76/222	1.01	0.71–1.42	0.98	0.66	0.42	NA	NA
		2	HH vs RH	Codominant		34/87	3.39	1.04–11.03	0.04*	0.00	0.98	NA	NA
		2	HH vs RR + RH	Recessive		83/228	3.11	1.01–9.59	0.048*	0.00	0.77	NA	NA
		2	RR vs HH + RH	Dominant		83/228	0.94	0.56–1.57	0.80	0.00	0.38	NA	NA
		2	RR + HH vs RH	Over-dominant		83/228	1.18	0.69–2.02	0.54	0.00	0.32	NA	NA
LTA 252	rs909253	2	AA vs GG + AG	Dominant	A	26/189	0.56	0.23–1.38	0.21	0.00	0.60	NA	NA
IL-1	RN	2	CC vs TT	Codominant	T	28/157	1.42	0.47–4.27	0.53	0.00	0.66	NA	NA
		2	CC vs CT	Codominant		34/106	0.91	0.35–2.36	0.85	0.00	0.81	NA	NA
		2	TT vs CT	Codominant		40/219	0.73	0.34–1.56	0.41	0.00	0.65	NA	NA
		2	CC + CT vs TT	Dominant		51/241	1.31	0.63–2.73	0.47	0.00	0.56	NA	NA
		2	CC vs CT + TT	Recessive		51/241	1.09	0.45–2.64	0.86	0.00	0.78	NA	NA
		2	CC + TT vs CT	Over-dominant		51/241	0.79	0.41–1.54	0.49	0.00	0.95	NA	NA
IL-1β	-511	2	CC vs TT	Codominant	C	27/59	0.81	0.30–2.16	0.67	0.00	0.97	NA	NA
		2	CC vs CT	Codominant		26/60	0.81	0.30–2.17	0.67	0.00	0.87	NA	NA
		2	TT vs CT	Codominant		35/71	1.01	0.44–2.33	0.99	0.00	0.89	NA	NA
		2	CC + CT vs TT	Recessive		44/95	0.92	0.43–1.95	0.82	0.00	0.94	NA	NA
		2	CC vs CT+TT	Dominant		44/95	0.81	0.33–1.97	0.64	0.00	0.91	NA	NA
		2	CC + TT vs CT	Over-dominant		44/95	0.93	0.44–1.98	0.85	0.00	0.84	NA	NA
IL-6	-174	3	GG vs CC	Codominant	C	46/167	0.64	0.31–1.32	0.23	0.00	0.82	0.74	0.36
		3	GG vs GC	Codominant		65/254	0.87	0.49–1.55	0.63	0.00	0.65	-0.08	0.96
		3	CC vs GC	Codominant		53/171	1.44	0.71–2.94	0.31	0.00	0.75	-0.75	0.42
		3	GG + GC vs CC	Dominant		82/296	0.68	0.35–1.30	0.24	0.00	0.78	0.93	0.0068**
		4	GG vs GC + CC	Recessive		136/322	0.92	0.57–1.47	0.72	0.00	0.61	0.44	0.78
		3	GG + CC vs GC	Over-dominant		82/296	1.01	0.61–1.70	0.96	0.00	0.58	-0.47	0.78
IL-10 promoter gene	-1082 -818 -592	2	GCC vs non GCC	NA	Non GCC haplotype	52/270	0.44	0.17–1.13	0.09	0.00	0.83	NA	NA
Vitamin D receptor gene	BsmI	3	B allele positive vs B allele negative	NA	B allele positive	32/453	0.79	0.21–2.91	0.72	36.11	0.21	2.05	0.77

* $p < 0.05$. ** $p < 0.01$. SLE: systemic lupus erythematosus; NPSLE: neuropsychiatric SLE; TNF-α: tumor necrosis factor-α; MBL: mannan-binding lectin; ITGAM: integrin alpha M; LTA: lymphotoxin alpha; IL: interleukin; NA: not available because the metaanalysis program requires at least 3 studies to calculate Egger's regression coefficient for publication bias.

A.



B.



C.





Figure 2. Forest plots describing the metaanalysis of the association between (A) the FCγR IIIa 158 V/F polymorphism (FF vs VV + FV), (B) FCγR IIIb NA 1/2 polymorphism (NA 1/2 vs NA 1/1), and (C) FCγR IIIb NA 1/2 polymorphism (NA 1/2 vs NA 1/1+NA 2/2) and the risk of neuropsychiatric systemic lupus erythematosus. FCγR: γ Fc region.

A.



B.




Figure 3. Forest plot describing the metaanalysis of the association between the ITGAM rs1143679 HR polymorphism (HH vs RR + RH) and the risk of NPSLE. (A) HH versus RH, and (B) HH versus RR + RH. ITGAM: integrin alpha M; NPSLE: neuropsychiatric systemic lupus erythematosus.

polymorphisms affect the structure and function of Fcγ receptors. There are 3 different isoforms of Fcγ receptors (Fcγ RI, Fcγ RII, and Fcγ RIII), and each isoform has its own cell-anchoring mechanisms and allelic variations. Defective Fcγ receptor-mediated clearance leads to the subsequent deposition of immune complexes and contributes to the pathogenesis of SLE⁵⁹. The gene of *Fcγ RIII* is located on chromosome 1q23. For the γγ RIIIa-V/F 158 polymorphism, F is the low-binding allele and V is the high-binding allele. The homozygosity of low-affinity FF genotype is associated with defective binding of immune complexes, increased tissue deposition, and accelerated organ damage⁶⁰. VV

homozygotes are more likely to bind to IG1 and IG3 compared with VF heterozygotes and FF homozygotes⁶¹. Li, *et al*⁶² performed a metaanalysis on the involvement of the Fcγ receptor IIIA-V/F 158 polymorphism in SLE and found that the homozygosity of FF genotype was associated with lupus nephritis and SLE. Our findings suggest that the homozygosity of FF genotype is associated with NP symptoms.

The Fcγ IIIb receptor is a low-affinity receptor that is expressed in neutrophils and that removes small immune complexes from the circulation. There are bi-allelic neutrophil-specific antigens (NA1 and NA2) in the *Fcγ IIIb* receptor gene. The NA1 allele is associated with autoimmune

neutropenia⁶³, and the NA2 allele is associated with a reduction in the capacity for phagocytosis⁶⁴. There are significant ethnic differences in allele frequencies among patients with SLE. NA1 is the prominent allele in Chinese and Malay patients⁶⁵, while NA2 is the prominent allele in Indian patients⁵⁷. The FcγRIIIb NA1/NA2 heterozygote demonstrates a reduction in the affinity for its ligand, a higher risk of encapsulated bacterial infections, and association with lupus nephritis in Japanese patients and with endstage renal disease in Chinese patients^{51,66,67}.

ITGAM. CD11b-integrin (ITGAM) is a member of the immune-complex processing pathway that enhances the adherence of neutrophils and monocytes to stimulated endothelium in the phagocytosis of complement cleavage fragment particles. ITGAM encodes integrin-αM (CD11b+), which is a molecule that combines with integrin-β2 to form a leukocyte-specific integrin¹⁴. The αMβ2-integrin forms cell surface receptors on monocytes and neutrophils to bind intracellular adhesion molecule 1⁶⁸. The *ITGAM* gene is associated with SLE susceptibility in Europeans⁶⁹, Hong Kong Chinese, and Thai¹⁴. The ITGAM genotype polymorphism was proposed to be one of the strongest genetic risk factors⁷⁰ and predicts disease manifestations in SLE³⁵. The rs1143679 variant of the *ITGAM* gene encodes an arginine-to-histidine amino acid change at position 77 (R77H) in the β propeller domain of CD11b. The presence of the H allele increases CR3 expression and aggravates the inflammatory process in patients with SLE by activating complements, antiphospholipid autoantibodies, and neutrophils, causing arterial and venous thrombosis and the risk of lupus nephritis⁷¹. In addition to renal disease and discoid rash⁷², the ITGAM genotype polymorphism may increase the risk of NP manifestations by cell-mediated immunological reactions.

Other genotypes. Our results suggest that there is no significant association of the polymorphisms of MBL2 rs7096206, exon1, LTA 252 rs909253, IL-6 -174GC, IL-1βC-511TC, and IL-1 RN with the susceptibility to develop NPSLE. IL-10 is a major immunoregulatory cytokine, and IL-10 genotypes are strongly associated with renal disease⁹. Rood, *et al*¹² found that IL-10 promoter haplotypes are implicated in the pathogenesis of NPSLE, but another study in Thai patients with SLE did not support those findings³⁴. Other studies on the IL-10 promoter genes in patients with SLE did not assess NP symptoms^{73,74,75,76}. Further studies are required to investigate the involvement of IL-10 promoter haplotypes in NPSLE.

Our metaanalysis has inherited the limitations of the case-control genetic association studies. First, the number of studies for each genotype was small, and our metaanalysis was underpowered to detect differences in some of the genotypes between patients with NPSLE and SLE. Second, Li, *et al*⁶⁰ found that the Fcγ RIII a-FF genotype was associated with a high risk of developing SLE in Asians and Europeans, but not in Africans. However, we were not able

to perform an ethnic subgroup analysis for each genetic polymorphism as a result of the small number of published studies. Third, we were not able to address intergenetic and gene-environment interactions (e.g., smoking and ultraviolet light exposure). Fourth, NPSLE consists of a broad spectrum of very different clinical manifestations. In our metaanalysis, we were not able to study the relationship between specific NP manifestation and its association with specific allele. We hope that researchers will attempt to study the association between specific NP symptom and genotype in the future. Finally, replication in multiple studies is essential for the validation of genetic associations, and metaanalysis, which combines results across multiple samples from independent studies, is a key component for genetic findings. The reliability of results is not reduced when a metaanalysis pools summary statistics from individual studies; however, unavailability of individual-level data and technical and statistical differences limit the available options for multiple-testing adjustments. Thus, our study used sample-size-weighted meta statistics, which is a common approach for combining results in metaanalysis⁵⁰.

Our metaanalysis serves as a preliminary overview of the involvement of various genes as a potential etiological factor in NPSLE and sheds light on the pathogenesis of NPSLE. Our results also provide a foundation for future genetics studies to focus on endophenotypes. Boackle⁷⁷ proposed the use of clinical features such as NP status to classify patients with SLE and to reduce genetic heterogeneity. The findings in our metaanalysis require further replication. Further longitudinal studies with larger sample sizes and different ethnicities are required to study the interaction between the Fcγ RIIIa, Fcγ RIIIb, and ITGAM genotypes in the causation of NPSLE. With the publication of more genetic studies in NPSLE, a future metaanalysis should attempt to combine the cumulative number of risk alleles and to assess the risk of developing NP complications in patients with SLE.

Our metaanalysis suggests that polymorphisms in the pathways of immune complex clearance might contribute to the susceptibility of NPSLE. The homozygous FcγR IIIa -158 FF genotype, the heterozygous FcγR IIIb NA1/2 genotype, and the homozygous ITGAMHH genotype were associated with NPSLE. The defects in the immune complex clearance pathway may shed light on new therapeutic targets for NPSLE. Further genetic studies focusing on the NP endophenotype with larger sample sizes are required to replicate the findings of our metaanalysis.

ONLINE SUPPLEMENT

Supplementary data for this article are available online at jrheum.org.

REFERENCES

1. Enocsson H, Sjöwall C, Wirestam L, Dahle C, Kastbom A, Rönnelid J, *et al*. Four anti-dsDNA antibody assays in relation to systemic lupus erythematosus disease specificity and activity. *J Rheumatol* 2015;42:817-25.

2. Mak A, Ho RC, Lau CS. Clinical implications of neuropsychiatric systemic lupus erythematosus. *Adv Psychiatr Treat* 2009;15:451-8.
3. The American College of Rheumatology nomenclature and case definitions for neuropsychiatric lupus syndromes. *Arthritis Rheum* 1999;42:599-608.
4. Frangou EA, Bertsias GK, Boumpas DT. Gene expression and regulation in systemic lupus erythematosus. *Eur J Clin Invest* 2013 Oct;43:1084-96.
5. Guerra SG, Vyse TJ, Cunninghame Graham DS. The genetics of lupus: a functional perspective. *Arthritis Res Ther* 2012;14:211.
6. Mok CC, Lau CS. Pathogenesis of systemic lupus erythematosus. *J Clin Pathol* 2003;56:481-90.
7. Tsao BP. An update on genetic studies of systemic lupus erythematosus. *Curr Rheumatol Rep* 2002;4:359-67.
8. Lau YL, Lau CS, Chan SY, Karlberg J, Turner MW. Mannose-binding protein in Chinese patients with systemic lupus erythematosus. *Arthritis Rheum* 1996;39:706-8.
9. Mok CC, Lanchbury JS, Chan DW, Lau CS. Interleukin-10 promoter polymorphisms in Southern Chinese patients with systemic lupus erythematosus. *Arthritis Rheum* 1998;41:1090-5.
10. Wang AY, Poon P, Lai FM, Yu L, Choi PC, Lui SF, et al. Plasminogen activator inhibitor-1 gene polymorphism 4G/4G genotype and lupus nephritis in Chinese patients. *Kidney Int* 2001;59:1520-8.
11. Chong WP, Ip WK, Wong WH, Lau CS, Chan TM, Lau YL. Association of interleukin-10 promoter polymorphisms with systemic lupus erythematosus. *Genes Immun* 2004;5:484-92.
12. Rood MJ, Keijsers V, van der Linden MW, Tong TQ, Borggreve SE, Verweij CL, et al. Neuropsychiatric systemic lupus erythematosus is associated with imbalance in interleukin 10 promoter haplotypes. *Ann Rheum Dis* 1999;58:85-9.
13. Koga M, Kawasaki A, Ito I, Furuya T, Ohashi J, Kyogoku C, et al. Cumulative association of eight susceptibility genes with systemic lupus erythematosus in a Japanese female population. *J Hum Genet* 2011;56:503-7.
14. Yang W, Zhao M, Hirankarn N, Lau CS, Mok CC, Chan TM, et al. ITGAM is associated with disease susceptibility and renal nephritis of systemic lupus erythematosus in Hong Kong Chinese and Thai. *Hum Mol Genet* 2009;18:2063-70.
15. May LA, Huang Q, Morris D, Danis V, Manolios N. Relationship of tumour necrosis factor alpha gene polymorphisms and neuropsychiatric lupus. *Lupus* 2002;11:114-8.
16. Sestak AL, Nath SK, Sawalha AH, Harley JB. Current status of lupus genetics. *Arthritis Res Ther* 2007;9:210.
17. Lu MM, Xu WD, Yang J, Ye QL, Feng CC, Li J, et al. Association of TNFSF4 polymorphisms with systemic lupus erythematosus: a meta-analysis. *Mod Rheumatol* 2013;23:686-93.
18. Lee YH, Witte T, Momot T, Schmidt RE, Kaufman KM, Harley JB, et al. The mannose-binding lectin gene polymorphisms and systemic lupus erythematosus: two case-control studies and a meta-analysis. *Arthritis Rheum* 2005;52:3966-74.
19. Lee YH, Harley JB, Nath SK. Meta-analysis of TNF-alpha promoter -308 A/G polymorphism and SLE susceptibility. *Eur J Hum Genet* 2006;14:364-71.
20. Lee YH, Choi SJ, Ji JD, Song GG. Association between the angiotensin-converting enzyme insertion/deletion polymorphism and susceptibility to systemic lupus erythematosus: a meta-analysis. *J Renin Angiotensin Aldosterone Syst* 2013;14:248-54.
21. Zhai JX, Zou LW, Zhang ZX, Fan WJ, Wang HY, Liu T, et al. CTLA-4 polymorphisms and systemic lupus erythematosus (SLE): a meta-analysis. *Mol Biol Rep* 2013;40:5213-23.
22. Song GG, Kim JH, Seo YH, Choi SJ, Ji JD, Lee YH. Associations between interleukin 1 polymorphisms and susceptibility to systemic lupus erythematosus: a meta-analysis. *Hum Immunol* 2014; 75:105-12.
23. Yang Z, Liang Y, Qin B, Zhong R. A meta-analysis of the association of IL-6 -174 G/C and -572 G/C polymorphisms with systemic lupus erythematosus risk. *Rheumatol Int* 2014;34:199-205.
24. Steiman AJ, Urowitz MB, Ibañez D, Li TT, Gladman DD, Wither J. Anti-dsDNA and Antichromatin Antibody Isotypes in Serologically Active Clinically Quiescent Systemic Lupus Erythematosus. *J Rheumatol* 2015;42:810-6.
25. Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA* 2000;283:2008-12.
26. Han B, Eskin E. Random-effects model aimed at discovering associations in meta-analysis of genome-wide association studies. *Am J Hum Genet* 2011;88:586-98.
27. Tsai LJ, Hsiao SH, Tsai JJ, Lin CY, Tsai LM, Lan JL. Higher genetic susceptibility to inflammation in mild disease activity of systemic lupus erythematosus. *Rheumatol Int* 2009;29:1001-11.
28. Tahmasebi Z, Akbarian M, Mirkazemi S, Shahlaee A, Alizadeh Z, Amirzargar AA, et al. Interleukin-1 gene cluster and IL-1 receptor polymorphisms in Iranian patients with systemic lupus erythematosus. *Rheumatol Int* 2013;33:2591-6.
29. Huang CM, Wu MC, Wu JY, Tsai FJ. Interleukin-1 receptor antagonist gene polymorphism in Chinese patients with systemic lupus erythematosus. *Clin Rheumatol* 2002;21:255-7.
30. Santos MJ, Fernandes D, Capela S, da Silva JC, Fonseca JE. Interleukin-6 promoter polymorphism -174 G/C is associated with nephritis in Portuguese Caucasian systemic lupus erythematosus patients. *Clin Rheumatol* 2011;30:409-13.
31. Schotte H, Schluter B, Rust S, Assmann G, Domschke W, Gaubitz M. Interleukin-6 promoter polymorphism (-174 G/C) in Caucasian German patients with systemic lupus erythematosus. *Rheumatology* 2001;40:393-400.
32. Hristova M, Dourmishev L, Kamenarska Z, Nikolova S, Kaneva R, Vinkov A, et al. Role of the promoter polymorphism IL-6 -174G/C in dermatomyositis and systemic lupus erythematosus. *Biomed Res Int* 2013;2013:315365.
33. Asano NM, Angelo HD, da Silva HA, Maia MM, Lins OG, Souza PE. Interleukin-6 promoter polymorphisms -174 G/C in Brazilian patients with systemic lupus erythematosus. *Hum Immunol* 2013;74:1153-6.
34. Hirankarn N, Wongpiyabovorn J, Hanvivatvong O, Netsawang J, Akkasilpa S, Wongchinsri J, et al. The synergistic effect of FC gamma receptor IIa and interleukin-10 genes on the risk to develop systemic lupus erythematosus in Thai population. *Tissue Antigens* 2006;68:399-406.
35. Toller-Kawahisa JE, Vigato-Ferreira IC, Pancoto JA, Mendes-Junior CT, Martinez EZ, Palomino GM, et al. The variant of CD11b, rs1143679 within ITGAM, is associated with systemic lupus erythematosus and clinical manifestations in Brazilian patients. *Hum Immunol* 2014;75:119-23.
36. Warchol T, Lianeri M, Łacki JK, Olesińska M, Jagodziński PP. ITGAM Arg77His is associated with disease susceptibility, arthritis, and renal symptoms in systemic lupus erythematosus patients from a sample of the Polish population. *DNA Cell Biol* 2011;30:33-8.
37. Monticeliolo OA, Chies JA, Mucenic T, Rucatti GG, Júnior JM, da Silva GK, et al. Mannose-binding lectin gene polymorphisms in Brazilian patients with systemic lupus erythematosus. *Lupus* 2010;19:280-7.
38. Tsai YC, Yao TC, Kuo ML, Cheng TT, Huang JL. Lack of association of mannose-binding lectin gene polymorphisms with development and clinical manifestations of systemic lupus erythematosus in Chinese children. *Lupus* 2009;18:372-6.
39. Piao W, Liu CC, Kao AH, Manzi S, Vogt MT, Ruffing MJ, et al. Mannose-binding lectin is a disease-modifying factor in North

- American patients with systemic lupus erythematosus. *J Rheumatol* 2007;34:1506-13.
40. Garred P, Pressler T, Madsen HO, Frederiksen B, Svejgaard A, Hoiby N, et al. Association of mannose-binding lectin gene heterogeneity with severity of lung disease and survival in cystic fibrosis. *J Clin Invest* 1999;104:431-7.
 41. Garred P, Voss A, Madsen HO, Junker P. Association of mannose-binding lectin gene variation with disease severity and infections in a population-based cohort of systemic lupus erythematosus patients. *Genes Immun* 2001;2:442-50.
 42. Sandrin-Garcia P, Brandao LA, Coelho AV, Guimaraes RL, Pancoto JA, Segat L, et al. Mannose binding lectin gene (MBL2) functional polymorphisms are associated with systemic lupus erythematosus in southern Brazilians. *Hum Immunol* 2011;72:516-21.
 43. Jakab L, Laki J, Sallai K, Temesszentandrás G, Pozsonyi T, Kalabay L, et al. Association between early onset and organ manifestations of systemic lupus erythematosus (SLE) and a down-regulating promoter polymorphism in the MBL2 gene. *Clin Immunol* 2007;125:230-6.
 44. Rood MJ, van Krugten MV, Zanelli E, van der Linden MW, Keijsers V, Schreuder GM, et al. TNF-308A and HLA-DR3 alleles contribute independently to susceptibility to systemic lupus erythematosus. *Arthritis Rheum* 2000;43:129-34.
 45. Lin YJ, Chen RH, Wan L, Sheu JC, Huang CM, Lin CW, et al. Association of TNF-alpha gene polymorphisms with systemic lupus erythematosus in Taiwanese patients. *Lupus* 2009;18:974-9.
 46. Ahmed HH, Taha FM, Darweesh Hel-S, Morsi HM. Association between TNF promoter -308 G>A and LTA 252 A>G polymorphisms and systemic lupus erythematosus. *Mol Biol Rep* 2014;41:2029-36.
 47. Angelo HD, da Silva HA, Asano NM, Muniz MT, de Mascena Diniz Maia M, de Souza PR. Tumor necrosis factor alpha promoter polymorphism -308 G/A in Brazilian patients with systemic lupus erythematosus. *Hum Immunol* 2012;73:1166-70.
 48. Dijkstra-Hoem HM, Bijl M, Fijnheer R, Scheepers RH, Oost WW, Jansen MD, et al. Fc gamma receptor polymorphisms in systemic lupus erythematosus: association with disease and in vivo clearance of immune complexes. *Arthritis Rheum* 2000;43:2793-800.
 49. Chen JY, Wang CM, Tsao KC, Chow YH, Wu JM, Li CL, et al. Fc gamma receptor IIa, IIIa, and IIIb polymorphisms of systemic lupus erythematosus in Taiwan. *Ann Rheum Dis* 2004;63:877-80.
 50. Hong CH, Lee JS, Lee HS, Bae SC, Yoo DH. The association between fc gammaRIIIB polymorphisms and systemic lupus erythematosus in Korea. *Lupus* 2005;14:346-50.
 51. Hatta Y, Tsuchiya N, Ohashi J, Matsushita M, Fujiwara K, Hagiwara K, et al. Association of Fc gamma receptor IIIB, but not of Fc gamma receptor IIA and IIIA polymorphisms with systemic lupus erythematosus in Japanese. *Genes Immun* 1999;1:53-60.
 52. Warchol T, Mostowska A, Lianeri M, Łącki JK, Jagodziński PP. XRCC1 Arg399Gln gene polymorphism and the risk of systemic lupus erythematosus in the Polish population. *DNA Cell Biol* 2012;31:50-6.
 53. Lin YJ, Wan L, Huang CM, Chen SY, Huang YC, Lai CH, et al. Polymorphisms in the DNA repair gene XRCC1 and associations with systemic lupus erythematosus risk in the Taiwanese Han Chinese population. *Lupus* 2009;18:1246-51.
 54. Huang CM, Wu MC, Wu JY, Tsai FJ. No association of vitamin D receptor gene start codon fok 1 polymorphisms in Chinese patients with systemic lupus erythematosus. *J Rheumatol* 2002;29:1211-3.
 55. Sakulpipattin W, Verasertniyom O, Nantiruj K, Totemchokchayakarn K, Lertsrisatit P, Janwityanujit S. Vitamin D receptor gene BsmI polymorphisms in Thai patients with systemic lupus erythematosus. *Arthritis Res Ther* 2006;8:R48.
 56. Luo XY, Yang MH, Wu FX, Wu LJ, Chen L, Tang Z, et al. Vitamin D receptor gene BsmI polymorphism B allele, but not BB genotype, is associated with systemic lupus erythematosus in a Han Chinese population. *Lupus* 2012;21:53-9.
 57. Pradhan V, Deshpande N, Nadkarni A, Patwardhan M, Surve P, Ghosh K. Fc gamma R IIIB polymorphisms: their association with clinical manifestations and autoantibodies in SLE patients from Western India. *Int J Rheum Dis* 2010;13:138-43.
 58. Kimberly RP, Salmon JE, Edberg JC, Gibofsky A. The role of Fc gamma receptors in mononuclear phagocyte system function. *Clin Exp Rheumatol* 1989;7 Suppl 3:S103-8.
 59. Rúa-Figueroa I, Nóvoa J, García-Laorden MI, Erausquin C, García-Bello M, Rodríguez de Castro F, et al. Clinical and immunogenetic factors associated with pneumonia in patients with systemic lupus erythematosus: a case-control study. *J Rheumatol* 2014;41:1801-7.
 60. Li X, Ptacek TS, Brown EE, Edberg JC. Fc gamma receptors: structure, function and role as genetic risk factors in SLE. *Genes Immun* 2009;10:380-9.
 61. Salmon JE, Pricop L. Human receptors for immunoglobulin G: key elements in the pathogenesis of rheumatic disease. *Arthritis Rheum* 2001;44:739-50.
 62. Li LH, Yuan H, Pan HF, Li WX, Li XP, Ye DQ. Role of the Fc gamma receptor IIIA-V/F158 polymorphism in susceptibility to systemic lupus erythematosus and lupus nephritis: a meta-analysis. *Scand J Rheumatol* 2010;39:148-54.
 63. Yates M, Hamilton LE, Elender F, Dean L, Doll H, MacGregor AJ, et al. Is etanercept 25 mg once weekly as effective as 50 mg at maintaining response in patients with ankylosing spondylitis? A randomized control trial. *J Rheumatol* 2015;42:1177-85.
 64. Salmon JE, Edberg JC, Kimberly RP. Fc gamma receptor III on human neutrophils. Allelic variants have functionally distinct capacities. *J Clin Invest* 1990;85:1287-95.
 65. Yap SN, Phipps ME, Manivasagar M, Bosco JJ. Fc gamma receptor IIIB-NA gene frequencies in patients with systemic lupus erythematosus and healthy individuals of Malay and Chinese ethnicity. *Immunol Lett* 1999;68:295-300.
 66. Bredius RG, Fijen CA, De Haas M, Kuijper EJ, Weening RS, Van de Winkel JG, et al. Role of neutrophil Fc gamma RIIa (CD32) and Fc gamma RIIb (CD16) polymorphic forms in phagocytosis of human IgG1- and IgG3-opsonized bacteria and erythrocytes. *Immunology* 1994;83:624-30.
 67. Xu G, He Q, Shou Z, Wang H, Wang R, Jiang H, et al. Association of Fc gamma receptor IIIB polymorphism with renal-allograft in Chinese. *Transpl Immunol* 2007;18:28-31.
 68. Kim-Howard X, Maiti AK, Anaya JM, Bruner GR, Brown E, Merrill JT, et al. ITGAM coding variant (rs1143679) influences the risk of renal disease, discoid rash and immunological manifestations in patients with systemic lupus erythematosus with European ancestry. *Ann Rheum Dis* 2010;69:1329-32.
 69. Nath SK, Han S, Kim-Howard X, Kelly JA, Viswanathan P, Gilkeson GS, et al. A nonsynonymous functional variant in integrin-alpha(M) (encoded by ITGAM) is associated with systemic lupus erythematosus. *Nat Genet* 2008;40:152-4.
 70. Rhodes B, Furnrohr BG, Roberts AL, Tziricotis G, Schett G, Spector TD, et al. The rs1143679 (R77H) lupus associated variant of ITGAM (CD11b) impairs complement receptor 3 mediated functions in human monocytes. *Ann Rheum Dis* 2012;71:2028-34.
 71. Cid MC. Endothelial cell biology, perivascular inflammation, and vasculitis. *Cleve Clin J Med* 2002;69 Suppl 2:SII45-9.
 72. Fagerholm SC, MacPherson M, James MJ, Sevier-Guy C, Lau CS. The CD11b-integrin (ITGAM) and systemic lupus erythematosus. *Lupus* 2013;22:657-63.
 73. Gibson AW, Edberg JC, Wu J, Westendorp RG, Huizinga TW, Kimberly RP. Novel single nucleotide polymorphisms in the distal IL-10 promoter affect IL-10 production and enhance the risk of systemic lupus erythematosus. *J Immunol* 2001;166:3915-22.

74. Khoa PD, Sugiyama T, Yokochi T. Polymorphism of interleukin-10 promoter and tumor necrosis factor receptor II in Vietnamese patients with systemic lupus erythematosus. *Clin Rheumatol* 2005;24:11-3.
75. Rosado S, Rua-Figueroa I, Vargas JA, Garcia-Laorden MI, Losada-Fernandez I, Martin-Donaire T, et al. Interleukin-10 promoter polymorphisms in patients with systemic lupus erythematosus from the Canary Islands. *Int J Immunogenet* 2008;35:235-42.
76. Sobkowiak A, Lianeri M, Wudarski M, Łacki JK, Jagodziński PP. Genetic variation in the interleukin-10 gene promoter in Polish patients with systemic lupus erythematosus. *Rheumatol Int* 2009;29:921-5.
77. Boackle SA. Advances in lupus genetics. *Curr Opin Rheumatol* 2013;25:561-8.