Lumbar Spine Chondrocalcinosis

HÉLÈNE PETIT, HP, MD, Rheumatology Department, Strasbourg University Hospital, Hautepierre Hospital; LUC MARCELLIN, LM, MD, Pathology Department, Strasbourg University Hospital, Hautepierre Hospital; EMMANUEL CHATELUS, EC, MD, Rheumatology Department, Strasbourg University Hospital, Hautepierre Hospital, Strasbourg, France. Address correspondence to Dr. H. Petit, Service de rhumatologie, Hôpital Hautepierre, 1 avenue Molière, 67098 Strasbourg, France. E-mail: helene.bauer@chru-strasbourg.fr. J Rheumatol 2017;44:1288–9; doi:10.3899/jrheum.161452. The patient gave her written informed consent for this study.

Crystal arthropathy of the lumbar facet joint is a rare but benign differential diagnosis of spondylodiscitis.

A 70-year-old woman with a recent sigmoid diverticulitis was examined for a suspicion of infectious spondylodiscitis. She had inflammatory low back pain for a month with low-grade fever and C-reactive protein (CRP) up to 3 mg/dl. Magnetic resonance imaging of the spine suggested a septic localization with T2 short-tau inversion recovery hypersignal on the left lumbar facet joints L3–L4 and L4–L5 (Figure 1).

A bone biopsy of lumbar facet joints was performed with computed tomography guidance and showed no micro-organism on bacteriological direct examination, and culture

was sterile at 48 h. CRP was 1.6 mg/dl. Pathological examination revealed that bone, cartilage, and synovial tissues contained basophilic material including numerous rhomboidal crystals birefringent in polarized light (calcium pyrophosphate crystal deposits; Figure 2). Symptoms disappeared with nonsteroidal antiinflammatory drugs.

Calcium pyrophosphate crystal deposition in the spine is well known in 2 cases: crown dens syndrome, which is a classic differential diagnosis of meningitis in older people¹, and paralyzing sciatica due to a synovial facet joint cyst filled with crystal deposits².

Septic arthritis of lumbar facet joints is rare, but is more

Figure 1. Magnetic resonance imaging on axial T2 short-tau inversion recovery sequences. Edema on L3–L4 left facet joints, suggesting septic arthritis.

Figure 2. Calcium pyrophosphate (CPP) crystal deposition in bone, cartilage, and synovial tissue. H&E coloration. (A) Without polarized light. Arrow indicates CPP crystal and arrowhead a chondrocyte. (B) With polarized light. Arrow indicates CPP crystal.

frequently described than the microcrystalline etiology. A review counted 34 cases of septic arthritis of facet joints³, whereas a single case of crystal lumbago was reported by a Japanese team, with involvement in the L4–L5 facet joints⁴.

Arthritis of lumbar facet joints could be due to crystal lumbago, if no microorganism is found.

ACKNOWLEDGMENT

We thank Dr. Anna Ludig and Professor Jean Sibilia.

REFERENCES

1. Sekijima Y, Yoshida T, Ikeda S. CPPD crystal deposition disease of the cervical spine: a common cause of acute neck pain encountered in the neurology department. *J Neurol Sci* 2010;296:79-82.
2. Mahmud T, Basu D, Dyson P. Crystal arthropathy of the lumbar spine: a series of six cases and a review of the literature. *J Bone Joint Surg Br* 2005;87:513-7.
3. Derouet N, Haettich B, Temmar Z, Dugard D, Puechal X. [Septic arthritis of a lumbar facet joint. A case report]. [Article in French] *Ann Med Interne* 2001;152:279-82.
4. Fujishiro T, Nabeshima Y, Yasui S, Fujita I, Yoshiya S, Fujii H. Pseudogout attack of the lumbar facet joint: a case report. *Spine* 2002;27:E396-8.