

initiative that seeks to better understand the current state of people with axSpA. It uses an integrative approach based on scientific evidence, expert knowledge, and patient opinion. The Atlas population represents the entire spectrum of axSpA and is equally distributed in terms of sex. Additionally, the risk of mental disorders was assessed by using the 12-item General Health Questionnaire (GHQ-12). This is a validated screening tool for identifying minor psychiatric disorders in the general population within nonpsychiatric clinical settings such as primary care or general medical outpatient facilities. Therefore, the Atlas 2017 provides an opportunity to further examine the effect of disease outcomes on mental health in patients with axSpA. Moreover, the screening tool can shed light on the influence of explanatory variables related to mental disorders in these patients.

The objective of our present study was to determine the extent of risk of mental disorders in patients with axSpA and any associated factors. Our hypothesis was that greater disease activity is associated with a higher risk of having mental disorders.

MATERIALS AND METHODS

Study design and population. As part of the Atlas of Axial Spondyloarthritis in Spain project⁹, 680 patients diagnosed with axSpA were interviewed between May 1 and August 15, 2016. The Atlas seeks to promote early diagnosis, improved care, and the use of effective treatments in patients with axSpA. The anonymous questionnaire included a series of items related to disease status [disease activity as measured by the Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) self-administered questionnaire, spinal stiffness, and functional limitation] and psychological aspects related to coping with the disease. All the methodologies upon which the Atlas is based have been previously published in detail. This specific study includes data from 474 patients with recorded information about risk for mental disorders using the GHQ-12.

Disease activity. Disease activity was measured using the BASDAI with a range between 0 and 10¹⁰.

Spinal stiffness. In addition, spinal stiffness levels in the 3 areas of the vertebral column (cervical, dorsal, and lumbar) were collected, measured on a scale from lower to higher levels using an ordinal variable (1 = without limitation, 2 = mild limitation, 3 = moderate limitation, and 4 = severe limitation). Using this information, an index of spinal stiffness was created by adding, but without weighting, the degree of stiffness in each of the 3 areas. The resulting values ranged between 3 and 12. The Cronbach's alpha coefficient of this scale was 0.85, which attests to its consistency for assessing general stiffness.

Functional limitations in daily activities. To determine the degree of functional limitation, a composite index was used and included the sum of recorded limitation in 18 daily activities (dressing, grooming, bathing, tying shoelaces, moving about the home, walking up and down stairs, getting into/out of bed, using the toilet, shopping, preparing meals, eating, housework, walking, using public transportation, going to the doctor, driving, physical exercise, and sexual relations) using an ordinal variable (0 = none, 1 = little, 2 = some, and 3 = moderate). The resulting values oscillate between 0 and 54; thus, a value of 0–18 was set as a low limitation, 18–36 as a medium limitation, and 36–54 as an indication of high limitation. The Cronbach's alpha coefficient of this scale was 0.964, guaranteeing its reliability as a means for assessing limitation.

Sociodemographic aspects. The following demographic aspects were also collected: sex, age, living as a couple, education level, employment status, and whether the individual belonged to a patient association.

Comorbidity of mental health. Finally, patients were queried about any previous mental health diagnosis (made by psychologist or psychiatrist), including anxiety or depression. Such diagnoses of depression and/or anxiety referred to some point(s) in the past and patients were asked to provide a positive answer only when said diagnosis had been made by a psychologist/psychiatrist based on clinical criteria.

Risk of mental disorders. The risk of mental disorders was determined by using the GHQ-12¹¹, which consists of 12 items, each one assessing the severity of a mental health problem during the previous 4 weeks using a Likert scale (0–3). Thus, a maximum score of 36 was obtained; higher scores reflected an increased risk of psychiatric morbidity. However, to determine the risk of patients with poor mental health, these scores were transferred to a GHQ scale, where 0 or 1 = 0, and 2 or 3 = 1. Following this methodology, the majority of studies found that scores ≥ 3 indicated a high risk of poor mental health; therefore, we used this cutoff point for the present study¹². The GHQ-12 offers reliability in studies developed within the Spanish population, resulting in a Cronbach's alpha coefficient that oscillated between 0.76 and 0.90^{11,13}. In our study, a Cronbach's alpha coefficient of 0.934 was obtained, which supported the reliability hypothesis of this scale.

Statistical analysis. The statistical analysis was conducted in 3 stages. The first step focused on the descriptive statistics for the sample and the study variables. The second step consisted of bivariate regression analyses of each predictor regarding the risk of poor mental health (GHQ-12). Here, nonparametric contrasts were made because the GHQ-12 values obtained did not follow a normal distribution pattern. For quantitative variables, Mann-Whitney U tests for 2 categories, and Kruskal-Wallis tests for more than 2 categories were performed. For categorical variables, chi-square independence test or Fisher's exact test was used. In addition, the Spearman correlation coefficient was used to assess whether there was a relationship between mental health status (GHQ-12) and the different quantitative factors, such as the level of disease activity as measured by BASDAI.

The third step evaluated the independent influence of disease outcomes on mental health by using a hierarchical multiple regression analysis with those variables or disease outcomes that had shown a significant relationship with mental health status in the bivariate analyses conducted during the second step. For the hierarchical regression analysis, variables were grouped into 3 conceptual categories: (1) sociodemographic variables, (2) comorbidities, and (3) physical variables. This analysis introduced the variables of each category in a step-by-step fashion, thereby allowing for model comparison by building several regression models. Each model added variables to the previous one at each step; later models always included smaller models in their previous steps. In this case, our interest was to determine whether newly added conceptual categories of variables showed a significant improvement in R^2 (the proportion of explained variance in GHQ-12 by the model). Finally, a stepwise forward regression multivariable analysis was carried out, which included the variables that passed an F test, to ascertain the best associated factors.

All contrasts were bilateral and considered statistically significant when the p value was < 0.05 . The data were analyzed using SPSS software, version 24 for 64 bits (IBM Corp.).

RESULTS

Characteristics of the study participants. A total of 838 patients with axSpA anonymously accessed the online questionnaire between May 1 and August 15, 2016. After validation and normalization of the information, the sample consisted of 680 patients who responded to the majority of the questionnaire (the completion rate exceeded 75%) and made up the total valid sample of the Spanish Atlas of Axial Spondyloarthritis⁹. From these, a total of 474 patients had valid data for the GHQ-12 and comprised the group under evaluation for this study; characteristics of the sample are summarized in Table 1. The characteristics of this subgroup

Table 1. Sociodemographic data, disease outcomes, and comorbidity sample characteristics.

Variables	Values
Sociodemographic	
Age, yrs	45.4 ± 10.7
Sex, male	233 (49.1)
Having a partner, n = 444	386 (86.9)
Education level, university	185 (39.0)
Job status, employee	234 (49.4)
Patient association membership	227 (47.8)
Disease outcomes	
BASDAI (0–10), n = 442	5.4 ± 2.1
Spinal stiffness, moderate or severe	
Cervical (1–4), n = 447	201 (44.9)
Thoracic (1–4), n = 435	186 (42.7)
Lumbar (1–4), n = 458	288 (62.8)
Functional limitations in daily activities (0–54), n = 473	27.5 ± 12.7
Treatment	
NSAID	363 (76.6)
sDMARD	181 (38.2)
bDMARD	240 (50.3)
Comorbidity	
Self-reported depression diagnostic	99 (20.8)
Self-reported anxiety diagnostic	134 (28.2)
GHQ-12 score (0–36)	18.3 ± 8.0

N = 474 unless otherwise specified. Results are expressed as mean ± SD or n (%). BASDAI: Bath Ankylosing Spondylitis Disease Activity Index; NSAID: nonsteroidal antiinflammatory drug; sDMARD: synthetic disease-modifying antirheumatic drug; bDMARD: biological DMARD; GHQ-12: 12-item General Health Questionnaire.

were similar compared with the entire population, with the exception of more frequently having a relationship with a partner. In the studied population, the mean GHQ-12 score was 18.3 ± 8.0. Further, 310 (65.4%) patients were classified as having risk of mental disorders, while 164 (34.6%) had no risk of mental disorders.

Bivariate analyses. Table 2, Table 3, and Table 4 depict the results for the bivariate analyses. Mental health showed a statistically significant association with the following variables: older age, male sex, no relationship with a partner, unemployment, no patient association member-

Table 2. Bivariate analysis of sociodemographic factors associated with the risk of mental disorders measured by GHQ-12.

Sociodemographic Factors	Homogeneity Test: Kruskal-Wallis or Mann-Whitney	Chi-square
Age, yrs	0.001* (Spearman = -0.174)	≤ 0.001
Sex, male	0.013 [†]	0.025
Having a partner	0.009 [†]	0.021
Education level, university	0.607*	0.43
Job status, employee	≤ 0.001*	≤ 0.001
Patient association membership	0.001 [†]	0.001

* Kruskal-Wallis test. [†] Mann-Whitney test. GHQ-12: 12-item General Health Questionnaire.

Table 3. Bivariate analysis for comorbidity factors associated with risk of mental disorders measured by GHQ-12.

Comorbidity Factors	Homogeneity Test: Kruskal-Wallis or Mann-Whitney	Chi-square
Self-reported depression diagnostic	≤ 0.00 [†]	≤ 0.001
Self-reported anxiety diagnostic	≤ 0.001 [†]	≤ 0.001

[†] Mann-Whitney test. GHQ-12: 12-item General Health Questionnaire.

Table 4. Bivariate analysis for disease activity and functional limitation factors associated with risk of mental health disorders measured by GHQ-12.

Factors	Spearman Correlation Coefficient	p
Disease activity		
BASDAI (0–10)	0.543	≤ 0.001
Spinal stiffness, moderate or severe, n (%)		
Cervical (1–4)	0.120	0.011
Thoracic (1–4)	0.073	0.130
Lumbar (1–4)	0.165	≤ 0.001
Limitations in daily activities (0–54)	0.378	≤ 0.001

GHQ-12: 12-item General Health Questionnaire; BASDAI: Bath Ankylosing Spondylitis Disease Activity Index.

ship, anxiety and/or depression diagnosis, high level of disease activity (BASDAI), high degree of spinal stiffness (cervical and lumbar), and worse functional limitations in daily activities.

Hierarchical modeling with successive conceptual categories. To evaluate the independent influence of disease outcomes on mental health, a hierarchical multiple regression analysis with those variables that had shown a significant relationship with GHQ-12 in the bivariate analyses was performed (Table 5). This analysis demonstrated how, among sociodemographic variables, patient association membership had a significant regression coefficient while the rest did not.

In the first step, all sociodemographic attributes were included and established as control variables. That model explained patient mental health to a great extent ($R^2 = 83.2\%$). In the second step, diagnoses of depression and anxiety were added to this model but the R^2 value increased by only 0.6% ($p = 0.001$) and yielded no significant regression coefficients (p values = 0.08 and 0.8, respectively). In the third step, disease activity (BASDAI and stiffness) and function were also included in the model, which added 5.5% ($p < 0.001$) to the variance of the GHQ-12 scores. However, of these last variables only BASDAI and functional limitations experienced in daily life had a coefficient significantly different from zero: 0.52, $p < 0.001$ and 0.14, $p = 0.004$, respectively. This means that once the sociodemographic characteristics and psychiatric/psychological comorbidity (depression and anxiety) were established, a change in BASDAI levels or functional limitation reflects a change in the GHQ-12.

Table 5. Hierarchical multivariate analyses of sociodemographic, comorbidity, and disease status factors in relation to the risk of mental disorders, measured by GHQ-12 (n = 474).

Step	Factors	β -Weights	95% CI	Standard β -Weights	p	R ² (%), p+*	Δ R ² (%), p+**
1	Sociodemographic					0.832	0.832
	Age, yrs	-0.137	-1.143, 0.870	-0.016	0.789		
	Sex, male	0.424	-0.953, 1.800	0.033	0.545		
	Being in a relationship	0.606	-0.548, 1.760	0.042	0.302		
	Education level, university	0.663	-0.192, 1.518	0.107	0.128		
	Job status, employee	0.237	-0.083, 0.557	0.041	0.146		
	Patient association membership	2.113	0.770, 3.456	0.168	0.002		
2	Comorbidity					0.838	0.006
	Self-reported depression diagnostic	-1.705	-3.590, 0.181	-0.155	0.076		
	Self-reported anxiety diagnostic	-0.260	-2.020, 1.500	-0.023	0.772		
3	Disease outcomes					0.893	0.055
	BASDAI (0–10)	1.796	1.424, 2.167	0.527	< 0.001		
	Functional limitations in daily activities (0–54)	0.092	0.030, 0.154	0.140	0.004		
	Cervical stiffness (1–4)	0.392	-0.439, 1.224	0.051	0.354		
	Lumbar stiffness (1–4)	0.429	0.393, 1.250	0.063	0.306		

Only values that were statistically significant in the bivariate analysis (or marginally significant: $p < 0.09$) were included in this analysis. All β -weights, 95% CI, and p values for individual variables were estimates derived in the context of the full model (i.e., with all 3 conceptual blocks entered into the equation). * Overall R² (%) after adding each conceptual block and accompanying p value for the test of the overall R². ** Incremental R² change due to the addition of the conceptual block and accompanying p value for the test of the incremental R² change. BASDAI: Bath Ankylosing Spondylitis Disease Activity Index; GHQ-12: 12-item General Health Questionnaire.

Stepwise regression analysis without successive conceptual categories. Stepwise regression analysis resulted in the inclusion of 4 variables: BASDAI, functional limitations in daily activities, membership in a patient association, and cervical stiffness. All 4 of these variables contributed to a significant change in R² ($p < 0.001$ for BASDAI, belonging to a patient association, and functional limitations in daily activities, and $p = 0.03$ for cervical stiffness, where p is the p value for a change test in R²). Out of these variables, BASDAI was found to have remarkably more explanatory power (contribution to R² = 0.875; Table 6).

DISCUSSION

According to the data from the 2017 Atlas survey, 45.6% of patients with axSpA experience risk of mental disorders. Further, our study identifies several disease-specific factors associated with the risk of mental disorders. Among all factors, the most relevant is disease activity. Higher disease activity is associated with higher risk of mental disorders. In

addition, other factors associated with risk of mental disorders are worse functional limitation in daily life, not belonging to a patient association, and a higher degree of cervical stiffness. Moreover, 28% of patients self-reported having a clinical diagnosis of anxiety and 21% of depression. Although the prevalence of anxiety and depression varied depending on the definition used, the prevalence of these diseases in our study lies within the published ranges: 11–64% for depression⁸ and 16–20% for anxiety^{14,15,16}.

These findings are consistent with other evidence suggesting a relationship between disease activity and negative psychological factors associated with axSpA. This relationship has been proven to have a bidirectional character. On the one hand, it shows how negative psychological factors (depression, anxiety, passive coping style, insomnia, and negative affective states) explain the significant variability of disease activity when self-reported by patients, beyond what is explained by the clinical and demographic variables^{17,18,19,20}. On the other hand, it is evident how

Table 6. Stepwise forward multivariate analyses of sociodemographic and disease status variables in relation to GHQ-12 (n = 474).

Step	Predictors	β -Weights	CI (95%)	Standard β -Weights	p	R ² (%), p+*	Δ R ² (%), p+**
1	BASDAI (0–10)	1.912	1.551, 2.274	0.562	< 0.001	0.875	0.875
2	Patient association membership	2.028	1.012, 3.044	0.161	< 0.001	0.883	0.008
3	Functional limitations (0–54)	0.112	0.053, 0.171	0.170	< 0.001	0.888	0.005
4	Cervical stiffness (1–4)	0.610	0.054, 1.166	0.080	0.032	0.889	0.001

Only values that were statistically significant in the bivariate analysis (or marginally significant: $p < 0.09$) were included in the analysis. All β -weights, 95% CI, and p values for individual variables were estimates derived in the context of the full model (i.e., with all 3 conceptual blocks entered into the equation). * Overall R² (%) after adding each conceptual block and accompanying p value for the test of the overall R². ** Incremental R² change due to the addition of the conceptual block and accompanying p value for the test of the incremental R² change. BASDAI: Bath Ankylosing Spondylitis Disease Activity Index; GHQ-12: 12-item General Health Questionnaire.

disease activity and functional limitation can explain some variability in patient mental health status (anxiety, depression, and insomnia)^{7,16,21,22}.

The high prevalence of risk of mental disorders found in our study is discouraging. Nevertheless, one must take into account that the identified factors associated with risk of mental disorders, especially the most influential (disease activity), are not fixed and can be modified with a more appropriate and intensive treatment strategy. Indeed, these results should encourage physicians to develop strategies for improving this situation during the coming years.

In the field of chronic disease, of great importance is an adequate assessment of the information given by patients in addition to their quality of life, because a patient's perspective can guide resource allocation, the design of interventions, and pharmacological treatment^{23,24,25}. The results of our study add to the evidence supporting the relationship between disease status (disease activity, spinal stiffness), psychosocial factors (functional limitation in daily activities), and the mental health status of patients with axSpA. However, it is important to understand the mechanism underlying this type of relationship before considering how to use this knowledge in clinical practice. These findings also raise the question of whether a psychological intervention, perhaps directed at certain subgroups of patients, may be useful in the treatment of axSpA. In this context, there is some evidence as to the influence of certain psychological interventions on these pathologies, although they are more commonly observed in patients who have rheumatoid arthritis than those with axSpA^{26,27,28,29}.

This study has several limitations. First, the use of previously nonvalidated scales or indices for assessing certain factors, such as functional limitations in daily life and stiffness, should be thoroughly considered. The reason for using such scales or composite indices originated during the preliminary phase of the survey development, when patients expressed their concern about not being able to report all aspects of their disease if other scales or indices were to be used. In any case, a good Cronbach's alpha coefficient was obtained for the index used in our study, which testifies to the reliability of these instruments. Second, the bidirectional features of these relationships are another limitation of this study, because it is not possible to establish causality using this cross-sectional approach. In fact, it is difficult to assess whether the risk of mental disorders stems from the problems related to the disease, the underlying inflammatory processes, or other factors associated with this chronic pathology (spinal stiffness or functional limitations in daily life). Conclusions can only be drawn based on the relationship between variables. To establish a causality, it would be necessary to carry out longitudinal studies to evaluate over time the evolution of physical variables and their relationship to mental health or vice versa. Third, a possible overlap between GHQ-12 and the diagnosis of depression and anxiety cannot be completely excluded. However, the diagnosis of depres-

sion and anxiety in these respondents was made by a psychologist/psychiatrist based on clinical criteria, while GHQ-12 is a self-reported measure of the risk of mental disorders. The GHQ-12 values show the current risk of mental disorders during the survey period while the diagnoses of depression and anxiety referred to points in the past. Finally, the possible effect of biological therapy on depressive symptoms and mental health-associated quality of life could have influenced the results of our study.

The results of our study underscore the high prevalence of mental disorders in patients with axSpA and its clear association with disease status. About 1 of every 2 patients with axSpA reports risk of mental disorders. Further, this risk of mental disorders seems to be explained to a great extent by the degree of disease activity and to a lesser extent to such factors as the degree of functional limitation in daily life, cervical stiffness, and patient association membership. These findings highlight the benefit of rheumatologists promoting psychiatric evaluations of patients with high disease activity and risk of mental disorders. It is posited that this will contribute to a more integral treatment strategy.

ACKNOWLEDGMENT

We thank the Spanish Society of Rheumatology for editing the manuscript.

REFERENCES

1. Kiltz U, van der Heijde D. Health-related quality of life in patients with rheumatoid arthritis and in patients with ankylosing spondylitis. *Clin Exp Rheumatol* 2009;4 Suppl 55:S108-11.
2. Lim HJ, Lim HS, Lee MS. Relationship between self-efficacy and exercise duration in patients with ankylosing spondylitis. *Clin Rheumatol* 2005;24:442-3.
3. Penhoat M, Saroux A, Le Goff B, Augereau P, Maugars Y, Berthelot JM. High pain catastrophizing scores in one-fourth of patients on biotherapy for spondylarthritis or rheumatoid arthritis. *Joint Bone Spine* 2014;81:235-9.
4. Yagiz AE, Kokacaya MH, Copoglu US, Uruc V, Paksoy H, Yengil E, et al. The effect of social support and severity of the disease on posttraumatic growth in ankylosing spondylitis. *Acta Medica Mediterr* 2014;30:1355-9.
5. DiMatteo MR, Lepper HS, Croghan TW. Depression is a risk factor for noncompliance with medical treatment: meta-analysis of the effects of anxiety and depression on patient adherence. *Arch Intern Med* 2000;160:2101-7.
6. Jang JH, Green CE, Assassi S, Reveille JD, Ward MM, Weisman MH, et al. The contribution of disease activity on functional limitations over time through psychological mediators: a 12-month longitudinal study in patients with ankylosing spondylitis. *Rheumatology* 2011;50:2087-92.
7. Kilic G, Kilic E, Ozgocmen S. Relationship between psychiatric status, self-reported outcome measures, and clinical parameters in axial spondyloarthritis. *Medicine* 2014;93:e337.
8. Zhao S, Thong D, Miller N, Duffield SJ, Hughes DM, Chadwick L, et al. The prevalence of depression in axial spondyloarthritis and its association with disease activity: a systematic review and meta-analysis. *Arthritis Res Ther* 2018;20:140.
9. Garrido-Cumbrera M, Navarro-Compán V, Zarco P, Collantes-Estévez E, Gálvez-Ruiz D, Braçe O, et al. Atlas of Axial Spondyloarthritis in Spain 2017: study design and population. [Article in English, Spanish] *Reumatol Clin* 2019;15:127-32.

10. Garrett S, Jenkinson T, Kennedy LG, Whitelock H, Gaisford P, Calin A. A new approach to defining disease status in ankylosing spondylitis: the Bath Ankylosing Spondylitis Disease Activity Index. *J Rheumatol* 1994;21:2286-91.
11. Golderberg D, Williams P. A user's guide to the General Health Questionnaire. Windsor, UK: NFER-Nelson; 1988.
12. Bones Rocha K, Pérez K, Rodríguez-Sanz M, Borrell C, Obiols JE. [Prevalence of mental health problems and their association with socioeconomic, work and health variables: Findings from the Spain National Health Survey]. [Article in Spanish] *Psicothema* 2010;22:389-95.
13. Sánchez-López M del P, Dresch V. The 12-item General Health Questionnaire (GHQ-12): reliability, external validity and factor structure in the Spanish population. *Psicothema* 2008;20:839-43.
14. Chan CYY, Tsang HHL, Lau CS, Chung HY. Prevalence of depressive and anxiety disorders and validation of the Hospital Anxiety and Depression Scale as a screening tool in axial spondyloarthritis patients. *Int J Rheum Dis* 2017;20:317-25.
15. Xu X, Shen B, Zhang A, Liu J, Da Z, Liu H, et al. Anxiety and depression correlate with disease and quality-of-life parameters in Chinese patients with ankylosing spondylitis. *Patient Prefer Adherence* 2016;10:879-85.
16. Baysal O, Durmuş B, Ersoy Y, Altay Z, Senel K, Nas K, et al. Relationship between psychological status and disease activity and quality of life in ankylosing spondylitis. *Rheumatol Int* 2011;31:795-800.
17. Brionez TF, Assassi S, Reveille JD, Learch TJ, Diekman L, Ward MM, et al. Psychological correlates of self-reported functional limitation in patients with ankylosing spondylitis. *Arthritis Res Ther* 2009;11:R182.
18. Brionez TF, Assassi S, Reveille JD, Green C, Learch T, Diekman L, et al. Psychological correlates of self-reported disease activity in ankylosing spondylitis. *J Rheumatol* 2010;37:829-34.
19. Domínguez CJ, Ugalde PF, Vilchez DR, Carretero-Dios H, Estévez EC. Positive and negative affective states and disease activity in ankylosing spondylitis. *Rheumatol Int* 2015;35:519-24.
20. Jiang Y, Yang M, Wu H, Song H, Zhan F, Liu S, et al. The relationship between disease activity measured by the BASDAI and psychological status, stressful life events, and sleep quality in ankylosing spondylitis. *Clin Rheumatol* 2015;34:503-10.
21. Bagcivan G, Cinar FI, Cinar M, Oflaz F, Uzun S, Pay S. Living with pain in ankylosing spondylitis: a qualitative study. *Contemp Nurse* 2015;51:135-47.
22. Shen CC, Hu LY, Yang AC, Kuo BI, Chiang YY, Tsai SJ. Risk of psychiatric disorders following ankylosing spondylitis: a nationwide population-based retrospective cohort study. *J Rheumatol* 2016;43:625-31.
23. van Genderen S, Plasqui G, van der Heijde D, van Gaalen F, Heuft L, Luime J, et al. Social role participation and satisfaction with life: a study among patients with ankylosing spondylitis and population controls. *Arthritis Care Res* 2018;70:600-7.
24. Montacer Kchir M, Mehdi Ghannouchi M, Hamdi W, Azzouz D, Kochbati S, Saadellaoui K, et al. Impact of the ankylosing spondylitis on the professional activity. *Joint Bone Spine* 2009;76:378-82.
25. Betegnig AL, Gauchet A, Lehmann A, Grange L, Roustit M, Baudrant M, et al. Why do patients with chronic inflammatory rheumatic diseases discontinue their biologics? An assessment of patients' adherence using a self-report questionnaire. *J Rheumatol* 2016;43:724-30.
26. Salaffi F, Carotti M, Gasparini S, Intorcchia M, Grassi W. The health-related quality of life in rheumatoid arthritis, ankylosing spondylitis, and psoriatic arthritis: a comparison with a selected sample of healthy people. *Health Qual Life Outcomes* 2009;7:25.
27. Haywood KL, Garratt AM, Jordan KP, Healey EL, Packham JC. Evaluation of ankylosing spondylitis quality of life (EASI-QoL): reliability and validity of a new patient-reported outcome measure. *J Rheumatol* 2010;37:2100-09.
28. Rau J, Ehlebracht-König I, Petermann F. [Impact of a motivational intervention on coping with chronic pain: results of a controlled efficacy study]. [Article in German] *Schmerz* 2008;22:575-8.
29. Geenen R, Newman S, Bossema ER, Vriezেকolk JE, Boelen PA. Psychological interventions for patients with rheumatic diseases and anxiety or depression. *Best Pract Res Clin Rheumatol* 2012; 26:305-19.