

TITLE PAGE

Title:

Biomechanical properties of common carotid arteries assessed by circumferential two-dimensional strain and beta stiffness index in patients with ankylosing spondylitis

Manuscript type:

Full-length Manuscripts

Authors

Helena Forsblad-d'Elia ORCID iD 0000-0001-7871-5303, Lucy Law ORCID iD 0000-0002-8129-8771, Karin Bengtsson ORCID iD 0000-0001-5154-0144, Johan Smeds, Maria Ketonen, Björn Sundström ORCID iD 0000-0001-7226-0969, Lotta Ljung ORCID iD 0000-0001-8999-0925, Mats Geijer ORCID iD 0000-0003-0529-7723, Stefan Söderberg ORCID iD 0000-0001-9225-1306, Per Lindqvist ORCID iD 0000-0002-8192-9166

Key Indexing Terms

Ankylosing spondylitis, cardiovascular disease, common carotid artery, ultrasound

Departments/Institutions

Department of Public Health and Clinical Medicine, Rheumatology, Umeå University, Umeå, Sweden, Department of Rheumatology and Inflammation Research, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden; Region Västra Götaland, Sahlgrenska University Hospital,

This article has been accepted for publication in The Journal of Rheumatology following full peer review. This version has not gone through proper copyediting, proofreading and typesetting, and therefore will not be identical to the final published version. Reprints and permissions are not available for this version. Please cite this article as doi: 10.3899/jrheum.200146. This accepted article is protected by copyright. All rights reserved.

Department of Rheumatology, Gothenburg, Sweden, Department of Surgical and Perioperative Sciences, Clinical Physiology, Umeå University, Umeå, Sweden, Department of Radiology, Institute of Clinical Sciences, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden; Region Västra Götaland, Sahlgrenska University Hospital, Department of Radiology, Gothenburg, Sweden and Faculty of Medicine, Lund University, Lund, Sweden, Department of Public Health and Clinical Medicine, Medicine, Umeå University, Umeå, Sweden

The sources of support

This study was supported by grants from The Swedish Research Council, Västerbotten's Association Against Rheumatism, The Swedish Association Against Rheumatism, the County of Västerbotten (agreement concerning research and education of doctors), King Gustaf Vth 80-year Foundation, The Norrland's Heart Foundation, and Mats Kleberg's Foundation

Conflict of interest

None of the authors report any potential conflict of interest.

Authors, academic degrees and appointments

H. Forsblad-d'Elia, MD, Professor, Department of Public Health and Clinical Medicine, Rheumatology, Umeå University, Umeå, Sweden

L. Law, BSc, Department of Public Health and Clinical Medicine, Rheumatology, Umeå University, Umeå, Sweden

K. Bengtsson, MD, Department of Rheumatology and Inflammation Research, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg,

Sweden; Region Västra Götaland, Sahlgrenska University Hospital, Department of Rheumatology, Gothenburg, Sweden

J. Smeds, MD, Department of Public Health and Clinical Medicine, Rheumatology, Umeå University, Umeå, Sweden

M. Ketonen, MD, Department of Surgical and Perioperative Sciences, Clinical Physiology, Umeå University, Umeå, Sweden

B. Sundström, RPT, Associate Professor, Department of Public Health and Clinical Medicine, Rheumatology, Umeå University, Umeå, Sweden

L. Ljung, MD, Associate Professor, Department of Public Health and Clinical Medicine, Rheumatology, Umeå University, Umeå, Sweden

M. Geijer, MD, Professor, Department of Radiology, Institute of Clinical Sciences, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden; Region Västra Götaland, Sahlgrenska University Hospital, Department of Radiology, Gothenburg, Sweden and Faculty of Medicine, Lund University, Lund, Sweden

S. Söderberg, MD, Professor, Department of Public Health and Clinical Medicine, Medicine, Umeå University, SE-901 87 Umeå, Sweden

P. Lindqvist, BSc, Professor, Department of Surgical and Perioperative Sciences, Clinical Physiology, Umeå University, Umeå, Sweden

Corresponding author:

Helena Forsblad-d'Elia, Department of Public Health and Clinical Medicine, Rheumatology, Umeå University, SE-901 87 Umeå, Sweden

E-mail: helena.forsblad.delia@umu.se

Phone number: +46 761103883

ORCID iD 0000-0001-7871-5303

Short running head

Biomechanics of carotid arteries

ABSTRACT

Objective

Ankylosing spondylitis (AS) is associated with an elevated risk of cardiovascular disease (CVD) related to atherosclerosis, preceded by arterial stiffness. We aimed to examine common carotid artery (CCA) biomechanical properties using ultrasound to calculate β stiffness index (indicating arterial stiffness) and, a more recently developed technique, two-dimensional (2D) speckle tracking strain (indicating arterial motion and deformation, strain) to 1) compare with age- and sex-matched controls and to 2) analyze relationships between strain and stiffness with disease characteristics and traditional risk factors for CVD in AS patients.

Methods

In this cross-sectional study, a cohort of 149 patients with AS, mean age 55.3 ± 11.2 years, 102 (68.5%) men, 146 (98%) HLA-B27 positive, were examined. Bilateral CCAs were examined for circumferential 2D strain and β stiffness index. A subgroup of 46 patients were compared with 46 age- and sex-matched controls, both groups without hypertensive disease, diabetes, myocardial infarction or stroke.

Results

Mean bilateral circumferential 2D strain was lower in AS patients compared with controls, $7.9 \pm 2.6\%$ vs $10.3 \pm 1.9\%$, $p < 0.001$ whereas mean bilateral β stiffness index was higher, $13.1 \pm 1.6 \text{ mmHg/mm}$ vs $12.3 \pm 1.3 \text{ mmHg/mm}$, $p = 0.018$. In multivariable linear regression analyses strain was associated with age, erythrocyte sedimentation

rate, history of anterior uveitis and treatment with csDMARD and/or bDMARD (R^2 0.33), while stiffness was associated with age (R^2 0.19).

Conclusion

Both CCA circumferential 2D strain and β stiffness index differed between AS patients and controls. Strain was associated with AS-related factors and age while stiffness with age, suggesting that the obtained results reflect different pathogenic vascular processes.

INTRODUCTION

Ankylosing spondylitis (AS), a subgroup of the spondyloarthritis (SpA) diseases, is a chronic rheumatic inflammatory disease primarily affecting the sacroiliac joints and spine but also, to a lesser extent, peripheral joints (1). It has been demonstrated that patients with AS have an increased risk of cardiovascular disease (CVD) compared to the general population (2-8). The European League Against Rheumatism has recognized CVD as an important comorbidity in patients with inflammatory joint disorders including SpA and emphasizes the need for risk assessment and risk management (9). Several clinically important manifestations of CVD are related to atherosclerosis such as coronary artery disease, stroke, and peripheral arterial disease. It is suggested that chronic systemic inflammation, in addition to traditional cardiovascular (CV) risk factors, contributes to the atherosclerotic process (10). The atherosclerotic process is characterized by the degeneration of smooth muscle cells and elastin fibers in parallel with the proliferation of more rigid collagen fibers in the vessel walls and intra- and extracellular deposition of lipids. These changes might lead to the development of increased arterial stiffness and also to the increase of the intima-media thickness (IMT) and plaque formation which can be evaluated by

ultrasound of the common carotid arteries (CCAs) (11). However, limitations of measuring carotid (c)IMT have been recognized since the association between cIMT progression and CV risk in the general population has remained unproven (12). Therefore, the development of better methods to evaluate atherosclerotic changes is warranted. Biomechanical properties of the arteries, such as stiffness, can also be evaluated by ultrasound where the β stiffness index can be calculated from the relation between systemic blood pressure and arterial diameter. The β stiffness index has been found to be significantly associated with coronary atherosclerosis (13). Furthermore, the β stiffness index correlated with the carotid atherosclerotic grade, vessel wall area and wall thickness suggesting that the β stiffness index of the CCA reflects not only biomechanical properties of the artery but also its atherosclerotic damage (14). However, the ultrasound method used for β stiffness index calculation is angle-dependent and only measures mechanics in one dimension. Technological advancements in ultrasound have resulted in a method assessing two-dimensional (2D) strain, using a speckle tracking technique, which measures vascular motion and deformation biomechanics in two dimensions. Speckle tracking was originally developed for examining the myocardium, providing additional information to conventional cardiac ultrasound methods (15). Recently, the speckle tracking based 2D strain technique has been applied in vascular studies with the aim of improving the understanding of the atherosclerotic process and to detect early subclinical disease (16, 17). Previous studies demonstrated that 2D speckle tracking strain correlated with cIMT (18-20), and that strain, in contrast to cIMT, was associated with the severity and extent of coronary artery disease (20).

The primary aim of this cross-sectional study was to investigate, for the first time in patients with AS, the biomechanical properties of the CCAs with both circumferential

2D strain and β stiffness index and to compare the results with age- and sex-matched controls. A secondary aim was to explore relationships between circumferential 2D strain and β stiffness index with AS disease characteristics and traditional risk factors for CVD in order to estimate the explanatory value of these factors for the biomechanical properties of the CCAs.

MATERIALS and METHODS

Patients and Controls

All patients attending the rheumatology clinic in Region Västerbotten in northern Sweden with a diagnosis of AS (ICD-10 M45.9) between May 2002 and November 2015 were identified through the digital administrative system (n=523). The diagnosis of AS was validated through a review of the medical records and patients not fulfilling the modified New York criteria (21) were excluded, leaving 346 patients. Two-hundred forty-six patients between 18 and 75 years of age, still living in Region Västerbotten, with at least one visit at the rheumatology clinic within the last five years were invited between 2016 and 2017 to take part in a study called the Backbone study. A flow-chart over the inclusion process is shown in Figure 1. Exclusion criteria were dementia, other inflammatory rheumatic diseases, pregnancy, or difficulties in understanding the Swedish language. One-hundred and fifty-five (63%) patients, fulfilling the criteria, were willing to participate in the Backbone study, investigating severity and comorbidities in AS. For the current study, six patients were further excluded due to a lack of or inadequate imaging data required for speckle tracking 2D circumferential strain analysis, leaving 149 patients. Out of the 246 patients, the 149 patients included in this report had a similar median age 55.0 (45.5, 62.5) years compared with the 97 patients not taking part, 52.0 (38.5, 63.0) years, $p=0.070$ (December 31, 2015). There was a sex difference between the included patients,

102/149 (68.4%) men compared to those not taking part, 81/97 (83.5%) men, $p=0.008$.

The patients with AS underwent clinical examinations and answered questionnaires regarding lifestyle habits, medication, AS-related data such as a history of anterior uveitis, peripheral arthritis and CV-related factors such as previous myocardial infarction, surgical myocardial revascularization or stroke. Patients having been told by a physician to have hypertension and being on an antihypertensive drug were defined as having hypertensive disease. Patients having been told by a physician to have diabetes and being on an antidiabetic drug were defined as having diabetes mellitus. The Bath Ankylosing Spondylitis Activity Index (BASDAI), Ankylosing Spondylitis Disease Activity Score with C-reactive protein (ASDAS-CRP), Bath Ankylosing Spondylitis Functional Index (BASFI) and Bath Ankylosing Spondylitis Metrology Index (BASMI) were assessed (22). The patients answered a questionnaire regarding health-related quality of life (HRQoL), Short-Form 36 (SF-36) (23, 24) and we report herein the overall physical component summary (PCS) score and the mental component summary (MCS) score. Blood samples were drawn in the morning after an overnight fast and erythrocyte sedimentation rate (ESR), high-sensitivity C-reactive protein (hs-CRP) and lipids were analyzed by standard laboratory techniques, consecutively.

From the 149 patients in this report, a subset of 46 patients (31 men, 15 women) without diabetes, hypertensive disease and without a history of myocardial infarction, surgical myocardial revascularization or stroke were selected consecutively from the list of inclusion and compared with 46 age- and sex-matched controls recruited from the hospital staff. The same inclusion criteria were applied for the controls, besides not having any inflammatory rheumatic disease.

Radiography

Spinal radiographic changes were assessed from the lateral projection of the spinal radiographs and were graded using the modified Stoke Ankylosing Spondylitis Spinal Score (mSASSS). The anterior corners of vertebra C2-T1 and T12-S1 were graded with a score between 0 and 3 (0 = normal, 1 = erosion, sclerosis or squaring, 2 = syndesmophyte, 3 = bridging syndesmophyte). The overall scoring scale ranges from 0 to 72, with 72 representing complete ankylosis (25). To have an mSASSS score ≥ 2 at a vertebral corner was classified as having a syndesmophyte. Severe spinal radiographic changes were defined as ≥ 3 consecutive inter-vertebral bridges in the cervical spine and/or the lumbar spine, similar to the definition of grade 4=severe in the Bath Ankylosing Spondylitis Radiology Index (BASRI) (26). The x-rays were performed at a mean time of 32.9 ± 15.1 days after inclusion in the study. One experienced radiologist performed all scoring (MG).

Ultrasound examination

The same operator (LL) carried out bilateral CCA ultrasound examinations on all patients and controls. A GE Vivid E9 ultrasound system with a GE 9L 2.5-8 MHz linear transducer (GE, Boston, Massachusetts) was used. All participants were examined in a supine position, resting quietly with their head tilted at a 45-degree angle away from the side being assessed. Blood pressure (BP) was taken using the right upper arm and a manual sphygmomanometer after a 5 minutes' rest in a supine position. A superimposed ECG was used to identify end-systole (end of T wave) and end-diastole (Q-wave). Standard B-mode short-axis (SAX) and long-axis (LAX) views of the right and left CCA were obtained. Image optimization was performed as

appropriate for each examination. CCA images included the carotid bulb as a reference. CCA measurements were taken 1-2 cm into the proximal CCA from the bulb. A 5 beat loop of CCA from the short-axis view was stored for further analysis. All examinations were stored in the Digital Imaging and Communications in Medicine (DICOM) format. As previously described the ultrasound examinations were post-processed and analyzed using TomTec (27, 28). We used TomTec Arena™ version 4.0 (TomTec Imaging Systems GMBH, Germany) and the post-processing was performed by the same operator (LL).

Speckle tracking strain

The mid-left ventricular SAX circumferential strain option (based on the speckle tracking ultrasound method for left ventricular assessment) was used to measure CCA 2D circumferential strain parameters from SAX (specific equation used not provided by the software company). The clip was edited to exclude significant drift or movement. An average of 3 consecutive beats was analyzed. The internal vessel wall was outlined manually at end-systole and end-diastole, and the clip played to ensure accurate speckle tracking analysis. The average circumferential 2D strain value was then recorded (Figure 2a and b). A higher circumferential 2D strain value indicates more motion and deformation of the vessel wall.

β stiffness index

β stiffness index was calculated offline using the equation: $\beta = \ln(\text{SBP}/\text{DBP}) / ((\text{ESD} - \text{EDD})/\text{EDD})$ (29). The systolic blood pressure (SBP) and diastolic blood pressure (DBP) were taken at the time of the examination in a supine position, and the end-systolic diameter (ESD) and end-diastolic diameter (EDD) luminal diameters were taken from the CCA LAX 1-2 cm into the CCA from the bulb from 3 consecutive heartbeats. ESD and EDD were defined as the largest and smallest luminal diameters,

respectively (Figure 2c). A higher β stiffness index indicates an increased stiffness of the vessel.

Reliability Testing of Circumferential 2D Strain

An expert ultrasound operator (PL) analyzed the left CCA of 10 randomly selected individuals, blinded to whether the individual was a patient or a control person. The delineation of the wall for tracking was done independently by the two operators (LL and PL) and were compared by inter-observer reliability testing and the calculated coefficient of variation was 11.7%.

Ethics

The Regional Ethical Review Board at Umeå University, Sweden approved the study, (patients dnr 2015/352-31, 2016/208-31, controls dnr 2010-21-21, 2014/198-32M) which was performed in accordance with the Declaration of Helsinki. All patients included in the Backbone study gave written informed consent.

Consent for publication

Consent for publication has been obtained (Figure 2).

Statistics

Continuous variables are presented as mean (SD) or median (25th percentile (Q1), 75th percentile (Q3)) and categorical variables are shown as numbers and percentages. An independent *t*-test or the Mann-Whitney U test was used to compare continuous variables as appropriate and the Chi-square test was used for categorical comparisons. Correlations between variables were calculated using Pearson's bivariate correlation test. Univariable and multivariable linear regression analyses were used to analyze

factors associated with mean bilateral CCA circumferential 2D strain and mean bilateral β stiffness index. The dependent variables, mean bilateral CCA circumferential 2D strain and mean bilateral β stiffness index, were normally distributed. Independent variables with a univariable p -value ≤ 0.1 were considered for the multivariable models. Also, correlations between independent variables in the models were analyzed and the limit was set to $r < 0.7$ and the variable with best prediction in the univariable analysis was selected for the multivariable analysis. Residual plots were assessed for assumptions of linearity to be confirmed. To have a characteristic was coded 1 and to not have a characteristic was coded 0 in dichotomous variables. Female sex was coded 1 and male sex 0. Statistics were performed using SPSS version 24 (SPSS Inc., IBM, Chicago, USA). $P < 0.05$ was considered statistically significant.

RESULTS

Altogether 149 patients (68.5 % men) were included with a mean age of 55.3 ± 11.2 years and a mean symptom duration of 31.5 ± 11.6 years. HLA-B27 was present in 146 (98.0%) patients. Sixty-seven (45.0%) of the patients were ever smokers and 8 (5.4%) smoked regularly. The median mSASSS value was 8.0 (1.0, 30.0) and 82 (55.0%) AS patients had at least one syndesmophyte. In total, 36 (24.2%) patients were treated with a conventional synthetic disease-modifying anti-rheumatic drug (csDMARD) and/or a biologic (b)DMARD. Sixty-five (43.6%) of the patients had hypertensive disease and 21 (14.1%) were on medication against dyslipidemia, Table 1.

Comparisons between patients with ankylosing spondylitis and controls

The patients with AS had significantly lower CCA strain and higher stiffness identified by a lower circumferential 2D strain and higher β stiffness index compared to the controls, Table 2. Neither significant difference in circumferential 2D strain was found between men and women with AS (8.3 ± 2.5 , % vs 7.0 ± 2.8 , %, $p=0.12$) nor in β stiffness index between men and women with AS (13.1 ± 1.3 , mmHg/mm vs 13.2 ± 2.2 , mmHg/mm, $p=0.95$). Concerning controls, neither significant difference in circumferential 2D strain was found between men and women (10.2 ± 1.9 % vs 10.4 ± 2.1 %, $p=0.73$) nor in β stiffness index between control men and women (12.5 ± 1.45 , mmHg/mm vs 12.15 ± 1.15 , mmHg/mm, $p=0.39$). Three (6.5 %) of the patients with AS smoked regularly and 23 (50.0 %) had ever been smokers. No information about smoking was available for controls.

Linear regression analyses demonstrating factors associated with mean bilateral common carotid artery circumferential 2D strain, all AS patients

In the univariable analysis, CCA circumferential 2D strain associated significantly with age, symptom duration, ESR, hsCRP, history of anterior uveitis or peripheral arthritis, BASMI, severe spinal radiographic changes, SBP, DBP, and heart rate, Table 3. In the multivariable analysis, the mean circumferential 2D strain showed inverse significant associations with age, ESR, a history of anterior uveitis, and present treatment with a csDMARD and/or a bDMARD (R^2 0.33), Table 4.

Linear regression analyses demonstrating factors associated with mean bilateral common carotid artery β stiffness index, all AS patients

In the univariable analysis, the mean CCA β stiffness index associated significantly with age, symptom duration, BASMI, mSASSS, severe spinal radiographic changes,

and hypertensive disease, Table 3. In the multivariable analyses, only age associated with the mean β stiffness index (R^2 0.19), Table 4.

DISCUSSION

In this investigation of biomechanical properties by ultrasound of the CCAs in a contemporary cohort of patients with AS from northern Sweden, we demonstrated a reduced strain and increased stiffness in patients with AS compared with controls. We selected two methods, the recently developed speckle tracking circumferential 2D strain and the established method, β stiffness index, since we were interested in studying, for the first time in patients with AS, biomechanics on the same arteries with different methods and find out if they were comparable or not. Speckle tracking circumferential 2D strain assesses arterial motion and deformation while the β stiffness index assesses arterial stiffness. Furthermore, in the multivariable analyses among patients with AS, we found that AS-related factors and age associated with circumferential 2D strain, and only age associated with β stiffness index. Thus, age was the only common contributing determinant explaining some of the variations of the two measurements of biomechanical properties of CCAs in this cohort of patients with AS. Interestingly, the AS-related variables ESR, history of anterior uveitis, and present treatment with a csDMARD and/or a bDMARD were also significant determinants of circumferential 2D strain. Thus, our results indicate that the circumferential 2D strain method has the capacity to capture aspects of strain related to inflammation and the severity of the AS disease. The HLA-B27 positive rate is high, 98% in this cohort of patients with AS from northern Sweden which may be explained by the high rate in the population, 17% in this area (30).

There is a growing recognition that the prevalence of CVD is increased in patients

with AS, which contributes to increased mortality (7, 31-34). An elevated risk of cardiovascular and cerebrovascular diseases related to atherosclerosis has been demonstrated in patients with AS (2, 4-6, 35, 36). In addition, the prevalence of other typical AS-related cardiac manifestations such as aortic insufficiency and cardiac conduction disturbances is more common compared to the general population (3, 5, 37). Together with traditional CV risk factors for atherosclerotic CVD (38), inflammation itself is considered to play a role in AS (39). Decreased elasticity of the arterial wall may be present before the occurrence of clinical symptoms or atherosclerotic plaques. Biomechanical properties of the CCAs have only been investigated in a few studies on patients with inflammatory arthritis diseases, all being cross-sectional. Kaplanoglu et al. recently reported no difference in β stiffness index between 38 patients with AS (mean age 39.6 years) and 49 healthy controls (mean age 35.5 years) (40). The discrepancy with our results might be explained by the lower mean patient age in the study by Kaplanoglu et al. compared to our study. In other investigations assessing vessel biomechanics, pulse-wave velocity (PWV) used to assess aortic stiffness, augmentation index (AIx) measuring arterial stiffness, and echocardiographic evaluation of aortic distensibility were impaired in AS patients compared with controls (41-43), the results being in line with our findings. Moreover, in a study on patients with rheumatoid arthritis both β stiffness index and speckle tracking 2D strain showed results in the same direction as ours; the β stiffness index was increased and the strain reduced compared to controls (44). Likewise, in patients with psoriasis, out of which about 20% had psoriatic arthritis, β stiffness index was increased compared to controls (45).

We did not find a significant association between strain and stiffness in all AS patients ($r=-0.13$, $p=0.11$, data not shown). Such an association has been displayed in

persons without inflammatory rheumatic disease (19, 46). This discrepancy might possibly be explained by our findings that AS-related factors were associated with biomechanical properties measured by circumferential 2D strain but not with the β stiffness index. Concerning anterior uveitis, we have previously discovered it to be independently associated with aortic regurgitation in AS, believed to be induced by an inflammatory process in the aortic root (47). Interestingly, we now also show anterior uveitis to be related to the strain of the CCAs. Inflammation is known to accelerate atherosclerosis and in a longitudinal study on AS, CRP and ASDAS were associated with future elevated AIx (48). However, it remains to establish if inflammation is also related to the forthcoming impairment of the biomechanical properties of CCAs in AS.

Among risk factors for CVD, we found age to be associated with circumferential 2D strain and β stiffness index in the multivariable analyses. Kaplanoglu et al. showed, in univariate analyses, that the β stiffness index was associated with age, symptom duration, and BMI in AS, partly in agreement with our findings (40).

There are some limitations to the current study. It is cross-sectional; thus, we cannot draw any conclusions about causality. The R^2 values in the multivariable models are rather low meaning that other unknown factors contribute to explaining the variation of circumferential 2D strain and β stiffness index. The number of AS patients was somewhat limited (n=149) and also the number of controls. Mostly Caucasian men and women were included and the results cannot be generalized to other ethnicities. Furthermore, the proportion of men was lower compared to the non-participants. However, we do not think that the difference has influenced the results considerably since there were no significant differences in circumferential 2D strain (men 8.1 ± 2.8 % vs women 7.5 ± 3.0 %, $p=0.18$, data not shown) or β stiffness index (men 13.4 ± 1.4

mmHg/mm vs women 13.2 ± 1.8 mmHg/mm, $p=0.5$, data not shown) between the examined men and women with AS. Additionally, we did not have data on smoking habits, BMI and dyslipidemia in the controls, recruited from hospital staff, which is a major limitation. However, the number of smokers who smoked regularly among the AS patients was low. The well-characterized cohort of patients with AS with matched controls and the usage of appropriate ultrasound methods are some of the notable strengths of this study.

In conclusion, the circumferential 2D strain was reduced and the β stiffness index increased in patients with AS compared to matched controls indicating impaired biomechanical properties of CCAs in AS patients. Strain was associated with factors related to AS-disease severity, which was not observed for stiffness. This could imply that the process leading to impaired strain is more dependent on the course of the AS disease than is the development of stiffness. Larger and longitudinal studies are required to investigate the clinical importance of these markers of arteriosclerosis and subclinical atherosclerosis in AS and if they add predictive value besides already identified risk factors for CVD.

Availability of data and material

The data sets generated and/or analyzed during the current study are not publicly available due to the General Data Protection Regulation (GDPR), but a limited and fully anonymized data set that supports the main analyses is available from the corresponding author on request.

Author's contributions

HFdE. was responsible for study design, recruitment of patients, data collection, statistical analyses, interpretation of data and writing the manuscript.

L.L. assessed ultrasound examinations, participated in the analysis, interpretation of data and drafting the manuscript.

K.B. participated in the analysis, interpretation of data and drafting the manuscript.

J.S. performed clinical examinations, participated in the planning of study and data collection and interpretation of data.

M.K. performed clinical examinations, participated in the planning of the study and data collection and interpretation of data.

B.S. contributed to the validation of participants, participated in the planning of the study and interpretation of data.

L.Lj. contributed to the validation of participants, participated in the planning of the study and interpretation of data.

M.G. assessed the radiographic scoring, participated in the interpretation of data and drafting the manuscript.

S.S. participated in interpretation of data and drafting the manuscript.

P.L. assessed ultrasound examinations, participated in the interpretation of data and drafting the manuscript.

All authors have critically reviewed the manuscript, approved the final version to be published and agreed to be accountable for all aspects of the work.

ACKNOWLEDGMENT

We wish to thank all the patients who participated in the study. We also wish to thank the research nurses at the University Hospital of Umeå, Viktoria von Zweigbergk and Jeanette Beckman Rehnman, for assisting with the project.

REFERENCES:

1. Braun J, Sieper J. Ankylosing spondylitis. *Lancet* 2007;369:1379-90.
2. Bengtsson K, Forsblad-d'Elia H, Lie E, Klingberg E, Dehlin M, Exarchou S, et al. Are ankylosing spondylitis, psoriatic arthritis and undifferentiated spondyloarthritis associated with an increased risk of cardiovascular events? A prospective nationwide population-based cohort study. *Arthritis Res Ther* 2017;19:102.
3. Bengtsson K, Forsblad-d'Elia H, Lie E, Klingberg E, Dehlin M, Exarchou S, et al. Risk of cardiac rhythm disturbances and aortic regurgitation in different spondyloarthritis subtypes in comparison with general population: A register-based study from Sweden. *Ann Rheum Dis* 2018;77:541-8.
4. Bremander A, Petersson IF, Bergman S, Englund M. Population-based estimates of common comorbidities and cardiovascular disease in ankylosing spondylitis. *Arthritis Care Res (Hoboken)* 2011;63:550-6.
5. Szabo SM, Levy AR, Rao SR, Kirbach SE, Lacaille D, Cifaldi M, et al. Increased risk of cardiovascular and cerebrovascular diseases in individuals with ankylosing spondylitis: A population-based study. *Arthritis Rheum* 2011;63:3294-304.
6. Chou CH, Lin MC, Peng CL, Wu YC, Sung FC, Kao CH, et al. A nationwide population-based retrospective cohort study: Increased risk of acute coronary syndrome in patients with ankylosing spondylitis. *Scand J Rheumatol* 2014;43:132-6.

7. Haroon NN, Paterson JM, Li P, Inman RD, Haroon N. Patients with ankylosing spondylitis have increased cardiovascular and cerebrovascular mortality: A population-based study. *Ann Intern Med* 2015;163:409-16.
8. Han C, Robinson DW, Jr., Hackett MV, Paramore LC, Fraeman KH, Bala MV. Cardiovascular disease and risk factors in patients with rheumatoid arthritis, psoriatic arthritis, and ankylosing spondylitis. *J Rheumatol* 2006;33:2167-72.
9. Agca R, Heslinga SC, Rollefstad S, Heslinga M, McInnes IB, Peters MJ, et al. EULAR recommendations for cardiovascular disease risk management in patients with rheumatoid arthritis and other forms of inflammatory joint disorders: 2015/2016 update. *Ann Rheum Dis* 2017;76:17-28.
10. Libby P, Ridker PM, Hansson GK, Leducq Transatlantic Network on A. Inflammation in atherosclerosis: From pathophysiology to practice. *J Am Coll Cardiol* 2009;54:2129-38.
11. Patel AK, Suri HS, Singh J, Kumar D, Shafique S, Nicolaides A, et al. A review on atherosclerotic biology, wall stiffness, physics of elasticity, and its ultrasound-based measurement. *Curr Atheroscler Rep* 2016;18:83.
12. Lorenz MW, Polak JF, Kavousi M, Mathiesen EB, Volzke H, Tuomainen TP, et al. Carotid intima-media thickness progression to predict cardiovascular events in the general population (the prog-imt collaborative project): A meta-analysis of individual participant data. *Lancet* 2012;379:2053-62.
13. Hirai T, Sasayama S, Kawasaki T, Yagi S. Stiffness of systemic arteries in patients with myocardial infarction. A noninvasive method to predict severity of coronary atherosclerosis. *Circulation* 1989;80:78-86.

- Accepted Article
14. Wada T, Kodaira K, Fujishiro K, Maie K, Tsukiyama E, Fukumoto T, et al. Correlation of ultrasound-measured common carotid artery stiffness with pathological findings. *Arterioscler Thromb* 1994;14:479-82.
 15. Bansal M, Kasliwal RR. How do i do it? Speckle-tracking echocardiography. *Indian Heart J* 2013;65:117-23.
 16. Bjallmark A, Lind B, Peolsson M, Shahgaldi K, Brodin LA, Nowak J. Ultrasonographic strain imaging is superior to conventional non-invasive measures of vascular stiffness in the detection of age-dependent differences in the mechanical properties of the common carotid artery. *Eur J Echocardiogr* 2010;11:630-6.
 17. Oishi Y, Miyoshi H, Iuchi A, Nagase N, Ara N, Oki T. Vascular aging of common carotid artery and abdominal aorta in clinically normal individuals and preclinical patients with cardiovascular risk factors: Diagnostic value of two-dimensional speckle-tracking echocardiography. *Heart Vessels* 2013;28:222-8.
 18. Iino H, Okano T, Daimon M, Sasaki K, Chigira M, Nakao T, et al. Usefulness of carotid arterial strain values for evaluating the arteriosclerosis. *J Atheroscler Thromb* 2019;26:476-87.
 19. Catalano M, Lamberti-Castronuovo A, Catalano A, Filocamo D, Zimbalatti C. Two-dimensional speckle-tracking strain imaging in the assessment of mechanical properties of carotid arteries: Feasibility and comparison with conventional markers of subclinical atherosclerosis. *Eur J Echocardiogr* 2011;12:528-35.
 20. Kim SA, Park SM, Kim MN, Kim YH, Cho DH, Ahn CM, et al. The relationship between mechanical properties of carotid artery and coronary artery disease. *Eur Heart J Cardiovasc Imaging* 2012;13:568-73.

21. van der Linden S, Valkenburg HA, Cats A. Evaluation of diagnostic criteria for ankylosing spondylitis. A proposal for modification of the New York criteria. *Arthritis Rheum* 1984;27:361-8.
22. Sieper J, Rudwaleit M, Baraliakos X, Brandt J, Braun J, Burgos-Vargas R, et al. The assessment of spondyloarthritis international society (ASAS) handbook: A guide to assess spondyloarthritis. *Ann Rheum Dis* 2009;68 Suppl 2:ii1-44.
23. Ware JE, Jr., Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Medical Care* 1992;30:473-83.
24. Sullivan M, Karlsson J, Ware JE, Jr. The swedish sf-36 health survey--i. Evaluation of data quality, scaling assumptions, reliability and construct validity across general populations in Sweden. *Soc Sci Med* 1995;41:1349-58.
25. Creemers MC, Franssen MJ, van't Hof MA, Gribnau FW, van de Putte LB, van Riel PL. Assessment of outcome in ankylosing spondylitis: An extended radiographic scoring system. *Ann Rheum Dis* 2005;64:127-9.
26. MacKay K, Mack C, Brophy S, Calin A. The Bath ankylosing spondylitis radiology index (BASRI): A new, validated approach to disease assessment. *Arthritis Rheum* 1998;41:2263-70.
27. Yang EY, Dokainish H, Virani SS, Misra A, Pritchett AM, Lakkis N, et al. Segmental analysis of carotid arterial strain using speckle-tracking. *J Am Soc Echocardiogr* 2011;24:1276-84 e5.
28. Yang EY, Brunner G, Dokainish H, Hartley CJ, Taffet G, Lakkis N, et al. Application of speckle-tracking in the evaluation of carotid artery function in subjects with hypertension and diabetes. *J Am Soc Echocardiogr* 2013;26:901-9 e1.

29. Kawasaki T, Sasayama S, Yagi S, Asakawa T, Hirai T. Non-invasive assessment of the age related changes in stiffness of major branches of the human arteries. *Cardiovasc Res* 1987;21:678-87.
30. Bjelle A, Cedergren B, Dahlqvist SR. HLA B27 in the population of northern Sweden. *Scand J Rheumatol* 1982;11:23-6.
31. Nurmohamed MT, van der Horst-Bruinsma I, Maksymowych WP. Cardiovascular and cerebrovascular diseases in ankylosing spondylitis: Current insights. *Current Rheumatology Reports* 2012;14:415-21.
32. Exarchou S, Lie E, Lindstrom U, Askling J, Forsblad-d'Elia H, Turesson C, et al. Mortality in ankylosing spondylitis: Results from a nationwide population-based study. *Ann Rheum Dis* 2015.
33. Bakland G, Gran JT, Nossent JC. Increased mortality in ankylosing spondylitis is related to disease activity. *Ann Rheum Dis* 2011;70:1921-5.
34. Mok CC, Kwok CL, Ho LY, Chan PT, Yip SF. Life expectancy, standardized mortality ratios, and causes of death in six rheumatic diseases in Hong Kong, China. *Arthritis Rheum* 2011;63:1182-9.
35. Peters MJ, Visman I, Nielen MM, Van Dillen N, Verheij RA, van der Horst-Bruinsma IE, et al. Ankylosing spondylitis: A risk factor for myocardial infarction? *Ann Rheum Dis* 2010;69:579-81.
36. Keller JJ, Hsu JL, Lin SM, Chou CC, Wang LH, Wang J, et al. Increased risk of stroke among patients with ankylosing spondylitis: A population-based matched-cohort study. *Rheumatol Int* 2014;34:255-63.
37. Ward MM. Lifetime risks of valvular heart disease and pacemaker use in patients with ankylosing spondylitis [abstract]. *Journal of the American Heart Association* 2018;7:e010016.

38. Greenland P, Knoll MD, Stamler J, Neaton JD, Dyer AR, Garside DB, et al. Major risk factors as antecedents of fatal and nonfatal coronary heart disease events. *JAMA* 2003;290:891-7.
39. Hahn BH, Grossman J, Chen W, McMahon M. The pathogenesis of atherosclerosis in autoimmune rheumatic diseases: Roles of inflammation and dyslipidemia. *J Autoimmun* 2007;28:69-75.
40. Kaplanoglu H, Ozisler C. Evaluation of subclinical atherosclerosis using ultrasound radiofrequency data technology in patients diagnosed with ankylosing spondylitis. *J Ultrasound Med* 2019;38:703-11.
41. Bodnar N, Kerekes G, Seres I, Paragh G, Kappelmayer J, Nemethne ZG, et al. Assessment of subclinical vascular disease associated with ankylosing spondylitis. *J Rheumatol* 2011;38:723-9.
42. Moyssakis I, Gialafos E, Vassiliou VA, Boki K, Votteas V, Sfrikakis PP, et al. Myocardial performance and aortic elasticity are impaired in patients with ankylosing spondylitis. *Scand J Rheumatol* 2009;38:216-21.
43. Berg IJ, van der Heijde D, Dagfinrud H, Seljeflot I, Olsen IC, Kvien TK, et al. Disease activity in ankylosing spondylitis and associations to markers of vascular pathology and traditional cardiovascular disease risk factors: A cross-sectional study. *J Rheumatol* 2015;42:645-53.
44. Lee JH, Cho KI, Kim SM. Carotid arterial stiffness in patients with rheumatoid arthritis assessed by speckle tracking strain imaging: Its association with carotid atherosclerosis. *Clin Exp Rheumatol* 2012;30:720-8.
45. Kim SY, Yang HS, Lee YW, Choe YB, Ahn KJ. Evaluation of the beta stiffness index and carotid intima-media thickness in Asian patients with psoriasis. *Angiology* 2015;66:889-95.

46. Saito M, Okayama H, Inoue K, Yoshii T, Hiasa G, Sumimoto T, et al. Carotid arterial circumferential strain by two-dimensional speckle tracking: A novel parameter of arterial elasticity. *Hypertens Res* 2012;35:897-902.
47. Klingberg E, Svealv BG, Tang MS, Bech-Hanssen O, Forsblad-d'Elia H, Bergfeldt L. Aortic regurgitation is common in ankylosing spondylitis - time for routine echocardiography evaluation? *Am J Med* 2015.
48. Berg IJ, Semb AG, van der Heijde D, Kvien TK, Olsen IC, Dagfinrud H, et al. CRP and ASDAS are associated with future elevated arterial stiffness, a risk marker of cardiovascular disease, in patients with ankylosing spondylitis: Results after 5-year follow-up. *Ann Rheum Dis* 2015;74:1562-6.

FIGURE LEGENDS

Figure 1


Title: Flow-chart over the inclusion of patients with ankylosing spondylitis into the Backbone study

Legend: AS; ankylosing spondylitis

Figure 2

Title: Figures of ultrasound examinations of the common carotid artery. a) Outlining of the internal vessel wall at end-diastole. b) Speckle tracking strain output. C) Measurement of end-systolic diameter.

Legend: CCA; common carotid artery, ED; end-diastole, ES; end-systole, ESD; end-systolic diameter






Table 1. Descriptive characteristics of 149 patients with ankylosing spondylitis.

General characteristics	
Sex, women	47 (31.5)
men	102 (68.5)
Age, years	55.3 ± 11.2
BMI, kg/m ²	27.8 ± 5.2
Ever smoker	67 (45.0)
Current smoker, regular	8 (5.4)
SF-36, PCS [^] , score	39.8 ± 8.8
SF-36, MCS [^] , score	45.2 ± 11.6
AS-related variables	
Duration of symptoms, years	31.5 ± 11.6
HLA-B27 positive	146 (98.0)
ESR, mm/h	10.0 (5, 20)
	13.7 ± 11.9
CRP, mg/L	2.6 (1.0, 6.0)
	4.6 ± 6.1
History of anterior uveitis	77 (51.7)
History of peripheral arthritis	80 (53.7)
BASDAI, score	3.7 ± 1.9
ASDAS-CRP, score	1.8 ± 0.7
BASFI, score	2.9 ± 2.0
BASMI, score	4.1 ± 1.5
NSAID, regular use	94 (63.1)
csDMARD	19 (12.8)
bDMARD	25 (16.8)
csDMARD and/or bDMARD	36 (24.2)
mSASSS#, score	8.0 (1.0, 30.0)
	17.5 ± 20.4
≥ 1 syndesmophyte#	82 (55.0)
Severe spinal radiographic changes# □	30 (20.3)
Comorbidity and cardiovascular-related variables	
Systolic BP, mmHg	133 ± 17
Diastolic BP, mmHg	77 ± 10
Pulse pressure, mmHg	56 ± 13

Heart rate, bpm	68 ± 11
Right CCA circumferential strain, %	7.9± 3.1
Left CCA circumferential strain, %	7.9± 3.3
Mean CCA circumferential strain, %	7.9 ± 2.9
Right CCA β stiffness, mmHg/mm ^	13.3± 2.0
Left CCA β stiffness, mmHg/mm ^	13.4± 2.0
Mean CCA β stiffness, mmHg/mm ^	13.3 ± 1.5
MI, surgical myocardial revascularization or stroke	9 (6)
Hypertensive disease	65 (43.6)
Diabetes mellitus	8 (5.4)
Dyslipidemia, medication	21 (14.1)
Cholesterol, mmol/L	5.4 ± 1.1
HDL, mmol/L	1.6 ± 0.5
LDL#, mmol/L `	3.3 ± 0.9
Cholesterol/HDL	3.7 ± 1.2
Triglycerides, mmol/L	1.3 ± 0.6

Values are mean ±SD, median (Q1, Q3) or numbers of patients and percent (%). Number of missing data: #=1 missing, ^= 3 missing, ≥3 consecutive inter-vertebral bridges, cervical and/or lumbar spine.

BMI; body mass index, SF-36; Short Form-36, PCS; physical component summary, MCS; mental component summary, ESR; erythrocyte sedimentation rate; CRP; C-reactive protein, BASDAI; Bath Ankylosing Disease Activity Index, ASDAS; Ankylosing Spondylitis Disease Activity Score, BASFI; Bath Ankylosing Spondylitis Functional Index, BASMI; Bath Ankylosing Spondylitis Metrology Index, NSAID; non-steroidal anti-inflammatory drug, csDMARD; conventional synthetic disease modifying anti-rheumatic drug, b; biologic, mSASSS; Modified Stroke Ankylosing Spondylitis Score, BP; blood pressure, bpm; beats per minute, CCA; common carotid artery, MI; myocardial infarction, HDL; High Density Lipoprotein, LDL; Low Density Lipoprotein

Table 2. Comparisons between circumferential two-dimensional strain and β stiffness index in common carotid arteries in patients with ankylosing spondylitis and age- and sex-matched controls.

	AS patients (n = 46)	Controls (n = 46)	P-value
Sex, men	31 (64.7)	31 (64.7)	
women	15 (32.6)	15 (32.6)	
Age, years	50.4 \pm 8.7	49.8 \pm 9.2	0.75
Systolic BP, mmHg [^]	127 \pm 13	127 \pm 12	0.96
Diastolic BP, mmHg [^]	75 \pm 9	74 \pm 8	0.53
Circumferential 2D strain, right, %	7.8 \pm 2.8	10.1 \pm 2.1	<0.001
Circumferential 2D strain, left, %	8.0 \pm 3.1	10.5 \pm 2.6	<0.001
Mean circumferential 2D strain, %	7.9 \pm 2.7	10.3 \pm 1.9	<0.001
β stiffness index, right, mmHg/mm [^]	13.0 \pm 1.7	12.4 \pm 1.4	0.05
β stiffness index, left, mmHg/mm [^]	13.3 \pm 2.1	12.3 \pm 1.8	0.02
Mean β stiffness index, mmHg/mm [^]	13.1 \pm 1.7	12.3 \pm 1.3	0.02

Values are mean \pm SD or numbers of patients and percent (%)

BP; blood pressure, 2D; two-dimensional, [^]; Systolic and Diastolic BP; controls, n= 45, β stiffness index, right; AS patients and controls, n=44, 45 respectively, β stiffness index, left; AS patients and controls, n=45, 44 respectively, mean β stiffness index; AS patients and controls, n=43, 44 respectively

Table 3. Univariable linear regression analysis in 149 patients with ankylosing spondylitis with common carotid artery biomechanical measurements as dependent variables.

	Mean bilateral circumferential 2D strain, %		Mean bilateral β stiffness index, mmHg/mm	
General characteristics	B, unstandardized	P-value	B, unstandardized	P-value
Sex, women (vs men)	-0.67	0.18	-0.19	0.48
Age, years	-0.073	<0.001	0.052	<0.001
BMI, kg/m ²	-0.046	0.30	0.017	0.52
Ever smoker	-0.30	0.53	-0.18	0.65
SF-36, PCS, score	0.022	0.46	-0.011	0.44
SF-36, MCS, score	-0.005	0.81	-0.004	0.71
Covariates, AS-related				
Duration of symptoms, years	-0.063	0.002	0.029	0.008
HLAB-27 positive	-1.63	0.33	1.60	0.14
ESR, mm/h	-0.084	<0.001	0.005	0.68
CRP, mg/L	-0.092	0.016	0.005	0.80
History of anterior uveitis	-1.18	0.012	-0.023	0.93
History of peripheral arthritis	-1.13	0.016	0.30	0.23
BASDAI, score	-0.039	0.75	0.020	0.77
ASDAS-CRP, score	-0.17	0.61	0.060	0.75
BASFI, score	-0.19	0.098	0.10	0.11
BASMI, score	-0.49	0.001	0.32	<0.001
NSAID, regular use	0.44	0.37	0.056	0.83
csDMARD and/or bDMARD	-1.07	0.051	0.41	0.18
mSASSS, score	-0.016	0.18	0.015	0.016
≥ 1 syndesmophyte	-0.15	0.76	0.43	0.090
Severe spinal radiographic changes [□]	-1.16	0.047	0.64	0.044
Covariates, CV-related				
Systolic BP, mmHg	-0.037	0.006	0.009	0.20
Diastolic BP, mmHg	-0.070	0.003	0.00	0.98

Pulse pressure, mmHg	-0.023	0.19	0.015	0.10
Heart rate, bpm	-0.055	0.008	0.020	0.075
MI, surgical myocardial revascularization or stroke	1.15	0.24	-0.46	0.39
Hypertensive disease	-0.48	0.31	0.53	0.038
Diabetes mellitus	0.46	0.66	0.25	0.65
Dyslipidemia, medication	0.064	0.92	0.42	0.24
Cholesterol, mmol/L	-0.41	0.055	0.12	0.32
HDL, mmol/L	-0.91	0.063	0.11	0.70
LDL, mmol/L	-0.36	0.15	0.077	0.57
Triglycerides, mmol/L	-0.12	0.75	0.32	0.13

≥ 3 consecutive inter-vertebral bridges, cervical and/or lumbar spine. 2D; two-dimensional, CI; confidence interval, BMI; body mass index, Short Form-36, PCS; physical component summary, MCS; mental component summary, ESR; erythrocyte sedimentation rate; CRP; C-reactive protein, BASDAI; Bath Ankylosing Disease Activity Index, ASDAS; Ankylosing Spondylitis Disease Activity Score, BASFI; Bath Ankylosing Spondylitis Functional Index, BASMI; Bath Ankylosing Spondylitis Metrology Index, NSAID; non-steroidal anti-inflammatory drug, csDMARD; conventional synthetic disease modifying anti-rheumatic drug, b; biologic, mSASSS; Modified Stroke Ankylosing Spondylitis Score, BP; blood pressure, bpm; beats per minute, MI; myocardial infarction, HDL; High Density Lipoprotein, LDL; Low Density Lipoprotein. To have a characteristic was coded 1 and to not have a was coded 0. Female sex was coded 1 and male sex 0.

Table 4. Multivariable linear regression analysis in 149 patients with ankylosing spondylitis with mean bilateral common carotid artery a) circumferential two-dimensional strain and b) β stiffness index as dependant variables.

a)	Mean bilateral circumferential 2D strain, %		
Covariates	B, unstandardized	B, standardized	P-value
Age, years	-0.054	-0.21	0.023
ESR, mm/h	-0.070	-0.29	<0.001
Anterior uveitis	-0.95	-0.17	0.025
Peripheral arthritis ever	-0.71	-0.12	0.097
BASFI, score	0.036	0.026	0.76
BASMI, score	0.057	0.031	0.78
csDMARD and/or bDMARD	-1.06	-0.16	0.035
Severe spinal radiographic changes \square	-0.78	-0.11	0.22
Diastolic BP, mmHg	-0.035	-0.12	0.099
Heart rate, bpm	-0.038	-0.15	0.056
Cholesterol, mmol/L	-0.070	-0.027	0.72
HDL, mmol/L	-0.58	-0.096	0.21
	R² 0.33		
b)	Mean bilateral β stiffness index, mmHg/mm		
Covariates	B, unstandardized	B, standardized	P-value
Age, years	0.043	0.32	0.003
BASMI, score	0.13	0.13	0.27
Syndesmophyte	-0.23	-0.077	0.43
Severe spinal radiographic changes \square	0.12	0.034	0.74
Pulse pressure, mmHg	-0.002	-0.022	0.80
Heart rate, bpm	0.018	0.14	0.096
Hypertensive disease	0.17	0.057	0.50
	R² 0.19		

\square = ≥ 3 consecutive inter-vertebral bridges, cervical and/or lumbar spine.

2D; two-dimensional, ESR; erythrocyte sedimentation rate, BASFI; Bath Ankylosing Spondylitis Functional Index, BASMI; Bath Ankylosing Spondylitis Metrology Index, csDMARD; conventional synthetic disease modifying anti-rheumatic drug, b; biologic, BP; blood pressure, bpm; beats per minute, HDL; High Density Lipoprotein, BASMI; Bath Ankylosing Spondylitis Metrology Index. To have a characteristic was coded 1 and to not have a was coded 0.