

Ultrasound, Skin, and Joints in Psoriatic Arthritis

EMILIO FILIPPUCCI, ROSSELLA DE ANGELIS, FAUSTO SALAFFI, and WALTER GRASSI

ABSTRACT. Over the last decade, ultrasound has been increasingly used in rheumatology for assessing soft tissue involvement in patients with chronic arthritis. In spite of the high number of studies supporting the role and the validity of ultrasound in the assessment of patients with rheumatoid arthritis, the potential of ultrasound imaging in patients with psoriatic arthritis still waits to be adequately investigated. Our report illustrates the most relevant sonographic findings obtainable with the latest generation ultrasound equipment in patients with psoriatic arthritis. (J Rheumatol 2009;36 Suppl 83:35-38; doi:10.3899/jrheum.090220)

Key Indexing Terms:
ULTRASOUND
SKIN

PSORIATIC ARTHRITIS
JOINT

Over the last decade, ultrasound (US) has been increasingly used in rheumatology for assessing soft tissue involvement in patients with chronic arthritis¹⁻⁸. In spite of the high number of studies supporting the role and the validity of US in the assessment of patients with rheumatoid arthritis, the potential of US imaging in patients with psoriatic arthritis (PsA) still waits to be adequately investigated²⁻⁶. Our report illustrates the most relevant US findings obtainable with the latest generation US equipment in patients with PsA.

METHODS

US imaging of patients with PsA requires dedicated equipment and proper scanning technique. Very high frequency probes (> 13 MHz for B mode and > 8 MHz for Doppler mode) and an adequate amount of gel are needed to obtain the best visualization of superficial soft tissues. This is particularly true for the US examination of skin and nails.

Dynamic, multiplanar US examination of joints and tendons is mandatory for revealing the main US findings detectable in patients with PsA, including synovial effusion, synovial proliferation, tendon tears, enthesitis, and bone erosions. The US images presented here were obtained using a MyLab70 XVG system (Esaote Biomedica, Genoa, Italy) equipped with 6–18 MHz and 4–13 MHz broadband linear probes.

RESULTS

Skin. US appearance of normal skin consists of a bilaminar layer: the epidermis is visualized as a sharp hyper-echoic line and the underlying dermis appears as a thicker and less echogenic band. Epidermis and dermis are

clearly distinguishable due to their different echotexture. More difficult is the identification of the acoustic interface separating the dermis from the subcutaneous fat^{9,10}. US findings of psoriatic plaque include inhomogeneous thickening of the epidermis with or without acoustic shadow and hypoechoic swelling of the dermis with or without power Doppler signal (Figure 1)⁹.

Nails. On US examination, the normal nail plate appears as a trilaminar structure, characterized by 2 hyperechoic sharp margins with an interposed thin anechoic line. US features indicative of nail plate involvement in psoriatic onychopathy include the loss of the sharpness of the hyper-echoic lines, which may appear focally curved and/or thickened, and the loss of the intermediate anechoic layer, which may be focal or complete. The nail bed involvement is characterized by a variable degree of thickening associated with or without power Doppler signal.

Joints. Joint effusion and synovial proliferation can easily be distinguished by grey scale US imaging even at the distal interphalangeal joint¹¹. Dynamic assessment while compressing the soft tissues under examination with the probe is the most reliable method to distinguish synovial fluid and synovial tissue, the former being more easily compressible and displaceable¹². Power Doppler helps in the identification of inflamed synovial tissue and has been demonstrated to be of value in therapy monitoring of PsA patients¹³⁻¹⁶ (Figure 2, A–A'). For the detection of sacroiliitis, contrast enhanced Doppler US has shown high negative predictive value in the observation of sacroiliac joints of patients with inflammatory low back pain¹⁷.

Tendons. Tenosynovitis is a relatively common finding in patients with PsA, the hand and ankle being the most frequently involved anatomic regions. When examining tenosynovitis of the finger flexor tendons, particular attention must be paid to not misinterpreting finger tendon pulleys as signs of synovial proliferation. At ankle level, posterior tibialis and peroneal tendons represent common sites of tenosynovitis even in asymptomatic PsA

From the Clinica Reumatologica, Università Politecnica delle Marche, Ancona, Italy.

E. Filippucci, MD, Senior Lecturer; R. De Angelis, MD, Senior Lecturer; F. Salaffi, MD, Associate Professor; W. Grassi, MD, Full Professor.

Address correspondence to Dr. W. Grassi, Cattedra di Reumatologia, Università Politecnica delle Marche, Ospedale A. Murri, Via dei Colli 52, 60035 Jesi, Ancona, Italy. E-mail: walter.grassi@univpm.it

Personal non-commercial use only. The Journal of Rheumatology Copyright © 2009. All rights reserved.




Figure 1. Psoriasis. A. Psoriatic plaque at the anterior aspect of the leg. Black line indicates the exact point where the probe was placed. B. Grey scale imaging showing hypoechoic swelling of the dermis. B'. Power Doppler imaging showing increased perfusion of the dermis.




Figure 2. Psoriatic arthritis. A–A': proliferative synovitis. Longitudinal (A) and transverse (A') views on the dorsal aspect of the metacarpophalangeal joint showing marked intraarticular power Doppler signal. B–B': dactylitis. B. Proliferative tenosynovitis. Longitudinal view on the volar aspect of the proximal interphalangeal joint showing an evident tendon sheath widening. B'. Power Doppler imaging revealing increased perfusion of the synovial tissue surrounding the flexor tendons (t). mp: middle phalanx; pp: proximal phalanx; mc: metacarpal bone.




Figure 3. Psoriatic arthritis. Enthesitis of the Achilles tendon (t). A. Proliferative retrocalcaneal bursitis. Asterisk indicates an area of bursal effusion. B. Arrows indicate enthesophytes. C. Arrowhead indicates a calcaneal erosion. c: calcaneal bone.

patients¹⁸. In dactylitis or “sausage-like” digits, US imaging allows the identification of different patterns of soft tissue involvement generated by the variable association of flexor tenosynovitis, synovitis of interphalangeal and metacarpophalangeal joints, and surrounding soft tissue edema¹⁹⁻²¹ (Figure 2, B–B’). Achilles tendon, plantar fascia, and patellar tendon are the most frequently affected tendons without tendon sheath. US findings indicative of tendon

pathology include tendon fusiform swelling and focal derangement of tendon echotexture with or without intratendinous power Doppler signal.

Enthesis. At lower limb entheses, US has been shown to be a sensitive tool for revealing subclinical enthesal involvement^{22,23}. US allows detailed assessment of both tendinous and bony side involvement²⁵⁻³². At the Achilles

tendon insertion into the calcaneal bone, early US signs of enthesitis include hypoechoic swelling of the tendon insertion, increase of blood flow detectable with power Doppler, and retrocalcaneal bursal enlargement²⁴ (Figure 3). Although most studies have been conducted at the entheses of the lower limbs, there is also evidence of US ability to evaluate enthesitis of the upper limbs²⁵. In established disease, US changes at the bony level become more evident and include erosions and enthesophytes.

CONCLUSION

Grey scale and power Doppler US imaging with very high frequency probes allows a detailed assessment of a wide range of morphostructural and perfusional changes at skin, enthesis, tendon, and joint level. Several US patterns result from the variable association of these findings.

The data obtained in patients with rheumatoid arthritis are expected to be reproduced also in those with PsA. Studies investigating validity issues, including accuracy and reproducibility, are required to define the role of US in the assessment of psoriasis.

REFERENCES

- Kane D, Grassi W, Sturrock R, Balint PV. Musculoskeletal ultrasound – state of the art review in rheumatology. Part 2: Clinical indications for musculoskeletal ultrasound in rheumatology. *Rheumatology* 2004;43:829-38.
- Filippucci E, Iagnocco A, Meenagh G, et al. Ultrasound imaging for the rheumatologist. *Clin Exp Rheumatol* 2006;24:1-5.
- Meenagh G, Filippucci E, Kane D, Taggart A, Grassi W. Ultrasonography in rheumatology: developing its potential in clinical practice and research. *Rheumatology* 2007;46:3-5.
- Joshua F, Lassere M, Bruyn GA, et al. Summary findings of a systematic review of the ultrasound assessment of synovitis. *J Rheumatol* 2007;34:839-47.
- Wakefield RJ, D'Agostino MA, Iagnocco A, et al. The OMERACT Ultrasound Group: status of current activities and research directions. *J Rheumatol* 2007;34:848-51.
- Terslev L, von der Recke P, Torp Pedersen S, Koenig MJ, Bliddal H. Diagnostic sensitivity and specificity of Doppler ultrasound in rheumatoid arthritis. *J Rheumatol* 2008;35:49-53.
- Riente L, Delle Sedie A, Filippucci E, et al. Ultrasound imaging for the rheumatologist IX. Ultrasound imaging in spondyloarthritis. *Clin Exp Rheumatol* 2007;25:349-53.
- Filippucci E, Iagnocco A, Meenagh G, et al. Ultrasound imaging for the rheumatologist VII. Ultrasound imaging in rheumatoid arthritis. *Clin Exp Rheumatol* 2007;25:5-10.
- El Gammal S, El Gammal C, Kaspar K. Sonography of the skin at 100 MHz enables in vivo visualization of stratum corneum and viable epidermis in palmar skin and psoriatic plaques. *J Invest Dermatol* 1999;113:821-9.
- Hesselstrand R, Scheja A, Wildt M, Akesson A. High frequency ultrasound of skin involvement in systemic sclerosis reflects oedema, extension and severity in early disease. *Rheumatology* 2008;47:84-7.
- Grassi W, Filippucci E, Farina A, Cervini C. Sonographic imaging of the distal phalanx. *Semin Arthritis Rheum* 2000;29:379-84.
- Wakefield RJ, Balint PV, Szkudlarek M, et al. OMERACT 7 Special Interest Group. Musculoskeletal ultrasound including definitions for ultrasonographic pathology. *J Rheumatol* 2005;32:2485-7.
- Fiocco U, Cozzi L, Chieco Bianchi F, et al. Vascular changes in psoriatic knee synovitis. *J Rheumatol* 2001;28:2480-6.
- Fiocco U, Ferro F, Cozzi L, et al. Contrast medium in power Doppler ultrasound for assessment of synovial vascularity: comparison with arthroscopy. *J Rheumatol* 2003;30:2170-6.
- Unlu E, Pamuk On, Cakir N. Color and duplex Doppler sonography to detect sacroiliitis and spinal inflammation in ankylosing spondylitis. Can this method reveal response to anti-tumor necrosis factor therapy? *J Rheumatol* 2007;34:110-6.
- Fiocco U, Ferro F, Vezzu M, et al. Rheumatoid and psoriatic knee synovitis: clinical, grey scale, and power Doppler ultrasound assessment of the response to etanercept. *Ann Rheum Dis* 2005;64:899-905.
- Klauser A, Halpern EJ, Frauscher F, et al. Inflammatory low back pain: high negative predictive value of contrast-enhanced color Doppler ultrasound in the detection of inflamed sacroiliac joints. *Arthritis Rheum* 2005;53:440-4.
- Galluzzo E, Lischi Dm, Taglione E, et al. Sonographic analysis of the ankle in patients with psoriatic arthritis. *Scand J Rheumatol* 2000;29:52-5.
- Olivieri I, Scarano E, Padula A, Giasi V, Priolo F. Dactylitis, a term for different digit diseases. *Scand J Rheumatol* 2006;35:333-40.
- Kane D, Greaney T, Bresnihan B, Gibney R, Fitzgerald O. Ultrasonography in the diagnosis and management of psoriatic dactylitis. *J Rheumatol* 1999;26:1746-51.
- Fournié B, Margarit-Coll N, Champetier de Ribes TL, et al. Extrasynovial ultrasound abnormalities in the psoriatic finger. Prospective comparative power-Doppler study versus rheumatoid arthritis. *Joint Bone Spine* 2006;73:527-31.
- D'Agostino MA, Said-Nahal R, Hacquard-Bouder C, et al. Assessment of peripheral enthesitis in the spondyloarthropathies by ultrasonography combined with power Doppler: a cross-sectional study. *Arthritis Rheum* 2003;48:523-33.
- Gisondi P, Tinazzi I, El-Dalati G, et al. Lower limb enthesopathy in psoriatic patients without clinical signs of arthropathy: a hospital based case-control study. *Ann Rheum Dis* 2008;67:26-30.
- Grassi W, Salaffi F, Filippucci E. Ultrasound in rheumatology. *Best Pract Res Clin Rheumatol* 2005;19:467-85.
- Falsetti P, Frediani B, Filippou G, et al. Enthesitis of proximal insertion of the deltoid in the course of seronegative spondyloarthritis. An atypical enthesitis that can mime impingement syndrome. *Scand J Rheumatol* 2002;31:158-62.
- Alcalde M, Acebes JC, Cruz M. A sonographic enthesitis index (SEI) at lower limbs is a valuable tool in the assessment of ankylosing spondylitis. *Ann Rheum Dis* 2007;66:1015-9.
- Kiris A, Kaya A, Ozgocmen S, Kocakoc E. Assessment of enthesitis in ankylosing spondylitis by power Doppler ultrasonography. *Skeletal Radiol* 2006;35:522-8.
- de Miguel E, Cobo T, Muñoz-Fernández S, et al. Validity of enthesitis ultrasound assessment in spondylarthropathy. *Ann Rheum Dis* 2008;68:169-74.
- Frediani B, Falsetti P, Storri L, et al. Ultrasound and clinical evaluation of quadriceps tendon enthesitis in patients with psoriatic arthritis and rheumatoid arthritis. *Clin Rheumatol* 2002;21:294-8.
- Balint PV, Kane D, Wilson H, McInnes IB, Sturrock RD. Ultrasonography of enthesal insertions in the lower limb in spondyloarthropathy. *Ann Rheum Dis* 2002;61:905-10.
- Olivieri I, Barozzi L, Padula A, et al. Retrocalcaneal bursitis in spondyloarthropathy: assessment by ultrasonography and magnetic resonance imaging. *J Rheumatol* 1998;25:1352-7.
- Morel M, Boutry N, Demondion X, Legroux Gerot I, Cotten H, Cotten A. Normal anatomy of the heel enthesitis: anatomical and ultrasonographic study of their blood supply. *Surg Radiol Anat* 2005;27:176-83.