



ONLINE SUPPLEMENTARY DATA


Supplementary Figure 1. Forest plot of the effect of the body mass index (overweight and obesity > 25 kg/m² versus normal weight 18.5–25 or 20–25 kg/m²) on Disease Activity Score in 28 joints.




Supplementary Figure 2. Forest plot of the effect of the body mass index (overweight and obesity > 25 kg/m² versus normal weight 18.5–25 or 20–25 kg/m²) on Health Assessment Questionnaire score.



Supplementary Figure 3. Forest plot of the effect of the body mass index (overweight and obesity ≥ 25 kg/m² versus normal weight < 25 kg/m²) on radiographic joint damage (standardized means).



Supplementary Figure 4. Funnel plot of publication bias for (A) Disease Activity Score in 28 joints (DAS28) and (B) radiographic joint damage by Egger test.



References

1. Choe J-Y, Bae J, Lee H, Park S-H, Kim S-K. Lack association of body mass index with disease activity composites of rheumatoid arthritis in Korean population: cross-sectional observation. *Clin Rheumatol* 2014;33:485-92 (online 2013)
2. Van der Helm-van Mil AHM, van der Kooij SM, Allaart CF, Toes REM, Huizinga TWJ. A high body mass index has a protective effect on the amount of joint destruction in small joints in early rheumatoid arthritis. *Ann Rheum Dis* 2008;67:769-74.
3. Westhoff G, Rau R, Zink A. Radiographic joint damage in early rheumatoid arthritis is highly dependent on body mass index. *Arthritis Rheum* 2007;56:3575-82.
4. Baker JF, George M, Baker DG, Toedter G, Von Feldt JM, Leonard MB. Associations between body mass, radiographic joint damage, adipokines and risk factors for bone loss in rheumatoid arthritis. *Rheumatology (Oxford)* 2011;50:2100-7.
5. Ajeganova S, Andersson ML, Hafström I, for the BARFOT Study Group. Association of obesity with worse disease severity in rheumatoid arthritis as well as with comorbidities: A long-term followup from disease onset. *Arthritis Care Res (Hoboken)* 2013;65:78-87.
6. Brown RA, Spina D, Butt S, Summers GD. Long-term effects of anti-tumour necrosis factor therapy on weight in patients with rheumatoid arthritis. *Clin Rheumatol* 2012;31:455-61.
35. Fukuda W, Omoto A, Ohta T, Majima S, Kimura T, Tanaka T, et al. Low body mass index is associated with impaired quality of life in patients with rheumatoid arthritis. *Int J Rheum Dis* 2013;16:297-302.
7. Velpula UD, Agrawal S, Thomas J, Prabu VN, Rajasekhar L, Narsimulu G. Low body mass index is adversely associated with radiographic joint damage in Indian patients with early rheumatoid arthritis. *J Rheumatol* 2011;38:434-8.
8. Fukuda W, Omoto A, Ohta T, Majima S, Kimura T, Tanaka T, et al. Low body mass index is associated with impaired quality of life in patients with rheumatoid arthritis. *Int J Rheum Dis* 2013;16:297-302.