

Supplementary Data. Association between clinical and serological features of patients with antisynthetase syndrome and nailfold capillaroscopy measures.

		Capillaries, mean number/mm	p	Giant Capillaries, %	p	Giant Capillaries, mean number/mm	p	Micro- hemorrhages, %	p	Avascular Areas, %	p	Ramified Capillaries, %	p	SSc-like Pattern, %	p
Female/male ratio		8.5/9	ns	32.7/27.9	ns	0.17/0.85	ns	21.4/27.9	ns	21.4/14	ns	49.1/53.5	ns	38.4/27.9	ns
RP	±	8.4/8.8	ns	38.5/25.7	ns	0.16/0.13	ns	24.2/21.1	ns	28.6/12.8	0.006	47.3/52.3	ns	40.7/32.1	ns
Mechanic's hands	±	8.5/8.7	ns	37.5/27.1	ns	0.15/0.14	ns	22.5/22	ns	23.8/16.9	ns	57.5/44.9	ns	43.8/30.5	ns
Arthritis	±	8.6/8.7	ns	31.5/37	ns	0.16/0.13	ns	20.8/30.4	ns	20.1/21.7	ns	47/60.9	ns	35.6/39.1	ns
Myositis	±	8.9/7.7	ns	31/35.1	ns	0.14/0.19	ns	22.2/24.3	ns	17.1/32.4	0.036	50.6/48.6	ns	34.8/43.2	ns
ILD	±	8.5/9.4	ns	33.3/20.7	ns	0.14/0.17	ns	23.6/17.2	ns	21.8/6.9	ns	53.4/27.6	0.01	37.9/24.1	ns
ANA	±	8.5/9.1	ns	33.3/34.5	ns	0.17/0.09	ns	25/19.1	ns	22.2/8.8	ns	51.9/47.1	ns	38/29.4	ns
Anti-SSA	±	8.8/8.2	ns	36.7/25.8	ns	0.16/0.11	ns	21.4/22.7	ns	27.6/13.4	0.014	55.1/46.4	ns	38.8/33	ns
Anti-Jo1	±	8.2/9.6	< 0.001	39.6/16.9	0.001	0.18/0.07	0.001	27.6/13.8	0.031	27.6/4.6	< 0.001	53.7/44.6	ns	44/21.5	0.002

RP: Raynaud phenomenon; ILD: interstitial lung disease; ANA: antinuclear antibodies; ns: not significant.