

Arthritis Induced by Corticosteroid Crystals

ENRICO SELVI, MD; RENATO De STEFANO, MD; SAURO LORENZINI, BSc; ROBERTO MARCOLONGO, Professor, Section of Rheumatology, Department of Clinical Medicine and Immunological Science, University of Siena, Viale Bracci, I-53100 Siena, Italy. Address reprint requests to Dr. Selvi. E-mail: eselvi@katamail.com

A 65-year-old man with bilateral osteoarthritis of the knee presented a joint effusion of the left knee. The swollen joint was treated with a 40 mg intraarticular injection of triamcinolone hexacetonide. The next day, he presented with acute arthritis in the injected knee; the joint was swollen and tender and he was unable to walk. Septic arthritis was considered and the joint fluid was aspirated. Gross examination showed 35 ml of a thick, turbid, yellowish synovial fluid (Figure 1A). Leukocyte count was 13,000 cells/mm³ with 95% neutrophils. Gram and acridine orange stainings were negative. Wet preparations of the specimen with polarizing compensated microscopy showed numerous birefringent, pleomorphic intra- and extracellular crystals of the

corticosteroid (Figure 1B). He underwent joint lavage with 1 l of normal saline and recovered completely within one day. The conclusive diagnosis was triamcinolone hexacetonide crystal-induced arthritis^{1,2}.

REFERENCES

1. Schumacher HR, Reginato AJ. Synthetic depot corticosteroid crystals. In: Schumacher HR, Reginato AJ. Atlas of synovial fluid analysis and crystal identification. Philadelphia: Lea & Febiger; 1991:183-7.
2. McCarty DJ, Hogan JM. Inflammatory reaction after intra-synovial injection of microcrystalline adreno-corticosteroid esters. Arthritis Rheum 1964;7:359-61.




Figure 1. A. Macroscopic appearance of the synovial fluid aspirated from the left knee. B. Intracellular triamcinolone hexacetonide crystals. Compensated polarized light, $\times 100$.