

Onset of Neck Pain After a Motor Vehicle Accident: A Case-Control Study

NICOLA J. WILES, GARETH T. JONES, ALAN J. SILMAN, and GARY J. MACFARLANE

ABSTRACT. Objective. To assess the relative contribution of constitutional (individual) factors, pre-accident health, psychological and workplace psychosocial factors, and accident related (mechanical) factors in the development of neck pain (whiplash) following a motor vehicle accident.

Methods. We conducted a case-control study of drivers (ages 17–70 yrs) who reported a motor vehicle accident to their insurance company. A self-report mailed questionnaire retrospectively collected information on the driver's pre- and post-accident health, details of the accident, and other exposure data. Case/control status (post-accident neck pain) was ascertained using a preshaded manikin.

Results. In total, 26% of drivers reported post-accident neck pain. Women, younger individuals, and those with a history of neck pain were more likely to report neck pain following their accident (OR 1.50, 95% CI 0.98, 2.28; OR 1.62, 95% CI 0.96, 2.74; OR 1.75, 95% CI 1.09, 2.81, respectively). In addition, a number of accident related and psychosocial factors were independently associated with reporting post-accident neck pain: collision from behind (OR 2.55, 95% CI 1.41, 4.62); vehicle stationary at impact (OR 1.93, 95% CI 1.12, 3.33); collision severity (upper vs lowest tertile: OR 16.1, 95% CI 8.64, 30.1); not being at fault (OR 2.61, 95% CI 1.49, 4.59); and monotonous work (OR 2.19, 95% CI 1.19, 4.04). Based on these 8 factors, the likelihood of having neck pain increased from 7% with ≤ 2 risk factors to 62% with ≥ 5 .

Conclusion. Development of neck pain after a motor vehicle accident is a complex phenomenon resulting from the combined effects of constitutional, mechanical, and psychosocial factors. Using 8 such variables it is possible to identify those at high risk of developing neck pain. (J Rheumatol 2005;32:1576–83)

Key Indexing Terms:

EPIDEMIOLOGY

NECK PAIN

WHIPLASH

RISK FACTORS

New episodes of neck pain are very frequent in the general population, and a number of population based studies have investigated risk factors for its occurrence. A combination of constitutional (individual) and psychological factors have been shown to predict a new episode of neck pain¹. The importance of psychological², psychosocial³, and mechanical factors^{3,4} has been confirmed by others.

In the UK, it has been estimated that roughly 250,000 cases of whiplash result from motor vehicle accidents each year⁵. Neck pain is the cardinal feature of this common problem. While the body of evidence on prognosis for individuals suffering whiplash has been growing in recent years, there remains a dearth of information about the factors associated with the onset of neck pain following such accidents.

Studies examining the etiology of neck pain following vehicle trauma have suggested that female sex^{6–9}, rear-end collision^{9,10}, and a history of neck pain⁷ are important risk factors. Age has also been implicated, although results are conflicting; some suggest that older age is a risk factor⁶, while others report an increased risk among younger individuals^{8,9}. The focus of earlier work has been the relationship between vehicle occupant and crash related factors and the onset of neck pain. Epidemiological investigations of the onset of a variety of regional pain syndromes have shown the multifactorial etiology of these syndromes, with constitutional, psychological, psychosocial and mechanical factors all being implicated^{11–14}. However, the interplay of such factors has not been examined in relation to the onset of neck pain following trauma. We assessed the relative contribution of constitutional (individual) factors, prior health, psychological and workplace psychosocial factors, and accident related (mechanical) factors in the development of neck pain (whiplash) following a motor vehicle accident.

From the ARC Epidemiology Unit and the Unit of Chronic Disease Epidemiology, Division of Epidemiology and Health Sciences, University of Manchester, Manchester, United Kingdom.

Supported by the Association of British Insurers and the Arthritis Research Campaign (ARC), Chesterfield, United Kingdom.

N.J. Wiles, PhD, Lecturer in Epidemiology; G.T. Jones, PhD, Lecturer in Rheumatic Disease Epidemiology; A.J. Silman, MD, Professor of Rheumatic Disease Epidemiology, Epidemiology Unit; G.J. Macfarlane, MD, Professor of Epidemiology, Unit of Chronic Disease Epidemiology.

Address reprint requests to Prof. G.J. Macfarlane, Aberdeen Pain Research (Epidemiology) Group, Department of Public Health, School of Medicine, Polwarth Building, Foresterhill, Aberdeen AB25 2ZD, Scotland. E-mail: g.j.macfarlane@abdn.ac.uk

Accepted for publication March 29, 2005.

MATERIALS AND METHODS

A case-control study design was adopted based on the recruitment of individuals reporting an accident to a national insurance company in the UK.

Recruitment. At the time of the initial post-accident telephone contact with the insurance company, consecutive driver-policyholders (ages 17–70 yrs) were asked for their consent for study participation. Recruitment commenced March 1, 2002, and continued for a 14 month period. Anonymized details (age, sex, and whether the driver was at fault for causing the acci-

dent) were recorded for those who refused to participate at the time of the initial telephone contact in order to assess the external validity of the study findings. The study was approved by the University of Manchester Committee on the Ethics of Research on Human Beings.

Questionnaire data. A questionnaire was mailed to all drivers who consented to contact from the university and retrospectively collected details of the accident itself and driver's pre- and post-accident health. Two reminders were mailed to subjects who failed to respond to the initial questionnaire mailing(s), with the option of completing a short questionnaire over the telephone or by post.

Subject identification: cases and controls. Drivers were asked to indicate whether they had experienced neck pain after the accident based on a shaded neck manikin (Figure 1). Specifically, drivers were asked, "Since the accident, have you had any ache or pain which lasted for one day or longer, in the area shaded below?". The duration of one day or longer was used in order to exclude any transient episodes of neck pain. Individuals who answered positively to this question were classified as cases and those without neck pain were designated controls.

Cases were asked whether they had consulted their general practitioner (GP) because of their neck pain and whether they had neck pain on the day of completing the questionnaire. Those positive for the latter were asked to rate the severity of, and distress caused by, their neck pain on the day of questionnaire completion using two 100 mm visual analog scales (VAS), and were also asked to complete the Neck Disability Index (NDI)¹⁵.

The NDI is a validated 10 item questionnaire designed to ascertain the influence of neck pain on daily living (e.g., on personal care, reading, work or usual activities, sleeping, recreational activities). The subject is asked to rate the effect of their neck pain on the day of questionnaire completion on each of the 10 areas on a scale from zero (no impact or no difficulty related to neck pain) to 5 (worst possible scenario — unable to do). A "not applicable" option was added to 2 sections (lifting and driving), as pilot studies showed that some drivers indicated that they did not lift weights because of other health problems (e.g., back pain) or no longer had access to a car to drive. The total score for the 10 NDI items is then summed and multiplied by 2 to give a score from 0 to 100% (prorating where applicable).

Exposure data collection. Data on the following potential etiologic risk factors was recorded for both cases and controls on the self-questionnaire.

Health before the accident. Self-rating of general health; number of GP consultations in the year prior; pain in the month before the accident (subjects were asked to shade the area affected on blank body manikins); history of neck pain ever and in the month before the accident (using a pre-shaded neck manikin; Figure 1); Health Anxiety (HA) and Illness Behavior (IB) [derived from the Illness Attitude Scales (IAS)¹⁶]; psychological distress [using the 12 item General Health Questionnaire (GHQ)¹⁷]; and the number of somatic symptoms (using the Somatic Symptom Checklist¹⁸).

Accident related data. Type of vehicle; whether a seat belt was worn; posi-

tion of the headrest; severity of the collision using a 100 mm VAS; direction of the collision; whether the driver anticipated the collision; and speed of the driver's vehicle at the time of the impact.

Work related data. Those in paid employment (full or part-time) were asked to complete psychosocial work related questions¹⁴.

Finally, all drivers were asked whether they had claimed compensation for any injuries resulting from the accident.

Statistical analysis. All analyses were conducted using Stata version 6¹⁹.

The primary analysis focused on identifying predictors of neck pain using logistic regression. All variables significant at $p \leq 0.20$ univariately were entered into a multivariable model to permit identification of independent associations. This model was simplified using the likelihood-ratio test²⁰. All variables significant at $p < 0.10$ were retained. Previously excluded variables (univariate; $p > 0.20$) were added to the multivariable model to determine whether they contributed significantly; any that were then significant at $p < 0.10$ were retained.

Compensation is an often controversial element in the development and persistence of neck pain (whiplash) following a motor vehicle accident. However, it may be postulated that the likelihood of initiating a claim for compensation is related to the severity of neck pain. This hypothesis was explored within this dataset. Differences between those who did and did not claim compensation were examined and predictors of compensation identified using logistic regression. Further analyses were conducted to identify predictors of neck pain among those individuals who had not claimed compensation. Statistical modeling techniques were employed as previously.

Differences between those invited to participate at the time of the first contact with the insurance company and those subjects included in the final dataset for analysis were identified. Final multivariable logistic models were adjusted for any differences using sampling (or probability) weights²¹. Unweighted and weighted results were then compared.

RESULTS

Recruitment and response to mailing. During the 14 month recruitment period, 2723 drivers (ages 17–70 yrs) were invited to participate in the study (Figure 2). Of these, 1901 (69.8%) agreed to being sent a questionnaire by post.

Of the 1843 "appropriate" referrals, a questionnaire was completed by 1499 individuals (81.3%). In total, 1003 drivers completed the full-length questionnaire and 496 drivers completed the short version. The majority of these drivers (97%) reported being involved in a collision and were eligible for the study. To permit identification of factors associated with the onset of neck pain after a motor vehicle accident, those drivers ($n = 154$) who reported pain in the neck area (using a pre-shaded manikin) during the month before the accident were excluded. Further analysis was therefore conducted among the 847 drivers who completed full-length questionnaires and 448 drivers with short questionnaire data.

Characteristics of participants. Of the 847 drivers who were involved in a collision, who completed a full-length questionnaire, and who did not report pain in the area of the neck manikin in the month before the accident, 434 were female (51.2%). The median age at the time of the accident was 40.6 years [interquartile range (IQR) 33.1, 49.9]. Eighty-six percent of drivers ($n = 726$) were in paid employment (full or part-time). The median interval from the accident to questionnaire completion was 24 days (IQR 14, 40).

Twenty-six percent of drivers ($n = 218$) reported having




Figure 1. Manikin with neck region shaded.




Figure 2. Study recruitment and response to mailing.

pain in the area of the neck manikin after their motor vehicle accident and were designated the cases. In the majority of cases the onset of this neck pain was within 24 hours of the accident (92%), had resulted in a consultation with a GP (64%), and was still present and reported on the day of questionnaire completion (69%). Overall, 11.3% of cases reported disabling neck pain (defined as NDI \geq 20%) and 46% were seeking compensation. In total, 183 cases and 505 of those without neck pain (the controls) had complete data for further analyses.

Predictors of neck pain. Women and younger individuals had increased odds of reporting post-accident neck pain (Table 1). In addition, markers of health prior to the accident (number of GP consultations, history of neck pain, and higher scores on the Illness Behavior subscale of the IAS) and psychosocial factors (no-fault claim and monotonous work) were associated with increased odds of post-accident neck pain. A number of accident related factors (height of head-rest, direction of collision, speed of vehicle at time of impact, severity of collision) were also univariately associated (Table 1).

On multivariable analysis, 8 factors were identified as

being independently associated with post-accident neck pain (Table 2): female sex (OR 1.50), age < 36 years (OR 1.62), history of neck pain before the accident (OR 1.75), collision from behind (OR 2.55), being stationary at the time of impact (OR 1.93), collision severity (OR upper tertile 16.1), not being at fault (OR 2.61), and monotonous work (OR 2.19). Based on these 8 factors, the likelihood of having neck pain increased from 7% with \leq 2 risk factors to 62% with \geq 5 risk factors. Excluding perceived severity of collision (which may be particularly influenced by symptoms) from the list of predictors had little effect on model performance (increasing from 11% to 60%, respectively). Overall, the model had an accuracy of 81.3% in predicting those who would experience neck pain (positive predictive value 69%, negative predictive value 84%).

Compensation. Women were more likely to claim compensation, as were younger individuals, those who were regarded as not at fault for the accident, those in more severe collisions, those who were hit from behind, and those who were stationary at the time of the impact (Table 3). Those claiming compensation were significantly more likely to report neck pain on the day of questionnaire completion [84.7% vs 54.3%: per-

Table 1. Univariate predictors of neck pain.

	Cases	Controls	OR	95% CI
Sex				
Male	77	266	1.00	
Female	106	239	1.53	1.09, 2.16
Age at time of accident, yrs				
≥ 46	44	160	1.00	
36–45	58	173	1.22	0.78, 1.91
< 36	81	172	1.71	1.12, 2.62
Self-rated health in year prior to accident				
Excellent/very good	147	395	1.00	
Good/fair/poor	36	110	0.88	0.58, 1.34
No. of GP consultations in year prior to accident				
0–1	81	269	1.00	
≥ 2	102	236	1.44	1.02, 2.02
History of neck pain				
No	128	410	1.00	
Yes	55	95	1.85	1.26, 2.73
GHQ score				
< 3	172	449	1.00	
≥ 3	11	56	0.51	0.26, 1.00
Somatic Symptom Checklist				
0	124	318	1.00	
≥ 1	59	187	0.81	0.57, 1.16
Health Anxiety (IAS)				
0–2	62	178	1.00	
3–7	61	164	1.07	0.71, 1.61
≥ 8	60	163	1.06	0.70, 1.60
Illness Behavior (IAS)				
0–2	62	211	1.00	
3–5	64	175	1.24	0.83, 1.86
≥ 6	57	119	1.63	1.07, 2.49
Headrest height				
Correct height	135	402	1.00	
Too low	48	103	1.39	0.94, 2.06
Direction of collision				
Other	31	164	1.00	
From front	30	169	0.94	0.54, 1.62
From behind	122	172	3.75	2.40, 5.88
Anticipated collision				
No	121	297	1.00	
Yes	62	208	0.73	0.51, 1.04
Speed of vehicle				
Moving	67	347	1.00	
Stationary	116	158	3.80	2.67, 5.42
Fault (insurance company)				
At fault	26	237	1.00	
Not at fault	157	268	5.33	3.40, 8.38
Perceived severity of collision (100 mm VAS)				
< 13	16	232	1.00	
13–28	51	162	4.56	2.51, 8.29
≥ 29	116	111	15.2	8.57, 26.8
Satisfaction with job				
Very/Satisfied	136	387	1.00	
Neither, Dissatisfied or Very dissatisfied	47	118	1.13	0.77, 1.67
Monotonous work				
Never/Occasionally	153	457	1.00	
Half/Most of the time	50	48	1.87	1.14, 3.05
Hectic work				
Never/Occasionally	132	342	1.00	
Half/Most of the time	51	163	0.81	0.56, 1.18
Stressful work				
Never/Occasionally	155	412	1.00	
Half/Most of the time	28	93	0.80	0.50, 1.27
Job decision making				
Very often/Often	155	436	1.00	
Sometimes/v/seldom	28	69	1.14	0.71, 1.84
Job learning				
Very often/Often	113	322	1.00	
Sometimes/v/seldom	70	183	1.09	0.77, 1.55

GP: general practitioner, GHQ: General Health Questionnaire, IAS: Illness Attitude Scales.

Table 2. Independent predictors of neck pain.

Variable	OR	95% CI
Sex		
Male	1.00	
Female	1.50	0.98, 2.28
Age, yrs		
≥ 46	1.00	
36–45	0.97	0.56, 1.68
< 36	1.62	0.96, 2.74
History of neck pain		
No	1.00	
Yes	1.75	1.09, 2.81
Direction of collision		
Other	1.00	
From front	1.03	0.55, 1.91
From behind	2.55	1.41, 4.62
Vehicle stationary		
No	1.00	
Yes	1.93	1.12, 3.33
At fault		
Yes	1.00	
No	2.61	1.49, 4.59
Severity of collision (VAS)		
< 13	1.00	
13–28	4.35	2.29, 8.26
≥ 29	16.1	8.64, 30.1
Monotonous work		
Never/Occasionally	1.00	
Half/Most of the time	2.19	1.19, 4.04

centage difference 30.4% (95% CI 18.9, 41.9)]. Compensation claimants also reported more severe, disabling, and more distressing neck pain (Table 3). In addition, those claiming compensation reported a greater number of health related symptoms after the accident (median 3 vs 0; $p < 0.0001$, Mann-Whitney U test). Three factors were identified as independent predictors of compensation: female sex, not being at fault for the accident, and being hit from behind (Table 4).

Multivariable modeling identified 6 factors that were independently associated with the odds of reporting post-accident neck pain in those not claiming compensation (Table 5). Five of these factors were previously identified as predictors of neck pain in all subjects (Table 2).

Comparison of predictors of post-accident neck pain in all subjects and in those not claiming compensation. In order to compare the predictors of post-accident neck pain in all subjects and in those not claiming compensation, multivariable models were constructed containing all predictors identified in either group (Table 6). Five factors were identified as predictors of post-accident neck pain in both groups (history of neck pain, being in a stationary vehicle, not being at fault, rating the collision as more severe, and monotonous work). Women, younger individuals, and those hit from behind were more likely to report post-accident neck pain in the entire dataset. However, these factors did not reach statistical significance among non-compensation claimants. Those who anticipated the collision had decreased odds of report-

ing neck pain in those not claiming compensation; when included in the list of predictors for all subjects, a weaker effect was observed.

External validity and adjustments for probability weights. Those who completed the short questionnaire were younger and more likely to be male. However, there was no significant difference in onset of neck pain after the accident among respondents to the short (24.4%) and full-length (25.8%) questionnaire [difference, 1.4% (95% CI -3.6%, 6.3%)].

The age, sex, and claim-type profile of those included in the main analysis ($n = 688$) and the original 2665 (appropriately referred) participants invited by the insurance company were compared. Those included in the main dataset were more likely to be female [50.2% vs 42.8%, percentage difference 7.3% (95% CI 3.0, 11.6)] and were more likely to have been not at fault for the accident [61.8% vs 48.4%, percentage difference 13.4% (95% CI 9.1, 17.6)]. There was no difference in age at the time of the accident between those included in and excluded from the main dataset ($p = 0.41$, Mann-Whitney U test).

A weighted analysis using probability weights to account for the differences in sex and claim type reported above showed no substantial differences in the associations observed in the various multivariable logistic regression models (data not shown).

DISCUSSION

Principal findings. This study has shown that the onset of neck pain after a motor vehicle accident has, in common with other regional pain syndromes, a multifactorial etiology. A combination of 8 constitutional, mechanical, and psychosocial factors could be used to identify those at high risk of developing neck pain. Specifically, younger individuals, women, and those with a history of neck pain had 50% to 80% increased odds of reporting post-accident neck pain. A 2- to 3-fold increased odds was seen for those whose vehicles were hit from behind and who were stationary at the time of impact. Greater severity of the collision was also associated with increased odds of post-accident neck pain. Those not at fault for the accident and those whose jobs were boring or monotonous also had 2- to 3-fold increased odds of reporting neck pain following their motor vehicle accident.

Comparison with data from previous studies. Female sex⁶⁻⁹ and being involved in a rear-end collision^{6,9,10} have emerged as the 2 most consistent predictors of the onset of neck pain after a motor vehicle accident. The balance of evidence⁷⁻⁹ would suggest, in common with our findings, that younger individuals are more likely to experience post-accident neck pain, although opposing evidence exists⁶. One study also observed⁷ that a history of neck pain is associated with an increased likelihood of reporting a new episode of neck pain following a motor vehicle accident.

Several studies have, in common with ours, been based

Table 3. Differences between those claiming and not claiming compensation.

	Not Claiming Compensation, n = 596		Claiming Compensation, n = 87		Difference, %
	n	%	n	%	
Sex					
Male	305	51.2	37	42.5	
Female	291	48.8	50	57.5	-8.6 (-19.8, 2.5)
Claim type					
Fault or 50:50	257	43.1	4	4.6	
No fault	339	56.9	83	95.4	38.5 (32.6, 44.4)
General health					
Excellent/v. good	466	78.2	72	82.8	
Good, fair, poor	130	21.8	15	17.2	-4.6 (-13.2, 4.0)
GP visits in year prior to accident					
0-1	307	51.5	42	48.3	
≥ 2	289	48.5	45	51.7	-4.6 (-13.2, 4.0)
History of neck pain					
No	469	78.7	65	74.7	
Yes	127	21.3	22	25.3	4.0 (-5.7, 13.7)
GHQ score					
< 3	532	89.3	84	96.6	
≥ 3	64	10.7	3	3.5	-7.3 (-2.7, -11.9)
Somatic Symptom Checklist					
0	376	63.1	62	71.3	
≥ 1	220	36.9	25	28.7	-8.2 (-18.4, 2.1)
Anticipated accident					
No	361	60.6	56	64.4	
Yes	235	39.4	31	35.6	-3.8 (-14.6, 7.0)
Head rest					
No	469	78.7	64	73.6	
Yes	127	21.3	23	26.4	5.1 (-4.7, 15.0)
Direction of collision					
Other direction	187	31.4	6	6.9	
Front	182	30.5	16	18.4	
Behind	227	38.1	65	74.7	p < 0.001**
Speed of vehicle					
Stationary	387	64.9	25	28.7	
Moving	209	35.1	62	71.3	36.2 (25.9, 46.4)
	Median	(IQR)	Median	(IQR)	p
Age, yrs	40.0	(32.8, 47.7)	37.5	(29.1, 44.5)	0.05*
Health Anxiety score	5	(1, 9)	5	(1, 8)	0.59*
Illness Behavior score	3	(1, 5.5)	3	(2, 6)	0.25*
Severity of collision, VAS	16	(8, 30)	48	(28, 64)	< 0.0001*
Neck pain NDI score	17.9	(12, 28)	33	(23, 42)	< 0.0001*
Severity of neck pain, VAS	18	(9, 31)	30	(19, 44)	0.0004*
Distress of neck pain, VAS	12	(5, 25)	31.5	(13, 49)	0.0001*

* Mann-Whitney U test for continuous non-normal variables. ** Chi-square test. GHQ: General Health Questionnaire, IQR: interquartile range, NDI: Neck Disability Index.

on data from insurance company claimants^{9,10}, but others have been restricted to hospital inpatients and outpatients^{6,8}, or to accidents reported to the police⁷. The generalizability of previous findings has also been debatable due to poor response rates⁷. Earlier studies have collected limited data on both the subject (age and/or sex) and the accident itself (most often, the direction of collision). Given the complex etiology of other regional pain syndromes, it is perhaps surprising that, with the exception of history of neck pain⁷, none of the above studies collected data on prior health or

psychosocial factors. Indeed, we have shown that a combination of constitutional, accident related, and psychosocial factors are associated with the onset of neck pain following a motor vehicle accident.

Compensation. The evolution of a “compensation culture” within society is often quoted in the media and indeed, as noted, compensation often forms a controversial aspect in discussions surrounding whiplash. However, our findings suggest that the initiation of a compensation claim is linked closely to the severity of the pain. Compensation claimants

Table 4. Independent predictors of claiming compensation.

Variable	OR	95% CI
Sex		
Male	1.00	
Female	1.58	0.97, 2.57
Claim type		
Fault	1.00	
No-fault	13.5	4.74, 38.4
Direction of collision		
Other	1.00	
From front	4.71	1.75, 12.7
From behind	7.35	3.08, 17.6

Table 5. Independent predictors of neck pain among those not claiming compensation.

Variable	OR	95% CI
Anticipated collision		
No	1.00	
Yes	0.56	0.34, 0.92
History of neck pain		
No	1.00	
Yes	1.96	1.16, 3.33
Vehicle stationary		
No	1.00	
Yes	2.58	1.49, 4.46
At fault		
Yes	1.00	
No	1.89	1.03, 3.48
Severity of collision, VAS		
< 13	1.00	
13–28	3.02	1.53, 5.96
≥ 29	8.63	4.49, 16.6
Monotonous work		
Never/Occasionally	1.00	
Half/Most of the time	2.24	1.14, 4.40

reported more severe, disabling, and distressing pain than nonclaimants. Others have also suggested that compensation is likely to be a consequence of severe pain rather than the cause²². True malingering is also regarded as uncommon²³. However, we are unable to exclude the possibility that those who decide to claim compensation may then report more severe symptoms.

When we restricted our analyses to those subjects who had not claimed compensation, 5 out of the 8 predictors previously identified on multivariable analysis remained in the model. The independent associations with sex, age, and direction of collision weakened. Given the association between compensation and severity of neck pain, this subgroup analysis is likely to identify predictors of less severe neck pain rather than neck pain *per se*.

Strengths and weaknesses of this study. The case-control study design allowed us to investigate a wide range of potential etiological factors in the development of neck pain. Use of a self-questionnaire eliminated the potential for

observer bias that may exist in interview based studies. Further, differential selection of cases and controls was avoided by the identification of case status from responses to the self-questionnaire.

The possibility of recall bias exists due to the retrospective nature of data collection. This is most likely to affect the individual's recall of their health before the accident and may explain the protective effect of prior psychological distress in predicting neck pain that was seen univariately. Similarly, rating the severity of collision is likely to be influenced by the consequences of the accident. However, when severity of collision was excluded from the final multivariable model, the performance of the model was not substantially affected. It could be argued that objective measures of the accident (for example, independent rating of vehicle damage) may be better than self-report data. However, it is likely that subjective health related outcomes, such as pain, will be more closely linked to the individual's perception of the accident rather than objective measures (which may in turn be related to unrelated factors such as vehicle monetary value). Finally, given that this study was conducted among drivers reporting a motor vehicle accident to their insurance company, it is possible that we have underestimated the incidence of neck pain.

As we restricted our main analyses to those with complete full-length questionnaire data, we explored the external validity of our study findings by using probability weights to adjust for the differences in sex and claim type between those included and excluded from the main dataset. No substantial differences were observed. Further evidence for the validity of our findings was provided by comparison of the incidence of neck pain among those who completed the short and the full-length questionnaires (24% vs 26%).

The development of neck pain following a motor vehicle accident is a complex phenomenon resulting from the combined effects of constitutional, mechanical, and psychosocial factors. Using 8 such variables it is possible to identify those at high risk of developing neck pain after a motor vehicle accident. The relationship between these etiological factors and markers of prognosis requires further investigation.

ACKNOWLEDGMENT

We thank the claims advisors at Direct Line Insurance, Manchester, who invited drivers to take part in the study. In particular, we extend our thanks to the team leaders, Jane Ogle and Angela Robson. We are grateful to Mark Blything for clerical assistance. Finally, we thank all drivers who participated in the study.

REFERENCES

1. Croft PR, Lewis M, Papageorgiou AC, et al. Risk factors for neck pain: a longitudinal study in the general population. *Pain* 2001;93:317-25.
2. Linton SJ. A review of psychological risk factors in back and neck pain. *Spine* 2000;25:1148-56.
3. Palmer KT, Walker-Bone K, Griffin MJ, et al. Prevalence and occupational associations of neck pain in the British population. *Scand J Work Environ Health* 2001;27:49-56.

Table 6. Comparison of independent predictors of neck pain in all subjects and those not claiming compensation.

Variable	All Subjects, n = 688		Variable	Not Claiming Compensation, n = 596	
	OR	95% CI		OR	95% CI
Sex			Sex		
Male	1.00		Male	1.00	
Female	1.49	0.98, 2.28	Female	1.38	0.84, 2.27
Age, yrs			Age, yrs		
≥ 46	1.00		≥ 46	1.00	
36–45	1.00	0.57, 1.73	36–45	0.90	0.47, 1.73
< 36	1.68	0.99, 2.84	< 36	1.55	0.84, 2.85
Anticipated collision			Anticipated collision		
No	1.00		No	1.00	
Yes	0.76	0.47, 1.22	Yes	0.65	0.37, 1.14
History of neck pain			History of neck pain		
No	1.00		No	1.00	
Yes	1.76	1.10, 2.82	Yes	1.98	1.16, 3.39
Direction of collision			Direction of collision		
Other	1.00		Other	1.00	
From front	1.15	0.60, 2.19	From front	0.77	0.36, 1.63
From behind	2.39	1.31, 4.38	From behind	1.67	0.87, 3.19
Vehicle stationary			Vehicle stationary		
No	1.00		No	1.00	
Yes	1.98	1.14, 3.42	Yes	1.84	0.99, 3.44
At fault			At fault		
Yes	1.00		Yes	1.00	
No	2.72	1.54, 4.81	No	1.81	0.96, 3.41
Severity of collision, VAS			Severity of collision, VAS		
< 13	1.00		< 13	1.00	
13–28	4.43	2.33, 8.45	13–28	2.90	1.45, 5.80
≥ 29	16.4	8.77, 30.6	≥ 29	8.43	4.34, 16.4
Monotonous work			Monotonous work		
Never/Occasionally	1.00		Never/Occasionally	1.00	
Half/Most/All	2.17	1.17, 4.01	Half/Most/All	2.17	1.09, 4.31

4. Ariens GA, van Mechelen W, Bongers PM, Bouter LM, van der Wal G. Physical risk factors for neck pain. *Scand J Work Environ Health* 2000;26:7-19.
5. Lovell ME, Galasko CSB. Whiplash disorders — a review. *Injury* 2002;33:97-101.
6. Otremski I, Marsh JL, Wilde BR, McLardy Smith PD, Newman RJ. Soft tissue cervical spine injuries in motor vehicle accidents. *Injury* 1989;20:349-51.
7. Dolinis J. Risk factors for 'whiplash' in drivers: a cohort study of rear-end traffic crashes. *Injury* 1997;28:173-9.
8. Versteegen GJ, Kingma J, Meijler WJ, ten Duis HJ. Neck sprain after motor vehicle accidents in drivers and passengers. *Eur Spine J* 2000;9:547-52.
9. Berglund A, Alfredsson L, Jensen I, Bodin L, Nygren A. Occupant- and crash-related factors associated with the risk of whiplash injury. *Ann Epidemiol* 2003;13:66-72.
10. Bourbeau R, Desjardins D, Maag U, Laberge-Nadeau C. Neck injuries among belted and unbelted occupants of the front seat of cars. *J Trauma* 1993;35:794-9.
11. Macfarlane GJ, Thomas E, Papageorgiou AC, Croft PR, Jayson MIV, Silman AJ. Employment and physical work activities as predictors of future low back pain. *Spine* 1997;22:1143-9.
12. Papageorgiou AC, Macfarlane GJ, Thomas E, Croft PR, Jayson MIV, Silman AJ. Psychosocial factors in the workplace — Do they predict new episodes of low back pain? Evidence from the South Manchester Back Pain Study. *Spine* 1997;22:1137-42.
13. Macfarlane GJ, Hunt IM, Silman AJ. Role of mechanical and psychosocial factors in the onset of forearm pain: Prospective population based study. *BMJ* 2000;321:676-9.
14. Nahit ES, Hunt IM, Lunt M, Dunn G, Silman AJ, Macfarlane GJ. Effects of psychosocial and individual psychological factors on the onset of musculoskeletal pain: common and site-specific effects. *Ann Rheum Dis* 2003;62:755-60.
15. Vernon H, Mior S. The Neck Disability Index: a study of reliability and validity. *J Manipulative Physiol Ther* 1991;14:409-15.
16. Speckens AEM, Spinhoven P, Sloekers PPA, Bolk JH, van Hemert AM. A validation study of the Whately Index, the Illness Attitude Scales, and the Somatosensory Amplification Scale in general medical and general practice patients. *J Psychosom Res* 1996;40:95-104.
17. Goldberg D, Williams P. A user's guide to the General Health Questionnaire. Windsor, UK: NFER-Nelson; 1988.
18. Othmer E, DeSouza C. A screening test for somatization disorder (hysteria). *Am J Psychiatry* 1985;142:1146-9.
19. Stata statistical software: release 6.0. College Station, TX: Stata Corp.; 1999.
20. Hosmer DWJ, Lemeshow S. Applied logistic regression. New York: John Wiley & Sons; 1989.
21. Brick JM, Kalton G. Handling missing data in survey research. *Stat Methods Med Res* 1996;5:215-38.
22. Dworkin RH. Compensation in chronic pain patients: cause or consequence? *Pain* 1990;43:387-8.
23. Barry M. Whiplash injuries. *Br J Rheumatol* 1992;31:579-81.

Personal non-commercial use only. The Journal of Rheumatology Copyright © 2005. All rights reserved.