

# Decrease in Pregnancy Loss Rates in Patients with Systemic Lupus Erythematosus Over a 40-Year Period

CHRISTINE A. CLARK, KAREN A. SPITZER, and CARL A. LASKIN

**ABSTRACT. Objective.** To determine if there has been a statistically significant change in pregnancy loss and preterm delivery rates in patients with systemic lupus erythematosus (SLE).

**Methods.** We analyzed the pregnancy outcomes of our SLE patients over the past 3 years and reviewed the literature over the past 40 years. We extracted pregnancy loss and preterm delivery data from reports of postdiagnosis SLE pregnancies. Studies were grouped into 5-year periods and weighted according to sample size. Group means, calculated for each study period, were plotted using linear regression to determine significance, and compared with population norms for the same periods.

**Results.** The rate of loss in SLE pregnancies over the past 40 years decreased from a mean of 43% in 1960–1965 to 17% in 2000–2003 ( $r^2 = 0.648$ ). This approximates the pregnancy loss rate in the general US population. Preterm deliveries were not uniformly reported and were rarely stratified into spontaneous or physician-initiated. Prior to 1980, it was not possible to derive group means for each time period. From 1980 to 2002, however, there was a trend toward a decrease in preterm births in SLE pregnancies, although they continue to be more frequent in SLE than in the general population.

**Conclusion.** Improvements in disease management and perinatal monitoring have resulted in a significant decrease in pregnancy loss in SLE over the last 40 years and a trend toward decreased preterm deliveries over the last 20 years in comparison to the general population. These advances highlight the importance of collaboration between rheumatologists and perinatologists. Given these data, the description of SLE-associated pregnancy could be revised to reflect a more positive prognosis for mother and fetus. (J Rheumatol 2005;32:1709–12)

*Key Indexing Terms:*

SYSTEMIC LUPUS ERYTHEMATOSUS

PRETERM DELIVERY RATES

PREGNANCY LOSS

Systemic lupus erythematosus (SLE) continues to be described as a condition associated with poor pregnancy outcome despite advances in disease management and perinatal care. We wanted to determine if there has been a statistically significant change in pregnancy loss and preterm delivery rates in women with SLE over the last 40 years in comparison with the general population.

## MATERIALS AND METHODS

**Patient selection.** We reviewed the pregnancy outcomes (pregnancy loss, live birth, and preterm delivery) of our patients with SLE over the last 3 years. Preterm deliveries were defined as those spontaneously occurring at less than 37 weeks' gestation, in contrast to those precipitated by the physician as the result of deteriorating maternal or fetal well being.

**Literature review.** In a literature review using the US National Library of Medicine PubMed database, we extracted pregnancy loss (including both early and late losses) and preterm delivery data (whether specified as spon-

aneous or not) from literature published over the last 40 years for women diagnosed with SLE. Values for the total number of pregnancies, fetal loss rates, and preterm delivery rates were elicited from each study.

For comparison, 5-year grouped US population values for pregnancy losses from 1960 to 2000 and preterm births (defined as deliveries at < 37 weeks' gestation) from 1980 to 2000 were calculated from data provided by National Vital Statistics Reports<sup>1,2</sup>.

**Analysis.** Data were analyzed using Sigma Stat version 4.0 (SPSS, Chicago, IL, USA). Mean values were calculated for 5-year periods from 1960 to 2003, and values were plotted using linear regression analysis (with 95% confidence intervals, CI) to determine the significance of any trends over time.

## RESULTS

**Pregnancy outcome in our clinic.** Of 83 pregnancies in women with SLE attending our clinic over the last 3 years, 73 (83%) resulted in a live birth. Of the 73 deliveries, 50 had spontaneous onset of labor, 17 of which (34%) occurred prior to 37 weeks' gestation.

**Literature review and analysis.** We identified more than 35 studies published since 1963 with pregnancy loss data available for patients diagnosed with SLE<sup>3-37</sup>.

Studies were collated into 5-year periods as some years had no available SLE pregnancy data. Because the sample mean is sensitive to a few unusually large or small observa-

---

From the Departments of Medicine, Immunology, and Obstetrics and Gynecology, the University of Toronto; and the LifeQuest Centre for Reproductive Medicine, Toronto, Ontario, Canada.

C.A. Clark, BSc, Research Associate; K.A. Spitzer, BSc, Study Coordinator; C.A. Laskin, MD, Director, Obstetric Medicine Programme. Address reprint requests to Dr. C.A. Laskin, Suite 1800, 655 Bay Street, Toronto, ON M5G 2K4. E-mail: claskin@rogers.com

Accepted for publication March 7, 2005.

---

Personal non-commercial use only. The Journal of Rheumatology Copyright © 2005. All rights reserved.

tions, we used a trimmed sample mean. Studies with results that differed significantly from other studies in the same 5-year time period (< 25th or > 75th percentile) were not included in the analysis. Results were weighted accordingly by combining raw data from studies grouped into the 5-year periods starting with 1963.

*Change in fetal loss rate in SLE pregnancies.* The rate of loss in SLE pregnancies decreased over the past 40 years from a mean of more than 40% to a mean of 17%. Despite interclinic variation within time periods, overall there has been a significant trend toward increased live birth rates (Figure 1;  $r^2 = 0.648$ ). In comparison, pregnancy loss rates in the US have remained relatively stable since 1960: in 2000, it was reported that the pregnancy loss rate was 16.1%<sup>1</sup> (Figure 1).

*Change in preterm delivery rates in SLE pregnancies.* The definition of preterm delivery varied from  $\leq 36$  weeks to  $\leq 38$  weeks among available SLE studies from 1980 to 2002, and was not generally reported in the earlier literature. There was seldom any delineation between spontaneous or induced onset of preterm labor. We therefore analyzed data from individual studies (in contrast to grouped data) and found a trend toward a decrease in preterm delivery rate (< 37 weeks' gestation) from 1980 to 2000 (37.3% to 32%, respectively), but this was neither consistent nor statistically significant ( $r^2 = 0.418$ ; Figure 2). In contrast, there has been a slight but significant increase in the percentage of preterm births in the US over the same period (from 9.4% in 1981 to 12.1% in 2002;  $r^2 = 0.997$ )<sup>2</sup>.

## DISCUSSION

Our review was hampered by inconsistent reporting and variable definitions of events in the SLE literature. The term "fetal loss" comprised losses at all stages of pregnancy (from embryonic to stillbirth), and in many instances, the term "preterm delivery" (defined as < 36 to < 38 weeks' gestation) was not stratified as spontaneous or induced. In addition, statistical analysis had to account for sample sizes that varied from 11 to 108. Our results must be interpreted in the context of these difficulties. Nevertheless, it is apparent that there has been a significant decline in pregnancy loss rates in SLE over the last 40 years compared to the relatively stable rate in the general population over the same time period.

There are a number of factors contributing to the decline in pregnancy loss (and potentially preterm births) among women with SLE from both the maternal and the fetal standpoint. Disease management, particularly the identification and treatment of secondary antiphospholipid syndrome (APS), may be the most significant clinical advance in the last 2 decades, although there is still debate regarding the optimum therapeutic regimen for women with a history of thrombosis and/or recurrent pregnancy loss. There is also recognition that inactive disease in contrast to stable disease is an important consideration in reducing both maternal and fetal morbidity<sup>37</sup>.

In addition, as timing of delivery is a highly critical aspect of antenatal management, measures including fetal biometry, amniotic fluid volume, heart rate patterns, arterial and venous Doppler, and biophysical variables are now inte-




Figure 1. Change in rate of fetal loss in SLE pregnancies and in the US general population over that last 40 years. Data were grouped into 5-year periods (except the first period, 1963-65, and last period, 2001-2003).




Figure 2. Change in rate of preterm deliveries (< 37 weeks' gestation, spontaneous and induced) in SLE pregnancies and in the US general population over the last 20 years.

gral parts of comprehensive fetal evaluation<sup>38</sup>. Clinicians are able to intervene and induce delivery in cases of deteriorating maternal or fetal health. Depending upon the gestational age, these induced deliveries might even be termed "fortuitous live births," as 20 years ago, no intervention would have been possible and the pregnancy would have terminated from either maternal or fetal causes with a resultant intrauterine death.

Kitridou and Goodwin<sup>39</sup> reviewed pregnancy outcomes in patients with SLE over the last half century and also reported rates approaching the population norm, but there was no mention of change in preterm deliveries. The increase in fetal survival due to improved pregnancy management was also raised by Petri and Allbritton<sup>40</sup>, although the authors concluded (in 1993) that adverse pregnancy outcomes in lupus, including loss and preterm delivery, were still very common. In the intervening 12 years, that perception has gradually changed.

Pregnancy in women with SLE continues to be contraindicated in the context of severe organ disease or a history of life-threatening complications in previous pregnancies. For SLE patients without those superimposed conditions, however, there should be no contraindication for pregnancy, providing disease is clinically quiescent at the outset and there is appropriate planning and close prenatal monitoring.

We continue to advocate that SLE pregnancies be treated as high risk. However, based upon our review of the literature, we propose that the prevailing description of pregnancy in patients with SLE be revised to reflect advances in disease and perinatal management over the last 40 years.

## REFERENCES

- Ventura SJ, Mosher WD, Curtin SC, Abma JC, Henshaw S. Trends in pregnancies and pregnancy rates by outcome: estimates for the United States, 1976-96. National Center for Health Statistics. Washington, DC: DHHS Publication No. 2000-1934. *Vital Health Stat* 21 2000;56:1-47.
- Martin JA, Hamilton BE, Sutton PD, Ventura SJ, Menacker F, Munson ML. Births: final data for 2002. *Natl Vital Stat Rep* 2003;52:1-113.
- Mund A, Simson J, Rothfield N. Effect of pregnancy on the course of systemic lupus erythematosus. *JAMA* 1963;183:917-20.
- Estes D, Larson DL. Systemic lupus erythematosus and pregnancy. *Coin Obstet Gynecol* 1965;8:307-21.
- McGee CD, Makowski EL. Systemic lupus in pregnancy. *Am J Obstet Gynecol* 1970;107:1008-12.
- Fraga A, Mintz G, Orozco J, et al. Sterility and fertility rates, fetal wastage and maternal morbidity in systemic lupus erythematosus. *J Rheumatol* 1974;1:293-8.
- Grigor RR, Shervington PC, Hughes FRV, et al. Outcome of pregnancy in systemic lupus erythematosus. *Proc Roy Soc Med* 1977;70:99-100.
- Zurier RB, Agyros TG, Urman JD, et al. Systemic lupus erythematosus: management during pregnancy. *Obstet Gynecol* 1978;51:178-80.
- Devoe LD, Taylor RL. Systemic lupus erythematosus in pregnancy. *Am J Obstet Gynecol* 1979;135:473-9.
- Tozman ECS, Urowitz MB, Gladman DD. Systemic lupus erythematosus and pregnancy. *J Rheumatol* 1980;7:624-32.
- Hayslett JP, Lynn RI. Effect of pregnancy in patients with lupus nephropathy. *Kidney Int* 1980;18:207-20.
- Houser MT, Fish AJ, Tagatz GE, et al. Pregnancy and systemic lupus erythematosus. *Am J Obstet Gynecol* 1980;138:409-13.
- Zulman JJ, Talal N, Hoffman GS, et al. Problems associated with the management of pregnancies in patients with systemic lupus erythematosus. *J Rheumatol* 1980;7:37-49.
- Fine LG, Barnett EV, Danovitch GM, et al. Systemic lupus erythematosus in pregnancy. *Ann Intern Med* 1981;94:667-77.

15. Jungers P, Dougados M, Pelisser C, et al. Lupus nephropathy and pregnancy; report of 104 cases in 36 patients. *Arch Intern Med* 1982;142:771-6.
16. Varner MW, Meehan RT, Syrop CH, et al. Pregnancy in patients with systemic lupus erythematosus. *Am J Obstet Gynecol* 1983;145:1025-40.
17. Imbasciati E, Surian M, Bottino S, et al. Lupus nephropathy and pregnancy: a study of 26 pregnancies in patients with systemic lupus erythematosus and nephritis. *Nephron* 1984;36:46-51.
18. Lockshin MD, Reinitz E, Ruzin ML, et al. Lupus pregnancy. I. Case-control prospective study demonstrating absence of lupus exacerbation during or after pregnancy. *Am J Med* 1984;77:893-8.
19. Gimovsky ML, Montoro M, Paul RH. Pregnancy outcome in women with systemic lupus erythematosus. *Obstet Gynecol* 1984;63:686-92.
20. Mintz G, Niz J, Gutierrez C, et al. Prospective study of pregnancy in systemic lupus erythematosus. *J Rheumatol* 1986;13:732-9.
21. Borbrie G, Liote F, Houllier P, et al. Pregnancy in lupus nephritis and related disorders. *Am J Kidney Dis* 1987;9:339-43.
22. Meehan RT, Dorsey JK. Pregnancy among patients with systemic lupus erythematosus receiving immunosuppressive therapy. *J Rheumatol* 1987;14:252-8.
23. Simaopoulou-Mavridou A, Manoussakis MN, Mavridis AK, et al. Outcome of pregnancy in patients with autoimmune rheumatic disease before the disease onset. *Ann Rheum Dis* 1988;47:892-7.
24. McHugh NJ, Reilly PA, MacHugh LA. Pregnancy outcome and autoantibodies in connective tissue disease. *J Rheumatol* 1989;16:42-6.
25. Lockshin MD. Pregnancy does not cause systemic lupus erythematosus to worsen. *Arthritis Rheum* 1989;32:665-70.
26. Nossent JC, Swaak AJG. Systemic lupus erythematosus. IV. Analysis of the interrelationship with pregnancy. *J Rheumatol* 1990;17:771-6.
27. Wong KL, Chan FY, Lee CP. Outcome of pregnancy in patients with systemic lupus erythematosus: a prospective study. *Arch Intern Med* 1991;151:269-73.
28. Le Thi Huong D, Wechsler B, Piette J-C, Bletry O, Godeau P. Pregnancy and its outcome in systemic lupus erythematosus. *QJM* 1994;87:721-9.
29. Lima F, Buchanan NMM, Khamashta MA, Kerslake S, Hughes GRV. Obstetric outcome in systemic lupus erythematosus. *Semin Arthritis Rheum* 1995;25:184-92.
30. Johnson MJ, Petri M, Witter FR, Repke JT. Evaluation of preterm delivery in a systemic lupus erythematosus pregnancy clinic. *Obstet Gynecol* 1995;86:396-9.
31. Le Thi Huong D, Wechsler B, Vauthier-Brouzes D, et al. Outcome of planned pregnancies in systemic lupus erythematosus: a prospective study on 62 pregnancies. *Br J Rheumatol* 1997;36:772-7.
32. Rahman P, Gladman DD, Urowitz MB. Clinical predictors of fetal outcome in systemic lupus erythematosus. *J Rheumatol* 1998;25:1526-30.
33. Carmona F, Font J, Cervera R, Munoz F, Cararach V, Balasch J. Obstetrical outcome of pregnancy in patients with systemic lupus erythematosus. A study of 60 cases. *Eur J Obstet Gynecol Reprod Biol* 1999;83:137-42.
34. Kobayashi N, Yamada H, Kisheda T, et al. Hypocomplementemia correlates with intrauterine growth retardation in systemic lupus erythematosus. *Am J Reprod Immunol* 1999;42:153-9.
35. Georgious PE, Polii EN, Katsimbri P, Sakka V, Drosos AA. Outcome of lupus pregnancy: a controlled study. *Rheumatology Oxford* 2002;39:1014-9.
36. Cortes-Hernandez J, Ordi-Ros J, Paredes F, Casellas M, Castillo F, Vilardell-Tarres M. Clinical predictors of fetal and maternal outcome in systemic lupus erythematosus: a prospective study of 103 pregnancies. *Rheumatology Oxford* 2002;41:643-50.
37. Clark CA, Spitzer KA, Nadler JN, Laskin CA. Preterm deliveries in women with systemic lupus erythematosus. *J Rheumatol* 2003;30:2127-32.
38. Baschat AA, Hecher K. Fetal growth restriction due to placental disease. *Semin Perinatol* 2004;28:67-80.
39. Kitridou RC, Goodwin TM. The fetus in systemic lupus erythematosus. In: Wallace DJ, Hahn BH, editors. *Dubois' lupus erythematosus*. Philadelphia: Lippincott Williams & Wilkins; 2002:1023-40.
40. Petri M, Allbritton J. Fetal outcome of lupus pregnancy: a retrospective case-control study of the Hopkins Lupus Cohort. *J Rheumatol* 1993;20:650-6.