

Correspondence



INSTRUCTIONS FOR LETTERS TO THE EDITOR

Editorial comment in the form of a Letter to the Editor is invited. The length of a letter should not exceed 800 words, with a maximum of 10 references and no more than 2 figures or tables; and no subdivision for an abstract, methods, or results. Letters should have no more than 4 authors. Financial associations or other possible conflicts of interest should be disclosed.

Letters should be submitted via our online submission system, available at the Manuscript Central website: <http://mc.manuscriptcentral.com/jrheum> For additional information, contact the Managing Editor, The Journal of Rheumatology, E-mail: jrheum@jrheum.com

Cytokines in Collagen Disease-Related Atherogenesis

To the Editor:

We read with interest the report of Asanuma, *et al*¹ and the excellent editorial of Shovman, *et al*² describing the role of proinflammatory cytokines in the pathogenesis of atherosclerosis in systemic lupus erythematosus (SLE). Two interconnected pathogenetic systems are involved: a complex array of proinflammatory factors found in collagen diseases per se, and a similar in part but different complex of proatherogenic factors in atherosclerosis. Indeed, in the last decade, our knowledge of proinflammatory and immunological factors in the pathogenesis of atherosclerosis has markedly expanded³. One of the important factors, not mentioned in the editorial², is the group of secretory phospholipases A₂ (sPLA₂). Low molecular weight nonpancreatic secretory PLA₂ catalyze the hydrolysis of the sn-2 group in glycerophospholipids, producing lysophospholipids and nonesterified fatty acids. The best investigated sPLA₂, called IIA, has been defined as an acute-phase reactant, since its circulating levels increase markedly in acute inflammatory and infectious conditions. It has, however, been recognized that sPLA₂ levels may remain elevated for long periods of time in several chronic diseases with inflammatory features. Several studies have shown that cytokines such as interleukin 1 (IL-1), IL-6, tumor necrosis factor, and some others induce *de novo* synthesis and extracellular release of sPLA₂, thus suggesting that sPLA₂ serves as a common distal effector of these cytokines⁴. sPLA₂ IIA and a few other low molecular weight sPLA₂ (such as V and X) were found not only to participate in the metabolism of arachidonic acid but also in the hydrolysis of human high-density lipoprotein and low-density lipoprotein, releasing in turn a battery of proinflammatory factors.

We reported that a large proportion of patients with systemic lupus erythematosus⁵ and rheumatoid arthritis⁶ have high circulating sPLA₂ IIA. sPLA₂ IIA as well as other sPLA₂ such as V and X, along with lipoproteins, were found in the vascular wall and more so in the atheromatous areas⁷. sPLA₂ are not only hydrolyzing lipoproteins, but also induce mitogenesis and proliferation of human vascular smooth muscle cells and the release of prostaglandin E₂ and leukotriene B₄ from the cells⁸. Recently, sPLA₂ was found to play an important role in the pathogenesis of atherosclerosis^{7,9}. Its level was found to serve as an independent predictive factor of recurrent

events in acute coronary syndromes¹⁰. Considering that the induction of sPLA₂ is mediated through a variety of cytokines and that sPLA₂ are released extracellularly along with the above cytokines, secretory PLA₂ should be included in the group of proinflammatory and proatherogenic factors, and may potentially serve as a novel target for the prevention of atherosclerosis in collagen diseases.

WALDEMAR PRUZANSKI, MD, FRCPC, FACP, FACR, Professor of Medicine (Em), University of Toronto; JULIA KOPILOV, MSc; ARNIS KUKSIS, PhD, Banting and Best Department of Medical Research, University of Toronto, Toronto, Ontario, Canada. Address reprint requests to Dr. W. Pruzanski, Rosedale Medical Centre, 600 Sherbourne Street, Suite 803, Toronto, Ontario M4X 1W4, Canada. E-mail: cpruzanski@stikeman.com

REFERENCES

1. Asanuma Y, Chung CP, Oeser A, et al. Increased concentration of proatherogenic inflammatory cytokines in systemic lupus erythematosus: Relationship to cardiovascular risk factors. *J Rheumatol* 2006;33:539-45.
2. Shovman O, Gilburd B, Shoenfeld Y. The role of inflammatory cytokines in the pathogenesis of systemic lupus erythematosus-related atherosclerosis: A novel target for treatment? [editorial]. *J Rheumatol* 2006;33:445-7.
3. Ross R. Atherosclerosis — An inflammatory disease. *N Engl J Med* 1999;340:115-26.
4. Vadas P, Pruzanski W, Stefanski E, et al. Extracellular phospholipase A₂ secretion is a common effector pathway of interleukin-1 and tumour necrosis factor action. *Immunol Letts* 1991;28:187-94.
5. Pruzanski W, Goulding NJ, Flower RJ, et al. Circulating group II phospholipase A₂ activity and antilipocortin antibodies in systemic lupus erythematosus. Correlative study with disease activity. *J Rheumatol* 1994;21:252-7.
6. Lin MKS, Farewell V, Vadas P, et al. Secretory phospholipase A₂ as an index of disease activity in rheumatoid arthritis. Prospective double blind study of 212 patients. *J Rheumatol* 1996;23:1162-6.
7. Webb NR. Secretory phospholipase A₂ enzymes in atherogenesis. *Curr Opin Lipidol* 2005;16:341-4.
8. Pruzanski W, Kopilov J, Kuksis A. Pathogenesis of atherosclerosis in rheumatic disease. *Ann Rheum Dis* 2006;65 Suppl 11:463.
9. Murakami M, Kudo I. New phospholipase A₂ isoenzymes with a potential role in atherosclerosis. *Curr Opin Lipidol* 2003;14:431-6.
10. Mallat Z, Steg G, Benessiano J, et al. Circulating secretory phospholipase A₂ activity predicts recurrent events in patients with severe acute coronary syndromes. *J Am Coll Cardiol* 2005; 46:1249-57.

Dr. Chung, *et al* reply

To the Editor:

We thank Dr. Pruzanski and colleagues for their letter regarding our report showing that interleukin 6 is associated with burden of atherosclerosis in patients with systemic lupus erythematosus (SLE). They comment on the relationship between inflammation and atherosclerosis and suggest that, in addition to the comprehensive list of biomarkers presented in the editorial by Shovman, *et al*, increased concentrations of secretory phospholipases A₂ (sPLA₂) should be a candidate marker or mediator for atherosclerosis in patients with inflammatory disease. This suggestion is based on the increased concentrations of sPLA₂ in patients with SLE and rheumatoid arthritis, and its role in atherosclerosis. Thus, indeed, sPLA₂ should be considered as a mediator and, if so, as a potential target. In addition, other mechanisms that may be common to inflammation and atherosclerosis such as oxidative stress are of interest. However, because associations do not always imply causation, the challenge facing future research will be defining whether markers are also mediators.

CECILIA P. CHUNG, MD, MPH, Department of Medicine;
YU ASANUMA, MD, PhD, Department of Pharmacology, Vanderbilt
University School of Medicine, Nashville, Tennessee; PAOLO RAGGI,
MD, Department of Cardiology, Emory University School of Medicine,
Atlanta, Georgia; C. MICHAEL STEIN, MD, Departments of Medicine
and Pharmacology, Vanderbilt University. Address reprint requests to Dr.
C. Chung, Vanderbilt University School of Medicine, T-3219 MCN, 1161
21st Ave., Nashville, Tennessee 37232-2681, USA.
E-mail: c.chung@Vanderbilt.Edu

Dr. Shovman, *et al* reply

To the Editor:

We thank Dr. Pruzanski and coworkers for their interest in our editorial¹ and welcome their comments. We agree that the group of secretory phospholipases A₂ (sPLA₂) and especially sPLA₂ IIA are involved in the pathogenesis of atherosclerosis through their proinflammatory and proatherogenic effects. Recently several new studies provided evidence that the protective function of sPLA₂ IIA in inflammation is also indicated. In particular, the efficient bactericidal properties of sPLA₂ IIA resulting in decreased persistence of microbial pathogens in the vessel wall have been reported²⁻⁴. It is presumed that the ability of sPLA₂ IIA to attack *Staphylococcus aureus* and other Gram-positive bacteria lies primarily in the enzyme, in the binding to the bacterial cell wall, the penetration of the wall, and the hydrolytic attack on the phospholipids of the bacterial cell membranes^{2,3}. The protective effect of sPLA₂ IIA against Gram-positive and Gram-negative bacteria was verified also *in vivo* through investigations of sPLA₂ IIA-transgenic mice. The transgenic mice showed a significantly higher resistance to *Staphylococcus aureus* compared with the control animals⁴.

Along with antibacterial characteristics, the antithrombotic properties of sPLA₂ IIA have been described and associated with the inhibition of thrombin synthesis, decreasing the probability of thrombus formation⁵. Moreover, it was established that sPLA₂ IIA may be responsible for enhanced clearance of oxidatively modified lipoproteins during inflammation via the liver and adrenals⁶⁻⁸. Recently, the antiinflammatory properties of sPLA₂ IIA were established in an experimental model of carrageenin-induced pleurisy in rats⁹.

The extent to which an expression of sPLA₂ IIA has pathogenic or protective functions with respect to atherosclerosis depends possibly on whether expression of the enzyme as the consequence of an inflammatory reaction is induced locally in the vessel wall or systemically as the result of an acute-phase reaction¹⁰. Thus, local sPLA₂ IIA expression in the vessel wall may be connected with several pathogenic effects, and generally through increased phospholipolysis of oxidatively modified lipoproteins by sPLA₂ IIA, resulting in cellular lipid accumulation and foam cell formation¹⁰. On the other hand, the local expression of sPLA₂ may have a protective effect associated with bactericidal activity and inhibition of thrombin synthesis. Systemic expression of sPLA₂ IIA may engender the protective effect through removal of oxidatively modified lipoproteins from the bloodstream via the liver and to a lesser extent via the adrenals⁶⁻⁸.

The insightful discussion by Pruzanski, *et al* is timely and highly relevant from the scientific and clinical points of view, and additional investigations regarding the role of the sPLA₂ group in the pathogenesis of atherosclerosis are required.

ORA SHOVMAN, MD; BORIS GILBURD, MD, Center for Autoimmune Diseases and Department of Medicine B, Sheba Medical Center;
YEHUDA SHOENFELD, MD, FRCP, Laura Schwarz-Kipp Chair for Research of Autoimmune Diseases, Tel-Aviv University, Center for Autoimmune Diseases and Department of Medicine B, Sheba Medical Center and Sackler Faculty of Medicine, Tel-Aviv University. Address reprint requests to Prof. Y. Shoenfeld, Sheba Medical Center, Department of Medicine B, Tel Hashomer, Ramat-Gan, 52621, Israel.
E-mail: shoenfel@post.tau.ac.il

REFERENCES

1. Shovman O, Gilburd B, Shoenfeld Y. The role of inflammatory cytokines in the pathogenesis of systemic lupus erythematosus-related atherosclerosis: a novel target for treatment? [editorial]. *J Rheumatol* 2006;33:445-7.
2. Buckland AG, Wilton DC. The antibacterial properties of secreted phospholipases A₂. *Biochim Biophys Acta* 2000;1488:71-82.
3. Foreman-Wykert AK, Weinrauch Y, Elsbach P, Weiss J. Cell-wall determinants of the bactericidal action of group IIA phospholipase A₂ against Gram-positive bacteria. *J Clin Invest* 1999;103:715-21.
4. Laine VJ, Grass DS, Nevalainen TJ. Protection by group II phospholipase A₂ against *Staphylococcus aureus*. *J Immunol* 1999;162:7402-8.
5. Mounier CM, Hackeng TM, Schaeffer F, Faure G, Bon C, Griffin JH. Inhibition of prothrombinase by human secretory phospholipase A₂ involves binding to factor Xa. *J Biol Chem* 1998;273:23764-72.
6. Eckey R, Menschikowski M, Lattke P, et al. Increased hepatic cholesterol accumulation in transgenic mice overexpressing human secretory phospholipase A₂ group IIA. *Inflammation* 2004;28:59-65.
7. Labeque R, Mullon CJ, Ferreira JP, et al. Enzymatic modification of plasma low density lipoproteins in rabbits: a potential treatment for hypercholesterolemia. *Proc Natl Acad Sci USA* 1993;90:3476-80.
8. Tietge UJ, Maugeais C, Cain W, Rader DJ. Acute inflammation increases selective uptake of HDL cholesteryl esters into adrenals of mice overexpressing human sPLA₂. *Am J Physiol Endocrinol Metab* 2003;285:403-11.
9. Gilroy DW, Newson J, Sawmynaden P, Willoughby DA, Croxtall JD. A novel role for phospholipase A₂ isoforms in the checkpoint control of acute inflammation. *FASEB J* 2004;18:489-98.
10. Menschikowski M, Hagelgans A, Siegert G. Secretory phospholipase A₂ of group IIA: is it an offensive or a defensive player during atherosclerosis and other inflammatory diseases? *Prostaglandins Other Lipid Mediat* 2006;79:1-33.



Muscle Cramps Associated with Localized Scleroderma Skin Lesions: Focal Dystonia, Neuromyotonia, or Nerve Entrapment?

To the Editor:

Localized scleroderma is a relatively benign and self-limited condition, with manifestations mostly restricted to the skin and subdermal tissue without vascular or visceral involvement¹. The pathogenesis and etiology of this disease remain controversial. It has been speculated that localized scleroderma may develop as a response to neurologic injury². We describe the occurrence of muscle cramps in the distribution of skin lesions in 3 patients with localized scleroderma and review the literature.

Case 1. A 36-year-old woman with established linear scleroderma developed a new hypo- and hyperpigmented skin lesion extending linearly from the dorsum of the right hand, over the forearm, and continuing in a band-like distribution to the biceps and deltoid areas. As the skin lesion enlarged over several months, she complained of muscle cramping involving the

right triceps, biceps, and forearm muscles. Forearm cramps were elicited with chopping motions, and writing precipitated wrist and finger flexor muscle cramping. She was treated with hydroxychloroquine 200 mg bid. D-penicillamine and prednisone were previously discontinued. Cranial nerves, muscle strength, coordination, sensory examination, and deep tendon reflexes were normal. Cervical spine magnetic resonance imaging (MRI) revealed mild C5–C6 central disc herniation. Electromyography documented co-contraction of the right biceps and triceps muscles. Laboratory testing showed normal creatine kinase (CK) and a positive anti-nuclear antibody in a titer of 1:160 (normal < 40) with speckled staining, positive single-strand DNA antibody test, and hypergammaglobulinemia.

Case 2. A 40-year-old woman with morphea complained of episodic right arm pain and cramping of the right third and fourth fingers, sometimes accompanied by sustained ulnar deviation of the wrist. These symptoms were elicited by writing. On examination she had hyperpigmentation and atrophy extending from just beneath the axilla down the inner aspect of the arm to the distal forearm. There was a pale, slightly indurated and thickened area over the right buttock. Cranial nerves, muscle strength, coordination, sensory examination, and deep tendon reflexes were normal. There were no spontaneous or inducible involuntary movements. Electromyography was normal. Cervical spine MRI was normal. Laboratory testing included normal CK and negative anti-single-strand DNA antibody test.

Case 3. A 19-year-old woman with linear scleroderma developed dysesthesias and muscle cramping of the right fourth and fifth fingers. Muscle cramping was not task-induced. She also complained of dysesthesias and “a pulling sensation” in the left upper quadrant of the abdomen. She was previously treated with hydroxychloroquine. Mental status, muscle strength, coordination, and deep tendon reflexes were normal. Hyperpigmented skin with atrophic subcutaneous tissue extended from the medial aspect of the right forearm to the right hypothenar area. Similar hyperpigmented skin lesions with atrophy were present in the left upper abdomen. There was also atrophy of the left side of the tongue. Sensation to light touch was decreased with hypersensitivity to pinprick in the right C8, T1 and T2, and left T8 and T9 dermatomes. There were no spontaneous or inducible involuntary movements. Three years earlier, at the onset of symptoms, nerve conduction studies showed mild distal right ulnar mononeuropathy. MRI of the cervical spine showed mild hypertrophy of the right C3-C4 and C5-C6 facets with mild foraminal narrowing at C6-C7. Laboratory testing showed normal CK and an elevated anti-single-strand DNA antibody (219 units/ml; normal < 60).

Muscle cramp is a sudden involuntary shortening of the muscle occurring at rest or with muscle activation (contraction). Shortened muscle is more susceptible to muscle cramps³, as seen in atrophic regions affected by localized scleroderma. The presence of agonist-antagonist co-contraction in one of our patients is suggestive of focal dystonia. The peripheral origin

of dystonia remains controversial, and proposed mechanisms include altered sensory input or increased spinal cord excitability after peripheral nerve injury (entrapment)^{4,5}. Muscle cramps caused by continuous muscle fiber activity⁶ and neuromyotonia^{7,8} have been reported in 3 other patients with contiguous scleroderma skin lesions (Table 1).

Localized scleroderma may affect subcutaneous and deeper tissues, including muscles, ligaments, and bone, leading to stretching, angulation, or compression of nerves, followed by focal demyelination of motor nerve fibers. Dermatomal distributions of localized scleroderma skin lesions have been observed by some authors, and the skin lesions may follow a nerve injury². The occurrence of muscle cramps in the distribution of skin lesions may be attributable to nerve hyperexcitability (leading to neuromyotonia or dystonia) or ephaptic transmission, similar to hemimasticatory spasm⁹. In systemic sclerosis, distal axonopathy and focal nerve entrapment usually do not correspond to the distribution of skin lesions and are not associated with prominent muscle cramps¹⁰.

Based on the temporal and spatial correlation of skin lesions and muscle cramps in our patients, we propose that localized scleroderma may precipitate muscle cramps, possibly caused by local nerve injury. Additional studies are needed to define the pathophysiology of such muscle cramps and to establish the spectrum of neurologic complications of localized scleroderma.

SAŠA A. ZIVKOVIC, MD, VA Pittsburgh Healthcare System and Department of Neurology, University of Pittsburgh School of Medicine; DAVID LACOMIS, MD, Department of Neurology and Department of Pathology (Neuropathology), University of Pittsburgh School of Medicine; THOMAS A. MEDSGER Jr, Department of Medicine, Division of Rheumatology and Clinical Immunology, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA.

Address reprint requests to Dr. S.A. Zivkovic, Department of Neurology, University of Pittsburgh Medical Center, PUH F875, 200 Lothrop St., Pittsburgh, PA 15213. E-mail: zivkovics@upmc.edu

REFERENCES

1. Falanga V, Medsger TA Jr, Reichlin M, Rodnan GP. Linear scleroderma. Clinical spectrum, prognosis, and laboratory abnormalities. *Ann Intern Med* 1986;104:849-57.
2. Littman BH. Linear scleroderma: a response to neurologic injury? Report and literature review. *J Rheumatol* 1989;16:1135-40.
3. Layzer RB. The origin of muscle fasciculations and cramps. *Muscle Nerve* 1994;17:1243-9.
4. Ross MH, Charness ME, Lee D, Logigian EL. Does ulnar neuropathy predispose to focal dystonia? *Muscle Nerve* 1995;18:606-11.

Table 1. Neurologic complications associated with cutaneous manifestations of systemic and localized scleroderma.

Study	Age/Sex	Type of Scleroderma (age of onset, yrs)	Latency, yrs	Symptoms	EMG results	Treatment Response
Our study	36 F	Linear (33)	2*	Cramps and posturing of right arm and hand	Dystonia	Baclofen; DNT Tizanidine; good
Our study	40 F	Morphea (19)	17*	Cramps of right hand	Normal	Baclofen; poor
Our study	19 F	Linear (9)	8	Cramps of right hand	Ulnar neuropathy	Gabapentin; DNT
Papadimitriou ⁶	17 M	Morphea (17)	None	Twitching, painless cramps of the leg	CMFA	Phenytoin; good
Kumar ⁸	32 M	Morphea (30)	None	Muscle twitching	Neuromyotonia	Phenytoin; good
Benito-Leon ⁷	19 F	SSc (12)	4	Cramps of left arm and leg, task-elicited	Neuromyotonia	Carbamazepine; good

* Muscle cramping in the setting of an exacerbation of scleroderma. SSc: systemic sclerosis; CMFA: continuous muscle fiber activity; DNT: did not tolerate.

5. Sankhla C, Lai EC, Jankovic J. Peripherally induced oromandibular dystonia. *J Neurol Neurosurg Psychiatry* 1998;65:722-8.
6. Papadimitriou A, Chroni E, Anastasopoulos I, Avramidis T, Hadjigeorgiou G, Koutroumanidis M. Continuous muscle fiber activity associated with morphea (localized scleroderma). *Neurology* 1998;51:1763-4.
7. Benito-Leon J, Miguelez R, Vincent A, Masjuan J, de Blas G. Neuromyotonia in association with systemic sclerosis. *J Neurol* 1999;246:976-7.
8. Kumar A, Jain R, Daga J. Simultaneous occurrence of neuromyotonia and morphea: a cause-effect relationship? *J Neurol Neurosurg Psychiatry* 2006;77:802.
9. Cruccu G, Inghilleri M, Berardelli A, et al. Pathophysiology of hemimasticatory spasm. *J Neurol Neurosurg Psychiatry* 1994;57:43-50.
10. Poncelet AN, Connolly MK. Peripheral neuropathy in scleroderma. *Muscle Nerve* 2003;28:330-5.

Azathioprine and Tubulointerstitial Nephritis in Henoch-Schönlein Purpura

To the Editor:

We read with interest the report of Bir, *et al*, which suggests that azathioprine might have induced tubulointerstitial damage in their patient¹. We have also experienced similar cases in patients with severe Henoch-Schönlein nephritis (HSN) treated with azathioprine².

We reported that azathioprine might be an effective therapy in children with severe HSN by ameliorating the progression of immunologic renal injury². Nevertheless, 2 of the 10 patients with HSN who had been treated

with azathioprine showed definite tubulointerstitial nephritis at followup biopsy. In these 2 patients, massive proteinuria rapidly improved after cessation of azathioprine therapy, and one of them also showed decreased mesangial depositions of IgG, IgA, IgM, and C3 at a second biopsy. However, mild proteinuria had persisted throughout the course of the disease, which might be related to tubulointerstitial nephritis. Although it is very difficult to prove that azathioprine might have caused severe tubulointerstitial damage, our patients did not receive any other nephrotoxic drugs, and the duration of severe proteinuria was not long enough to cause such a tubulointerstitial injury. However, we detected the tubulointerstitial nephritis by renal biopsy at the end of the course of azathioprine treatment, because our cases did not show the characteristics of rapidly progressive renal failure.

Therefore, clinicians should be more cautious in their use of azathioprine in patients with vasculitis, and further studies should be performed to elucidate the relationship between azathioprine and tubulointerstitial nephritis.

JAE IL SHIN, MD; JAE SEUNG LEE, MD, Department of Pediatrics, Institute of Kidney Disease; HYEON JOO JEONG, MD; Department of Pathology, Yonsei University College of Medicine, 134 Shinchon-Dong, Seodaemun-Ku, 120-752, CPO Box 8044, Seoul, Korea.
E-mail: jsyonse@yumc.yonsei.ac.kr

REFERENCES

1. Bir K, Herzenberg AM, Carette S. Azathioprine induced acute interstitial nephritis as the cause of rapidly progressive renal failure in a patient with Wegener's granulomatosis. *J Rheumatol* 2006;33:185-7.
2. Shin JI, Park JM, Shin YH, et al. Can azathioprine and steroids alter the progression of severe Henoch-Schonlein nephritis in children? *Pediatr Nephrol* 2005;20:1087-92.