

Extracranial Meningioma Presenting as a Neck Mass in a Patient with Underlying Ankylosing Spondylitis

TIRA GALM, FIONA MacGREGOR, ANNE-MARIE SINCLAIR, JOHN HUNTER, and ALLAN McPHADEN

ABSTRACT. Extracranial meningiomas are rare. They account for only 2% of meningiomas compared to the much more common intracranial site. We describe a rare case of extracranial meningioma presenting as a neck lump in a patient with preexisting neck disease due to ankylosing spondylitis. Extracranial meningioma should be considered in the differential diagnosis of atypical neck lumps. Investigations including magnetic resonance imaging and histopathology are discussed, as well as the results of a literature review on extracranial meningiomas. (J Rheumatol 2006;33:1883–5)

Key Indexing Terms:

MENINGIOMA

NECK

ANKYLOSING SPONDYLITIS

Meningioma is a well recognized tumor of the central nervous system (CNS) accounting for 20% of all intracranial tumors; it is the second most common tumor of the CNS^{1,2}. Meningiomas are more common in women than in men, with a ratio of 2:1 and peak incidence at age 45 years³. Meningiomas have been described in children, where they tend to be more aggressive, multiple, and occasionally malignant⁴. At autopsy asymptomatic meningiomas are found in 1.4% of cases and are usually isolated findings, except when associated with Von Recklinghausen disease^{2,3,5}. This same trend is also seen in primary extracranial meningioma, which also shows a strong female predominance³.

Primary extracranial tumors are often difficult to diagnose and must be thoroughly evaluated in order to rule out an intracranial component. In the head and neck area, extracranial meningiomas have been described in various sites, such as the temporal bone, middle ear^{3,6–8}, nasal cavity⁹, parotid gland¹⁰, and paranasal sinus^{1,6}.

CASE REPORT

A 51-year-old woman with a long-standing history of ankylosing spondylitis (AS) was referred from rheumatology with a 6-month history of a swelling in the right posterior triangle. There was very limited neck movement as a result of AS, but the swelling was not associated with any pain.

On examination there was a 3 × 5 cm lump, which was hard in consistency and fixed, with no focal neurology (Figure 1). Further ENT examination

was unremarkable. An incisional biopsy was performed, and histological examination revealed features consistent with meningioma (Figure 2). There were no histological features suggestive of a malignant melanoma. Overall, the morphological features were in keeping with a meningioma.

Magnetic resonance imaging (MRI) was required to differentiate between a peripheral extension of an intracranial meningioma and the less common occurrence of an extracranial meningioma. Head and neck MRI showed a mass measuring 2.2 cm in width, 3 cm in anteroposterior diameter, and 4.8 cm in vertical diameter. The mass was lying in the posterior triangle of the right neck, deep to the levator scapula muscle, which the mass lifted laterally. It was insinuating into several intravertebral foramina, but did not occlude any of these foramina (Figure 3). There was no evidence of any extension into the spinal canal or intracranial involvement.

Overall histopathology confirmed meningioma, and MRI showed that this was the much rarer extracranial type. The lesion is benign with no mitotic activity. Our patient has therefore been initially managed with a “wait and watch” policy, as she is pain-free with no neurological symptoms.

DISCUSSION

Meningiomas are slow-growing lesions that represent about 20% of all intracranial tumors; however, primary extracranial meningiomas are relatively infrequent.

Considered by most to be benign hamartomas, they arise from meningotheial arachnoid cells, which are derived from the neuroectoderm. These cells are found in clusters at the tip of arachnoid membrane through the walls of dural sinuses into the sinus lumen, where they absorb cerebrospinal fluid. They are concentrated along the major dural sinuses and their tributary veins^{2,4–6,11}.

Meningiomas have a variety of easily recognizable histomorphological features. The cells appear as round, polygonal, ovoid epithelial, or endothelial types of cells. The nuclei have a delicate appearance and occasionally large nuclei may be present but are usually uniform. Mitoses are extremely rare⁶. The most common histologic classification of these tumors is based on the predominant cell type and includes 5 groups: syncytial, fibroblastic, transitional, angioblastic, and sarcomatous^{1,3,8}.

From Royal Berkshire and Battle Hospital, Reading; Gartnavel Hospital, and Glasgow Royal Infirmary, Glasgow, UK.

T. Galm, MBChB, MRCS, DO-HNS, SHO ENT, Royal Berkshire and Battle Hospital; F. MacGregor, MBChB, MRCS, Consultant ENT Surgeon; A-M. Sinclair, MBChB, MRCP, FRCA, Consultant Radiologist; J. Hunter, MBChB, MRCP, FRCP, Consultant Rheumatologist, Gartnavel Hospital; A. McPhaden, MBChB, MD, FRCPATH, Consultant Pathologist, Glasgow Royal Infirmary.

Address reprint requests to Dr. T. Galm, 12 Lakin House, Lakin Road, Warwick CV34 5BW, UK. E-mail: tiragalm@yahoo.com

Accepted for publication April 17, 2006.



Figure 1. The right posterior triangle neck lump.




Figure 2. Histopathological specimen shows nests of epithelioid cells with a whorled pattern containing psammoma body formation. There is no evidence of cellular pleomorphism and no mitotic activity. Immunocytochemistry showed focal EMA positivity with no staining for cytokeratins or S100. Hematoxylin and eosin; original magnification $\times 400$.

Many theories have been suggested for the mechanism of occurrence of extracranial arachnoid cells: Arachnoid cellular rests have been proposed to be formed by aberrant migration of arachnoid cells during embryogenesis^{2,8,12}. Alternatively, neural crest cells, schwann cells, arachnoid cells, and fibroblasts, which are mesodermal in origin, are cell types with potential to give rise to meningiomas^{2,8}.

Extracranial meningiomas are categorized in 4 groups, as described by Hoyer, *et al*¹³: (1) primarily intracranial tumors with extracranial extension; (2) benign-appearing intracranial meningioma with extracranial metastasis; (3) originating from

arachnoid cell rests of cranial nerve sheaths with extracranial growth; or (4) extracranial growth with no apparent connection with foramina or cranial nerve, presumably from embryonic rest cells.

According to the Hoyer classification our case is an extracranial tumor that fits into either category 3 or 4. It is possible that the tumor may have arisen from arachnoid cell rests from the spinal nerve sheath, as the tumor was seen to insinuate into several intravertebral foramina, which would be category 3. It could also be category 4, as the tumor was shown not to be occluding the foramina.




Figure 3. T1 axial post-gadolinium magnetic resonance image shows a mass that appears to be insinuating into several intravertebral foramina, but not occluding them.

Our patient presented with a rare case of neck swelling in the presence of another pathology in the cervical spine, and due to the gross neck fixation and curvature, the lump was masked for some time. The initial reaction by the patient and her clinician was that the lump was due to her AS. It is important therefore that the neck is examined carefully. Additional investigations including histopathology and imaging are needed to make an accurate diagnosis to exclude other causes of neck swelling.

REFERENCES

1. Daneshi A, Asghari A, Bahramy E. Primary meningioma of the ethmoid sinus: a case report. *Ear Nose Throat J* 2003;82:310-1.
2. Rietz DR, Ford CN, Kurtycz DF, et al. Significance of apparent intratympanic meningiomas. *Laryngoscope* 1983;93:1397-403.
3. Jabor MA, Amedee RG, Gianoli GJ, et al. Primary meningioma of the fallopian canal. *Medscape Italia* 2002;93:717-20.
4. Michel RG, Woodard BH. Extracranial meningioma. *Ann Otolaryngol* 1970;88:407-11.
5. Nicholas RD, Knighton RS, Chason JL, et al. Meningioma of the parotid region. *Laryngoscope* 1987;97:693-6.
6. Granich MS, Pilch BZ, Goodman ML. Meningiomas presenting in the paranasal sinuses and temporal bone. *Head Neck Surg* 1983;5:319-28.
7. Rojas R, Palacios E, D'Antonia M. An unusual primary intratympanic meningioma. *Ear Nose Throat J* 2004;83:607-8.
8. Michel RG, Woodard BH. Extracranial meningioma. *Ann Otolaryngol* 1970;88:407-11.
9. McGavran MH, Biller HF, Ogura JH. Primary intranasal meningioma. *Arch Otolaryngol* 1971;93:95-7.
10. Wolff M, Rankow RM. Meningioma of the parotid gland: An insight into the pathogenesis of extracranial meningioma. *Hum Pathol* 1971;3:453-9.
11. Rose WS, Makhija MC, Sattenspiel S. Meningioma presenting as a tumor in the neck. *Am J Radiol* 1980;134:1070-2.
12. Hallgrímsson J, Björnsson A, Gudmundsson G. Meningioma of the neck. *J Neurosurg* 1970;32:695-9.
13. Hoyer SJ, Hoar CS, Murray JE. Extracranial meningioma presenting as a tumour of the neck. *Am J Surg* 1960;100:486-9.