

Association of ACE Gene Polymorphism with Genetic Susceptibility to Systemic Lupus Erythematosus in a Chinese Population: A Family-based Association Study

JIANHUA XU, YONG WANG, FAMING PAN, JIM STANKOVICH, DONGQING YE, LI LIAN, KECHUN ZHANG, and CHANGHAI DING

ABSTRACT. *Objective.* Family-based association analysis was performed to investigate whether the angiotensin-converting enzyme (ACE)-G261T and ACE-A592G polymorphisms are risk factors for systemic lupus erythematosus (SLE) in a Chinese population.

Methods. A total of 119 patients with SLE from 95 nuclear families, aged from 14 to 78 years, who met the American College of Rheumatology 1997 criteria, were recruited, as were 316 family members of these patients. A family-based association study was used to explore the association between ACE gene polymorphisms (ACE-G261T and ACE-A592G) and SLE. The ACE gene polymorphisms were genotyped with restriction fragment length polymorphism-polymerase chain reaction.

Results. Using family-based association tests, the T allele of the ACE-G261T single-nucleotide polymorphism (SNP) was significantly associated with genetic susceptibility to SLE in an additive model ($Z = 2.877$, $p = 0.004$), a dominant model ($Z = 2.557$, $p = 0.011$), and a recessive model ($Z = 2.202$, $p = 0.028$). No association between the ACE-A592G SNP and SLE susceptibility was determined.

Conclusion. The T allele of the ACE-G261T SNP was associated with SLE in this Chinese population. (First Release Oct 15 2007; *J Rheumatol* 2007;34:2408–11)

Key Indexing Terms:

ANGIOTENSIN-CONVERTING ENZYME
SUSCEPTIBILITY

SINGLE-NUCLEOTIDE POLYMORPHISM
SYSTEMIC LUPUS ERYTHEMATOSUS

FAMILY-BASED ASSOCIATION TEST

Systemic lupus erythematosus (SLE) is an autoimmune disorder of unknown etiology that is known to occur in genetically susceptible patients influenced by environmental factors¹. It is likely that the genetics of SLE is complex, with many genes influencing susceptibility². A number of genetic association studies have identified various candidate lupus susceptibility genes. These include members of the HLA family, complement genes, mannose-binding protein, Fcγ receptors, interleukin 10 (IL-10)^{3,4}, and recently, angiotensin-converting enzyme (ACE)^{5,6}. ACE is expressed in a wide range of tissues including lung, vascular endothelium, kidney, heart, and testes⁵. ACE activates angiotensin I into angiotensin II

and inactivates bradykinin via the kallikrein–kininogen system, and plays a major role in the rennin–angiotensin system (RAS). Angiotensin II increases vascular smooth-muscle cell contraction and affects smooth-muscle proliferation, monocyte adhesion, and platelet adhesion and aggregation, being mediated either directly or via various factors such as endothelin, nitric oxide, and others⁶. Angiotensin II is also a potent proinflammatory modulator with the ability to augment and perpetuate immune responses in renal and non-renal tissues⁷. Since immunological abnormalities appear to be directly related to the disease process of SLE, it is reasonable to predict that ACE gene polymorphisms may contribute to SLE susceptibility by affecting the function of the ACE protein or the systemic or local concentrations of ACE, which subsequently alters the immunological response in patients with lupus. However, the role of ACE gene polymorphisms such as insertion/deletions (I/D) in the pathogenesis of SLE remains unresolved^{5,6,8}. A search for additional ACE gene polymorphisms may be helpful to elucidate the role of ACE in SLE. The aim of our study was, therefore, to determine whether 2 ACE gene polymorphisms, ACE-G261T and ACE-A592G, are associated with SLE in a Chinese population.

From the Department of Rheumatology, First Affiliated Hospital; Department of Epidemiology and Biostatistics, School of Public Health, Anhui Medical University, Hefei, China; Menzies Research Institute, University of Tasmania, Hobart; and The Walter and Eliza Hall Institute of Medical Research, Melbourne, Australia.

J. Xu, MD, Professor; Y. Wang, MMed; L. Lian, MMed, Department of Rheumatology, First Affiliated Hospital; F. Pan, PhD; D. Ye, PhD; K. Zhang, MMed, Department of Epidemiology and Biostatistics, School of Public Health, Anhui Medical University; C. Ding, MD, Menzies Research Institute; J. Stankovich, PhD, Menzies Research Institute and The Walter and Eliza Hall Institute of Medical Research.

Address reprint requests to Dr. C. Ding, Menzies Research Institute, Private Bag 23, Hobart, Tasmania 7000, Australia.

E-mail: changhai.ding@utas.edu.au

Accepted for publication July 24, 2007.

MATERIALS AND METHODS

Study participants. A family-based association study was carried out in 119 subjects with SLE (classified according to the American College of

Rheumatology 1997 revised criteria⁹) from 95 nuclear families and 316 other family members (age 37.4 ± 14.6 yrs). They were recruited from the Department of Rheumatology at First Affiliated Hospital, Anhui Medical University. Of the families, 47 (39.5%) had both parents of patients available, 27 (22.7%) had 1 parent available, and 45 (37.8%) had no parent available. The family criteria used for subject selection were as follows: (1) 2 parents and proband; (2) if 1 parent was missing or deceased, at least 1 or 2 normal siblings were recruited. The structures of families with SLE in our study are presented in Table 1. All subjects gave their written informed consent, and the study protocol was approved by the Medical Human Research Ethics Committee at Anhui Medical University.

Markers. Two single-nucleotide polymorphisms (SNP) of ACE gene were selected from the National Center for Biotechnology Information dbSNP database that met the following criteria: (1) the SNP were located in exons; (2) the SNP had high heterozygosity (selective standard: heterozygosity > 5%); and (3) the SNP were nonsynonymous. Of the 2 SNP residing in the ACE gene we selected, rs4303 (G → T transition; Ala → Ser) is located in exon 5 (at amino acid position 261), and rs12709426 (A → G transition; Asp → Gly) is located in exon 12 (at amino acid position 592).

Phlebotomy. Forearm venous blood samples were collected from each subject by venipuncture into 10-ml vacuum tubes containing EDTA and citrate and put on ice. The tubes were centrifuged at 2000 rpm for 10 min, and then the plasma was excluded from the cell pellet by pipetting. All samples were stored at -80°C .

Genotyping of ACE polymorphisms. Genomic DNA was extracted from the cell pellet in whole blood (QIAamp blood Kit, Qiagen) and stored at -20°C until the genotype analysis was performed. ACE-G261T polymorphism was genotyped by restriction fragment length polymorphism-polymerase chain reaction (RFLP-PCR) method using the following primers: sense and anti-sense primers were 5'-TCT ACC TGA ACC TCC ATG CC-3' and 3'-GGT CCT TAC CCA GCA GAT GA-5', respectively. The 108-bp PCR amplicon was digested by the HhaI restriction enzyme [total reaction volume 15 μl , including PCR product 10 μl , 10 \times buffer 1.5 μl , restriction enzyme 0.4 μl (5 u/10 μl), H₂O 3.1 μl], and then was separated by electrophoresis on 2.5% agarose gel. The PCR fragment for the heterozygous ACE-261G allele was digested into 3 fragments of 108, 32, and 76 base pairs. Each reaction mixture contained 20 ng of genome DNA, 0.15 μM of each primer, 0.4 mM dNTPs, and 2.5 u rTaq (heat-resistant) DNA polymerase (Takara; Otsu, Shiga, Japan), using a T-gradient thermocycler (Biometa, Göttingen, Germany). The amplification procedure consisted of initial denaturation at 96°C for 3 min, 35 cycles of denaturation at 96°C for 30 s, annealing at 58°C for 30 s, and extension at 72°C for 90 s, followed by a final extension at 72°C for 7 min until genotype analysis was performed following a standard protocol. PCR was performed in 10 μl volumes.

Statistical analysis. Statistical analysis was performed using the family-based association test (FBAT)¹⁰, an extension of the transmission disequilibrium test. The test was performed using dominant, recessive, and additive genetic models. The number of informative families is dependent on the genetic model used, and the statistical power of the FBAT depends on the number of informative families. Single-marker FBAT analysis was used to estimate the single-loci frequencies. Each test counts how often a specific locus is present in informative families with SLE. A positive Z statistic in a single-locus FBAT indicates that a specific allele is more frequently transmitted into patients with SLE in informative families than expected under the null

Table 1. Systemic lupus erythematosus (SLE) families in this study.

Category	No. of Siblings					
	1	2	3	4	5	6
2 parents	15	13	4	8	1	6
1 parent	19	2	2	1	1	2
0 parent	0	5	34	4	2	0
Total	34	20	40	13	4	8

hypothesis of no linkage and no association. It was tested whether the genotypes were in Hardy-Weinberg equilibrium.

RESULTS

A total of 119 patients with SLE from 95 nuclear families that contained the complete genotype were used in the data analysis. Additionally, 316 family members of these patients were genotyped. Table 1 shows the family structures of the recruited patients with SLE.

We performed screening of ACE-G261T SNP and ACE-A592G SNP in 48 randomly selected subjects. If the heterozygosity degree of a SNP exceeded 5%, we then assessed this in all subjects. In the screening analysis, we found the SNP of ACE-A592G had lower heterozygosity degree (heterozygosity = 0.021), so we discontinued the assessment of this SNP in the whole sample.

The frequencies of the ACE-G261T GG, GT, and TT genotypes in the 119 patients were 13.9%, 62.0%, and 24.1%, respectively. The frequencies of the G and T alleles in cases were 44.8% and 55.2%, respectively. The genotype distributions of the ACE-G261T polymorphism in cases did not significantly deviate from the Hardy-Weinberg proportions (chi-squared = 3.434, $p = 0.180$; Table 2). Single-marker FBAT analysis showed that the ACE-261T allele was significantly associated with susceptibility to SLE in an additive model ($Z = 2.877$, $p = 0.004$), a dominant model ($Z = 2.557$, $p = 0.011$), and a recessive model ($Z = 2.202$, $p = 0.028$) (Table 3). Genotype analysis showed that individuals with the T/T genotype had greater susceptibility to SLE than those without the genotype ($Z = 2.202$, $p = 0.028$; Table 4).

DISCUSSION

In our study, we found that the ACE-G261T (rs4303) polymorphism was significantly associated with genetic susceptibility to SLE, and demonstrated that the ACE-261T allele was frequently transmitted to affected offspring ($p = 0.004$). Therefore, the ACE gene may play an important role in influencing susceptibility to SLE in the Chinese population. The results of our study suggest that DNA sequence variation in the ACE gene influences the risk of developing SLE.

Previous studies of gene polymorphisms in SLE have been mostly limited to the ACE I/D polymorphism, with relatively small numbers of patients from various ethnic populations and different disease characteristics, and thus yielding highly discordant results. For example, some studies have demonstrated association of SLE with the ACE D allele¹¹⁻¹³, while others

Table 2. Genotype distributions of ACE-G261T polymorphism and Hardy-Weinberg expectation test in systemic lupus erythematosus.

SNP	Genotype	Observed, n	Expected, n	Chi-square	p
rs4303	G/G	17	24	3.434	0.180
	G/T	73	59		
	T/T	29	36		

Table 3. Associations between the ACE polymorphism and systemic lupus erythematosus by family-based association test (single-marker FBAT analysis) in additive, dominant, and recessive models.

Allele	Additive Model			Dominant Model			Recessive Model		
	No. of Families [†]	Z	p	No. of Families	Z	p	No. of Families	Z	p
261T	60	2.877	0.004	40	2.557	0.011	43	2.202	0.028
261G	60	-2.877	0.004	43	-2.202	0.028	40	-2.557	0.011

[†] When the number of informative families was less than 10, the test was not computed. The positive Z statistic indicates that the frequency of transmitted allele is positively associated with susceptibility to SLE.
* Significant at level = 0.05.

Table 4. Associations between ACE genotype and SLE by family-based association test.

Marker	Genotype	Frequency	Families [†]	S	E (S)	Var (S)	Z	p
rs4303	G/G*	0.139	40	8.00	15.720	9.116	-2.557	0.011
	G/T	0.620	55	27.00	26.262	12.241	0.211	0.833
	T/T*	0.241	43	25.00	18.018	10.058	2.202	0.028

[†] When the number of informative families was less than 10, the test was not computed. The positive Z statistic indicates that the frequency of transmitted allele is positively associated with susceptibility to SLE.
* Significant at level = 0.05.

have demonstrated either association with the I allele¹⁴⁻¹⁶ or no association with either allele^{5,7,17-19}. Results have also varied in terms of the specificity of the association, with some studies demonstrating association with SLE in general¹²⁻¹⁵, and other associations restricted to specific clinical subgroups, such as lupus nephritis (LN), active LN, or disease activity^{11,16,20}. One study reported the ACE I/D polymorphism was associated with SLE among non-Caucasians and the 23949 (CT)_{2/3} polymorphism was associated with LN among non-Caucasians²¹. Another multivariate analysis demonstrated associations between the ACE I/D, ACE (CT)_{2/3}, and Atg C-532T functional polymorphisms and LN among Asians²². In stratified analyses among LN cases according to high versus low glomerular filtration rate (GFR), associations remained significant for the ACE D and (CT)₂ alleles among Asian subjects with low GFR²². Careful review of these previous studies in SLE suggests that much of the discordance may be secondary to heterogeneity, study designs, power, disease state, phenotype definition, and ethnic variability. This also suggests the need to genotype additional ACE gene polymorphisms in order to understand the role of ACE in SLE.

We report for the first time in our study the association of the G261T SNP of the ACE gene with SLE in a Chinese population using a family-based association method. Further studies with larger sample sizes, specific ethnic subgroups, and additional polymorphisms in the region are needed to confirm and extend these results.

ACKNOWLEDGMENT

We thank all families with SLE for their enthusiastic participation in the genetic epidemiology. We also acknowledge assistance from the Department of Epidemiology and Biostatistics, School of Public Health, Anhui Medical University.

REFERENCES

- Klippel JH, Dieppe PA. Rheumatology. 2nd ed. London: Mosby; 1998:731-54.
- Tan FK, Arnett FC. The genetics of lupus. *Curr Opin Rheumatol* 1998;10:399-408.
- Kelly JA, Moser KL, Harley JB. The genetics of systemic lupus erythematosus: putting the pieces together. *Genes Immun* 2002;3 Suppl 1:S71-85.
- Nath SK, Kilpatrick J, Harley JB. Genetics of human systemic lupus erythematosus: the emerging picture. *Curr Opin Immunol* 2004;16:794-800.
- Kaufman KM, Kelly J, Gray-McGuire C, et al. Linkage analysis of angiotensin-converting enzyme (ACE) insertion/deletion polymorphism and systemic lupus erythematosus. *Mol Cell Endocrinol* 2001;177:81-5.
- Uhm WS, Lee HS, Chung YH, et al. Angiotensin-converting enzyme gene polymorphism and vascular manifestations in Korean patients with SLE. *Lupus* 2002;11:227-33.
- Suzuki Y, Ruiz-Ortega M, Egido J. Angiotensin II: a double edged sword in inflammation. *Nephrol* 2000;13 Suppl 3:S101-10.
- Lee YH, Rho YH, Choi SJ, Ji JD, Song GG. Angiotensin-converting enzyme insertion/deletion polymorphism and systemic lupus erythematosus: a metaanalysis. *J Rheumatol* 2006;33:698-702.
- Hochberg MC. Updating the American College of Rheumatology revised criteria for the classification of systemic lupus erythematosus [letter]. *Arthritis Rheum* 1997;40:1725.
- Horvath S, Xu X, Lake SL, et al. Family-based tests for associating haplotypes with general phenotype data. Application to asthma genetics. *Genet Epidemiol* 2004;26:61-9.
- Harjacek M, Anic B, Bosnic D, et al. The role of genetic polymorphism of angiotensin-converting enzyme in lupus nephritis [abstract]. *Arthritis Rheum* 1999;42 Suppl:S302.
- Pullmann R Jr, Lukac J, Skerenova M, et al. Association between systemic lupus erythematosus and insertion/deletion polymorphism of the angiotensin converting enzyme (ACE) gene. *Clin Exp Rheumatol* 1999;17:593-6.
- Prkacin I, Novak B, Sertic J, et al. Angiotensin-converting enzyme gene polymorphism in patients with systemic lupus. *Acta Med Croatica* 2001;55:73-6.

14. Tassioulas IO, Aksentijevich I, Salmon JE, et al. Angiotensin converting enzyme gene polymorphisms in systemic lupus erythematosus: decreased prevalence of DD genotype in African-American patients. *Clin Nephrol* 1998;50:8–13.
15. Sato H, Akai Y, Iwano M, et al. Association of an insertion polymorphism of angiotensin-converting enzyme gene with the activity of systemic lupus erythematosus. *Lupus* 1998;7:530–4.
16. Akai Y, Sato H, Iwano M, et al. Association of an insertion polymorphism of angiotensin-converting enzyme gene with the activity of lupus nephritis. *Clin Nephrol* 1999;51:141–6.
17. Molad Y, Gal E, Magal N, et al. Renal outcome and vascular morbidity in systemic lupus erythematosus (SLE): lack of association with the angiotensin-converting enzyme gene polymorphism. *Semin Arthritis Rheum* 2000;30:132–7.
18. Douglas G, Reilly C, Dooley MA, et al. Angiotensin-converting enzyme (insertion/deletion) and endothelial nitric oxide synthase polymorphisms in patients with systemic lupus erythematosus. *J Rheumatol* 2004;31:1756–62.
19. Provieri SRS, Sens YAS. Polymorphisms of the renin-angiotensin system genes in Brazilian patients with lupus nephropathy. *Lupus* 2005;14:356–62.
20. Guan T, Lui Z, Chen Z. Angiotensin-converting enzyme gene polymorphism and the clinical pathological features and progression in lupus nephritis. *Zhonghua Nei Ke Za Zhi* 1997;36:461–4.
21. Parsa A, Peden E, Lum RF, et al. Association of angiotensin-converting enzyme polymorphisms with systemic lupus erythematosus and nephritis: analysis of 644 SLE families. *Genes Immun* 2002;3 Suppl:S42–6.
22. Parsa A, Lovett DH, Peden EA, et al. Renin–angiotensin system gene polymorphisms predict the progression to renal insufficiency among Asians with lupus nephritis. *Genes Immun* 2005;6:217–24.