

Different Genetic Effects of Interferon Regulatory Factor 5 (*IRF5*) Polymorphisms on Systemic Lupus Erythematosus in a Korean Population

HYOUNG DOO SHIN, IL KIM, CHAN-BUM CHOI, SOO OK LEE, HYE WON LEE, and SANG-CHEOL BAE

ABSTRACT. *Objective.* In an effort to replicate additional associations of interferon regulatory factor 5 (*IRF5*) polymorphisms with systemic lupus erythematosus (SLE) in an Asian population, we examined those genetic effects in a Korean SLE cohort.

Methods. Each *IRF5* polymorphism was genotyped in 1565 subjects using the TaqMan method and examined to determine whether it could explain the association with SLE.

Results. Three single-nucleotide polymorphisms (*IRF5-15-1*, *rs2070197*, and *rs10488631*), which showed strong and/or independent association in Caucasian populations, were not polymorphic in our Korean population. Association analysis revealed different genetic effects in Koreans compared with Caucasian populations. In addition, conditional analysis suggested independent genetic effects of 3 variant groups in the Korean population.

Conclusion. We demonstrate different genetic effects of *IRF5* polymorphisms on the risk of SLE according to ethnicity. (First Release Oct 1 2008; J Rheumatol 2008;35:2148–51; doi:10.3899/jrheum.080124)

Key Indexing Terms:

INTERFERON REGULATORY FACTOR 5

SYSTEMIC LUPUS ERYTHEMATOSUS

POLYMORPHISM

Recently, several studies have provided convincing evidences that interferon regulatory factor 5 [*IRF5* (MIM 607218)] polymorphisms are significantly associated with systemic lupus erythematosus [SLE (MIM 152700)]^{1–4}. We previously reconfirmed the genetic association of *IRF5* polymorphisms with the risk of SLE in a Korean population³. In a subsequent study of the same group from which *IRF5* association with SLE had been identified in Caucasian populations, 3 functional alleles of *IRF5* were found to be associated with SLE: the previously described exon 1B splice site variant; a 30-bp in-frame insertion/deletion variant of exon 6 that alters a proline-, glutamic acid-, serine-, and threonine-rich domain region; and a variant in a con-

served polyA₂ signal sequence that alters the length of the 3' untranslated region (UTR) and stability of *IRF5* mRNA among Caucasian populations⁵. In addition, the *CGGGG*indel located in the promoter region of *IRF5*, only 64 bp upstream of exon 1a of *IRF5*, was reported as an additional candidate for a causal variant of *IRF5*⁵.

In an effort to replicate the additional associations with SLE in an Asian population, we examined those genetic effects in our SLE cohort from the Korean population.

MATERIALS AND METHODS

A total of 593 Korean patients with SLE [mean age 32.36 yrs (6.99–70.7), male/female ratio 35/558] who fulfilled the 1997 American College of Rheumatology (ACR) criteria for SLE⁶ were consecutively enrolled in our study between September 1998 and February 2005 from the Hospital for Rheumatic Diseases, Hanyang University, Seoul. The following clinical and laboratory data were obtained: sex, age, ages at first symptom onset and clinical diagnosis, ACR diagnosis, and Systemic Lupus International Collaborating Clinics (SLICC)/ACR damage index⁷. As a control group, we included 972 healthy ethnically matched subjects [mean age 37.2 yrs (range 16.6–78.6), male/female ratio 139/832]. Written informed consent was obtained from each subject, and the protocol was approved by the Institutional Review Board of Hanyang University Hospital.

Six polymorphisms (*CGGGG*indel, *IRF5-15-1*, *Exon6*indel, *rs2070197*, *rs10954213*, and *rs10488631*), which showed strong and/or independent association in Caucasian populations^{5,8}, were genotyped using TaqMan^{®9} among our cases with SLE and controls from the Korean population.

Chi-squared analyses were used to evaluate the significance of differences in genotype and allele frequencies in the case-control samples using SAS. For conditional analysis of *IRF5* polymorphisms, the variants were subdivided into 3 groups according to linkage disequilibrium (LD) values

From the Department of Life Science, Sogang University; Department of Genetic Epidemiology, SNP Genetics, Inc.; and Department of Rheumatology, the Hospital for Rheumatic Diseases, Hanyang University, Seoul, Korea.

Supported in part by a grant of the Korea Health 21 R&D Project, Ministry of Health and Welfare, Republic of Korea (01-PJ3-PG6-01GN11-0002).

H.D. Shin, PhD, Laboratory of Genomic Diversity, Department of Life Science, Sogang University, and Department of Genetic Epidemiology, SNP Genetics Inc.; S.O. Lee, MS; H.W. Lee, MS, Department of Genetic Epidemiology, SNP Genetics, Inc.; I. Kim, MD; C-B. Choi, MD; S-C. Bae, MD, PhD, MPH, Department of Rheumatology, the Hospital for Rheumatic Diseases, Hanyang University.

Address reprint requests to Prof. S-C. Bae, Hospital for Rheumatic Diseases, Hanyang University Medical Center, 17 Haengdang-Dong, Sungdong-Gu, Seoul 133-792, Korea. E-mail: scbae@hanyang.ac.kr

Accepted for publication July 3, 2008.

Personal non-commercial use only. The Journal of Rheumatology Copyright © 2008. All rights reserved.

among them (Table 1). Conditional p values were estimated and permuted using the WHAP software¹⁰.

RESULTS

The genotype distributions of 10 markers, including 4 markers that had been reported in our previous study, are shown in Table 2. All polymorphisms except *rs729302* showed significant differences between Korean and Caucasian populations. Interestingly, 3 single-nucleotide polymorphisms (SNP; *IRF5-15-1*, *rs2070197*, and *rs10488631*) that showed strong and/or independent association in Caucasian populations were not polymorphic in the Korean population. By pairwise LD analysis using 7 polymorphic markers in the Korean population, 2 sets of tight linkages (*CGGGGindel*:

rs2004640: *rs752637* and *Exon6indel*: *rs10954213*: *rs2280714*) ($D' > 0.97$) were observed (Table 1).

First, genetic effects of the polymorphisms were individually analyzed. We found that 4 variants in the promoter region (*rs729302*, *CGGGGindel*, *rs2004640*, and *rs752637*) and 2 haplotypes (*ht2* and *ht3*) showed strong associations ($p = 0.003$ – 0.00002). However, the signals of 3 variants (*CGGGGindel*, *rs2004640*, and *rs752637*) might not be independent because of tight LD among them ($D' > 0.97$).

Next, we investigated whether each polymorphism could explain the association to SLE using conditional logistic regression. The variants were subdivided into 3 groups according to LD values among them (Table 1). The results of conditional analysis were as follows. (1) When condi-

Table 1. Linkage disequilibrium among *IRF5* polymorphisms (r^2 and ID').

	<i>rs729302A>C</i>	<i>CGGGGindel</i>	<i>rs2004640G>T</i>	ID' <i>rs752637T>C</i>	<i>Exon6indel</i>	<i>rs10954213G>A</i>	<i>rs2280714T>C</i>
<i>rs729302A>C</i>	—	0.878	0.833	0.321	0.713	0.71	0.762
<i>CGGGGindel</i>	0.063	—	0.981	0.978	0.666	0.668	0.943
<i>rs2004640G>T</i>	0.160	0.348	—	0.984	0.36	0.358	0.937
r^2 <i>rs752637T>C</i>	0.032	0.256	0.714	—	0.467	0.466	0.937
<i>Exon6indel</i>	0.223	0.083	0.068	0.155	—	1	0.996
<i>rs10954213G>A</i>	0.222	0.085	0.068	0.155	1	—	0.996
<i>rs2280714T>C</i>	0.168	0.114	0.312	0.419	0.663	0.661	—

Table 2. Frequencies of *IRF5* polymorphisms in Korean, Japanese, Chinese, and Caucasian populations.

rs	Region	Genotype			Allele	Korean (n = 1565)	Frequency			p*
							Caucasian (n = 934 ^a , 533 ^b , 1048 ^c)	Japanese ^d (n = 3540)	Chinese ^e (n = 190)	
<i>rs729302</i>	Promoter	A	AC	C	A	0.70	0.69 ^a	0.73		0.6163
<i>CGGGGindel</i>	Promoter	Del	Indel	Ins	Ins	0.16	0.50 (0.44) ^c	0.12		4.062×10^{-141}
<i>rs2004640</i>	Promoter	G	GT	T	T	0.345	0.57 ^a	0.33	0.29	5.2×10^{-28}
<i>rs752637</i>	Promoter	T	CT	C	C	0.42	0.68 ^a	0.41		3.7×10^{-36}
<i>IRF5-15-1</i>	Intron2	C	CT	T	T	0.00	0.18 (0.14) ^c			8.8312×10^{-136}
<i>Exon6indel</i>	Exon6	Del	Indel	Ins	Ins	0.50	0.51 ^b			0.5934
<i>rs2070197</i>	3'UTR	T	CT	C	C	0.00	0.13 ^b		0.00	4.5×10^{-32}
<i>rs10954213</i>	3'UTR	A	AG	G	A	0.50	0.63 ^b		0.49	4.6×10^{-9}
<i>rs10488631</i>	3'down	T	CT	C	C	0.00	0.17 (0.12) ^c			3.112×10^{-129}
<i>rs2280714</i>	3'down	T	CT	C	T	0.60	0.72 ^a	0.56		1.6×10^{-11}

^a Graham, et al¹. Data of 4 SNP are from Caucasian parental chromosomes used in transmission disequilibrium test analysis. ^b Kozyrev¹⁷. Data of 4 SNP are from Argentinian, Spanish, and German combined subjects. ^c Sigurdsson, et al⁴. Data of 3 SNP are from Swedish subjects (frequency of 563 controls in parentheses). ^d Shimane, et al¹⁸. ^e Siu, et al¹⁶. * Chi-square comparison of ethnic difference between Korean and Caucasian populations.

tioned by *CGGGGindel*, which showed the strongest association among first-group variants (*CGGGGindel*, *rs2004640*, and *rs752637*), 2 variants in the third group now appeared to have associations (*Exon6indel* and *rs10954213*). This might suggest that the genetic effects of the third-group variants were hidden by first-group variants. (2) When conditioned by the second-group SNP (*rs729302*), the effects of the first-group variants remained significant ($p = 0.03\text{--}0.003$). (3) When conditioned by a third-group variant [*Exon6indel*, which absolutely linked ($r^2 = 1$) with *rs10954213*], the effects of first and second-group variants remained significant ($p = 0.002\text{--}0.00002$). (4) Although no significant p values were obtained when conditioned on any 2 variants among each variant group simultaneously, *CGGGGindel* still remained significant ($p = 0.03$; Table 3), possibly due to the strongest association.

Altogether, conditional analysis in the Korean population suggested independent genetic effects of the 3 variant groups, although the causal polymorphism in each group could not be clearly identified.

DISCUSSION

IRF5 was originally identified as a regulator of type I interferon gene expression¹¹. Transcription factors of the IRF family play several roles in viral infection and immunostimulation, as well as cell growth regulation. *IRF5* is regulated by type I interferon, which indicates an important regulatory pathway for the controlled induction of multiple immunomodulatory genes. *IRF5* is phosphorylated in cells upon viral infections and translocates to the nucleus, which results in activation of a spectrum of interferon genes¹².

Thus, polymorphism within the *IRF5* gene may affect several cellular functions of importance for the development of an autoimmune disease such as SLE.

Genetic association studies provide a potentially powerful tool for identifying genetic variations that influence susceptibility to common diseases. However, there are many cases of associations that are not replicated afterward, which has led to skepticism about genetic epidemiology studies of complex diseases¹³⁻¹⁵. It is well known that ethnicity is one of the most important factors for evaluating genetic effects on common complex traits. Here, we demonstrate the different genetic effects of *IRF5* polymorphisms on the risk of SLE according to ethnicity.

In our study, different genetic effects of *IRF5* polymorphisms on SLE were observed in a Korean population compared with those in Caucasian populations. Three SNP (*IRF5-15-1*, *rs2070197*, and *rs10488631*), which showed strong/independent associations in Caucasian populations, had no effects (monomorphic) in the Korean population. Similar results were also reported in a Chinese population¹⁶. In addition, we have shown that the 3 groups of *IRF5* variants have independent genetic effects on the risk of SLE in a Korean population, which were different from those in Caucasian populations^{4,5}.

Association analysis of *IRF5* polymorphisms in a Korean population revealed different genetic effects of *IRF5* polymorphisms on SLE compared with those in Caucasian populations. These results indicate different genetic backgrounds of *IRF5* polymorphisms and different genetic effects on the risk of SLE according to ethnicity. The difference among ethnic groups should be considered for the clinical application of SNP.

Table 3. Conditional analysis of *IRF5* polymorphisms in Koreans with SLE and healthy controls (con).

Loci	Allele	Conditional Analysis Group	Frequency		p	<i>CGGGGindel</i>	<i>rs729302</i>	Conditional p Values			
			SLE	Con				<i>Exon6indel</i>	<i>CGGGGindel</i> and <i>rs729302</i>	<i>CGGGGindel</i> and <i>Exon6indel</i>	<i>rs729302</i> and <i>Exon6indel</i>
<i>rs729302</i>	C	Group 2	0.270	0.319	0.003	0.07	—	0.002	—	0.66	—
<i>CGGGGindel</i> *	Ins	Group 1	0.197	0.138	0.00006	—	0.003	0.00002	—	—	0.03
<i>rs2004640</i> *	T	Group 1	0.386	0.321	0.0002	0.16	0.03	0.0003	0.28	0.17	0.44
<i>rs752637</i>	C	Group 1	0.457	0.399	0.0008	0.28	0.03	0.002	0.52	0.10	0.33
<i>Exon6indel</i>	Ins	Group 3	0.503	0.495	0.73	0.02	0.13	—	0.35	—	—
<i>rs10954213</i> *	G	Group 3	0.505	0.496	0.72	0.02	0.19	NA	0.38	NA	NA
<i>rs2280714</i>	C	Group 3	0.395	0.402	0.68	0.27	0.10	0.24	0.55	0.20	0.47
Haplotype [†]											
<i>ht1 (A-Ins-G-T-Ins-G-C)</i>			0.349	0.357	0.59						
<i>ht2 (C-Ins-G-T-Del_A-T)</i>			0.157	0.205	0.0007						
<i>ht3 (A-Del-T-C-Del-A-T)</i>			0.170	0.116	0.00002						
<i>ht4 (A-Ins-T-C-Del-A-T)</i>			0.094	0.103	0.43						
<i>ht5 (A-Ins-T-C-Ins-G-T)</i>			0.077	0.066	0.37						
<i>ht6 (C-Ins-G-C-Del-A-T)</i>			0.064	0.067	0.76						

Conditional p values were estimated and permuted using WHAP software¹⁰. * Markers reported to have strong and/or independent effects on risk of SLE in Caucasian populations. † Haplotype consisting of 7 polymorphic markers. NA: No p values were available in conditional analysis because *Exon6indel* and *rs10954213* were absolutely linked ($r^2 = 1$).

ACKNOWLEDGMENT

We thank the study participants and their families who took part in the SLE cohort study of Hanyang University Hospital.

REFERENCES

1. Graham RR, Kozyrev SV, Baechler EC, et al. A common haplotype of interferon regulatory factor 5 (IRF5) regulates splicing and expression and is associated with increased risk of systemic lupus erythematosus. *Nat Genet* 2006;38:550-5.
2. Sigurdsson S, Nordmark G, Goring HH, et al. Polymorphisms in the tyrosine kinase 2 and interferon regulatory factor 5 genes are associated with systemic lupus erythematosus. *Am J Hum Genet* 2005;76:528-37.
3. Shin HD, Sung YK, Choi CB, Lee SO, Lee HW, Bae SC. Replication of the genetic effects of IFN regulatory factor 5 (IRF5) on systemic lupus erythematosus in a Korean population. *Arthritis Res Ther* 2007;9:R32.
4. Sigurdsson S, Goring HH, Kristjansdottir G, et al. Comprehensive evaluation of the genetic variants of interferon regulatory factor 5 (IRF5) reveals a novel 5 bp length polymorphism as strong risk factor for systemic lupus erythematosus. *Hum Mol Genet* 2008;17:872-81.
5. Graham RR, Kyogoku C, Sigurdsson S, et al. Three functional variants of IFN regulatory factor 5 (IRF5) define risk and protective haplotypes for human lupus. *Proc Natl Acad Sci USA* 2007;104:6758-63.
6. Hochberg MC. Updating the American College of Rheumatology revised criteria for the classification of systemic lupus erythematosus [letter]. *Arthritis Rheum* 1997;40:1725.
7. Gladman DD, Urowitz MB, Ong A, Gough J, MacKinnon A. Lack of correlation among the 3 outcomes describing SLE: disease activity, damage and quality of life. *Clin Exp Rheumatol* 1996;14:305-8.
8. Sigurdsson S, Padyukov L, Kurreeman FA, et al. Association of a haplotype in the promoter region of the interferon regulatory factor 5 gene with rheumatoid arthritis. *Arthritis Rheum* 2007;56:2202-10.
9. Livak KJ. Allelic discrimination using fluorogenic probes and the 5' nuclease assay. *Genet Anal* 1999;14:143-9.
10. Purcell S, Daly MJ, Sham PC. WHAP: haplotype-based association analysis. *Bioinformatics* 2007;23:255-6.
11. Barnes BJ, Moore PA, Pitha PM. Virus-specific activation of a novel interferon regulatory factor, IRF-5, results in the induction of distinct interferon alpha genes. *J Biol Chem* 2001;276:23382-90.
12. Barnes BJ, Richards J, Mancl M, Hanash S, Beretta L, Pitha PM. Global and distinct targets of IRF-5 and IRF-7 during innate response to viral infection. *J Biol Chem* 2004;279:45194-207.
13. Freely associating [editorial]. *Nat Genet* 1999;22:1-2.
14. Cardon LR, Bell JI. Association study designs for complex diseases. *Nat Rev Genet* 2001;2:91-9.
15. Lohmueller KE, Pearce CL, Pike M, Lander ES, Hirschhorn JN. Meta-analysis of genetic association studies supports a contribution of common variants to susceptibility to common disease. *Nat Genet* 2003;33:177-82.
16. Siu HO, Yang W, Lau CS, et al. Association of a haplotype of IRF5 gene with systemic lupus erythematosus in Chinese. *J Rheumatol* 2008;35:360-2.
17. Kozyrev SV, Alarcon-Riquelme ME. The genetics and biology of Irf5-mediated signaling in lupus. *Autoimmunity* 2007;40:591-601.
18. Shimane K, Kochi Y, Yamada R, et al. A single nucleotide polymorphism in the IRF5 promoter region is associated with susceptibility to rheumatoid arthritis in the Japanese patients. *Ann Rheum Dis* 2008 Apr 13. [Epub ahead of print]