

Single Nucleotide Polymorphism of COL6A1 in Patients with Ankylosing Spondylitis

TAE-HWAN KIM, TAE-JONG KIM, HYE-SOON LEE, WAN-SIK UHM, EUN-SOON SHIN, YOUNG-IN NA, and JAE-BUM JUN

ABSTRACT. *Objective.* To investigate the genetic association between ankylosing spondylitis (AS) and single nucleotide polymorphisms (SNP) of collagen 6A1 gene (COL6A1), the candidate gene for ossification of the posterior longitudinal ligament.

Methods. One-hundred thirty Korean patients with AS (M: 116, F: 14, age: 29.0 ± 4.6) and 130 age- and sex-matched healthy subjects were recruited. The SNP of G365G, IVS15+39 C/T, IVS21+18 A/C by Snap shot assay and the SNP of IVS32-29T/C, IVS33+15G/A, IVS33+20A/G, and IVS33+55A/G by direct sequencing were genotyped and analyzed. Bonferroni correction was applied to multiple comparisons.

Results. The observed allelic frequencies for these SNP met Hardy-Weinberg equilibrium in all AS and controls. We also found an additional 2 SNP (R783Q and IVS33+88C/T) during direct sequencing. Therefore, a total of 9 SNP were analyzed in this study. There were no significant associations of allelic and genotype variations between AS and controls. The presence of uveitis was marginally associated with a haplotype (CC in G365G + IVS15+39 C/T). The variation of allele or haplotype of COL6A1 is not significantly associated with "more ossified disease."

Conclusion. Because the genetic variations of COL6A1 could not be correlated with the occurrence of AS in Koreans, we conclude that despite common clinical features, AS and ossification of posterior longitudinal ligament are not genetically related, and the hyperostotic condition seen in the 2 diseases might be regulated differently. Further SNP of COL6A1 were not related to radiographic progression of AS. However, we found that the occurrence of uveitis might be related to the genetic variations of COL6A1 in patients with AS. (First Release July 15 2008; J Rheumatol 2008;35:1849–52)

Key Indexing Terms:

ANKYLOSING SPONDYLITIS COL6A1 SINGLE NUCLEOTIDE POLYMORPHISM
OSSIFICATION OF POSTERIOR LONGITUDINAL LIGAMENT

Ankylosing spondylitis (AS) is characterized by inflammatory back pain, peripheral arthritis, uveitis, and an association with HLA-B27. However, the most distinct feature is spinal involvement, which includes syndesmophytosis and ectopic ossification of spinal ligaments. AS shares common features of heterotopic ossification with other relatively common diseases characterized by calcification of spinal ligaments, such as ossification of posterior longitudinal ligament (OPLL)¹. AS is sporadically complicated with OPLL, and recently we reported that the frequency of OPLL in 544 Korean patients with AS was 3.5%². A single nucleotide polymorphism (SNP) of the collagen 6A1 gene (COL6A1)

has been revealed to be strongly associated with OPLL³ and diffuse idiopathic skeletal hyperostosis (DISH)⁴.

We hypothesized that these SNP of COL6A1 were implicated in development of AS and ectopic ossification in spinal ligaments.

MATERIALS AND METHODS

Subjects. One-hundred thirty AS patients, who met modified New York criteria⁵, were consecutively recruited (116 men, mean age: 29.0 ± 4.6 yrs). One-hundred thirty age- and sex-matched controls were recruited from health professionals and college students (mean age: 28.6 ± 3.4 yrs).

Seventy-seven patients with ≥ 10 years of disease duration and available cervical spine radiographs were selected to examine the relationship between radiographic progression and COL6A1 SNP. Lateral views were scored using the Bath Ankylosing Spondylitis Radiology Index (BASRI)⁶ and were examined for the existence of OPLL⁷ and syndesmophytosis. We classified AS patients as having "more ossified disease" if they had a BASRI score of 3 or 4, or had OPLL of any type. Our research work was approved by the Institutional Review Board.

Polymerase chain reaction (PCR) and genotyping. Genomic DNA was prepared from peripheral blood with written consent, and PCR was performed by a standard method. Genotyping was done with a single base primer extension assay (Table 1) using SNaPSHOT assay, and direct sequencing using electrophoresis. Results were analyzed using Gene Mapper software (Applied Biosystems, Foster City, CA, USA).

We investigated 7 SNP that are reportedly related to OPLL or DISH. An

From the Hospital for Rheumatic Diseases, Hanyang University; Bioinformatics Team, DNA Link, Inc; and Institute of Rheumatism, Hanyang University, Seoul, Republic of Korea.

T-H. Kim, MD, PhD; H-S. Lee, MD, PhD; W-S. Uhm, MD, PhD, Hospital for Rheumatic Diseases, Hanyang University; T-J. Kim, MD, Department of Rheumatology, Chonnam National University Hospital, Gwangju; E-S. Shin, PhD, Bioinformatics Team, DNA Link, Inc; Y-I. Na, MS, Institute of Rheumatism; J-B. Jun, MD, PhD, Hospital for Rheumatic Diseases, Hanyang University.

Address reprint requests to Dr. J-B. Jun, Hospital for Rheumatic Diseases, Hanyang University, 17 Haengdang-Dong, Sungdong-Gu, Seoul 133-792, Republic of Korea.

Accepted for publication April 8, 2008.

Personal non-commercial use only. The Journal of Rheumatology Copyright © 2008. All rights reserved.

Table 1. Primer sequences of SNP of COL6A1 gene.

Character	Method	Strand	Primer	Sequence (5'→3')
G365G	SnaPshot	Reverse	F	TTT TCT CCA GGG CAT TCA
			R	AGG GGC CGC AAG TCA CTC
			G	CCT TTT CTC CTT TCA GTC CAA AGG C
IVS15+39 C/T	SnaPshot	Forward	F	TTC TCC AGG GCA TTC AAG G
			R	GAC TTG CTT CCA GGA GAG G
			G	GGC CCC TGG AGG ACC AGG GCC TTC A
IVS21+18 A/C	SnaPshot	Forward	F	TAA AGG CTA CCG AGG CGA
			R	AGA CAG TGG GTT CTG GAT CC
			G	GGT CCG AGG TGA GTC CCA CTC CCC A
IVS32-29 T/C	Sequencing	Reverse	F	ATG CAG AGC CAT CAA GAG
R783Q	Sequencing	Reverse	R	CCC TGG AGG TAA CAG TGA
			F	ATG CAG AGC CAT CAA GAG
IVS33+15 G/A	Sequencing	Reverse	R	CCC TGG AGG TAA CAG TGA
			F	ATG CAG AGC CAT CAA GAG
IVS33+20 A/G	Sequencing	Reverse	R	CCC TGG AGG TAA CAG TGA
			F	ATG CAG AGC CAT CAA GAG
IVS33+55 A/G	Sequencing	Forward	R	CCC TGG AGG TAA CAG TGA
			F	ATG CAG AGC CAT CAA GAG
IVS33+88 C/T	Sequencing	Reverse	R	CCC TGG AGG TAA CAG TGA
			F	ATG CAG AGC CAT CAA GAG

F: PCR forward primer, R: PCR reverse primer, G: genotyping primer.

additional 2 SNP (SNP No. 5 and 9) were detected during direct sequencing. Therefore, 9 SNP (Table 2) were chosen for genotyping.

Haplotype estimation and defining haplotype blocks. Haplotypes and haplotype blocks were defined by Haploview. There were 2 blocks: the first included the first 2 SNP and the second included the others. Seven haplotypes with a frequency $\geq 5\%$ were submitted to further analysis.

Statistical analysis. Differences in frequency were determined using a 2-

tailed test or multiple logistic regression analysis. Odds ratios were calculated with 95% confidence interval for the relative risk for the development of AS and its clinical manifestations related to the specific genotypes by logistic regression analysis, including age and sex. Haplotype frequencies were calculated by using an expectation-maximization algorithm. Bonferroni correction was applied to multiple comparisons. Statistical analyses were performed with SAS (version 9-1-3, SAS Institute, Cary, NC, USA).

Table 2. Individual SNP and haplotype studied.

SNP	Character	Location	NCBI dbSNP ID	Nucleotide Substitution
1	G365G	Exon 15 (+39)	rs1980982	T/C
2	IVS15+39 C/T	Intron 15 (+39)	Non-dbSNP	C/T
3	IVS21+18 A/C	Intron 21 (+18)	rs2276254	A/C
4	IVS32-29 T/C	Intron 32 (-29)	Non-dbSNP	T/C
5*	R783Q	Exon 33 (+98)	Non-dbSNP	G/A
6	IVS33+15 G/A	Intron 33 (+15)	rs2236485	G/A
7	IVS33+20 A/G	Intron 33 (+20)	rs2236486	A/G
8	IVS33+55 A/G	Intron 33 (+55)	rs2236487	A/G
9*	IVS33+88 C/T	Intron 33 (+88)	rs2236488	C/T
Block	Haplotype	Selected SNP	Alleles	
Haplotype block 1	Haplotype 1	SNP 1 + 2	TC	
	Haplotype 2		CC	
	Haplotype 3		CT	
Haplotype block 2	Haplotype 4	SNP 3+4+6+7+8	ATGAA	
	Haplotype 5		CCGGG	
	Haplotype 6		CCAGG	
	Haplotype 7		ATGGG	

SNP 5 and 9: detected by chance during direct sequencing.

RESULTS

Allele and genotype frequency. All allele and genotype frequencies met Hardy-Weinberg equilibrium. There were no significant associations of allelic and genotype variations between AS and controls. Uveitis was associated with SNP No. 1 ($p = 0.03$, OR = 3.222, 95% CI = 1.119–9.278, recessive model) and SNP No. 6 ($p = 0.027$, OR = 2.654, 95% CI = 1.114–6.326, co-dominant model), but these associations did not show statistical significance after Bonferroni correction. No SNP was associated with peripheral arthritis and HLA-B27.

Haplotype analysis. Within the block, no haplotype frequencies in all haplotypes except Haplotype 1 showed statistical differences. There was a significant difference of haplotype distribution in patients according to presence of uveitis after Bonferroni correction for 3 haplotypes and 3 models (Table 3). Haplotype 1 occurred at a lower frequency in patients with uveitis compared to patients without uveitis ($p = 0.032$ before Bonferroni correction, OR = 0.315, 95% CI = 0.109–0.906, dominant model), but no significance remained after Bonferroni correction. In contrast, Haplotype 2 showed at a higher frequency in patients with uveitis ($p = 0.04978$ after Bonferroni correction, OR = 3.331, 95% CI = 1.001–11.086, co-dominant model). No haplotype had a significant correlation with peripheral arthritis.

“More ossified disease.” Among 77 patients, 27 patients were classified as having more ossified disease. There was no significant variation of alleles/genotypes or haplotypes of COL6A1 between the control disease and the more ossified disease (data not shown).

DISCUSSION

We found no significant associations of allelic and genotype variations of COL6A1 between AS and controls. Thus, COL6A1 may not have a role in development of spinal manifestations of AS. Especially SNP No. 4 (association with OPLL, $p = 0.000003$), and SNP No. 4, 6, and 3 (association with DISH, $p = 0.0005$, $p = 0.022$, $p = 0.024$, respectively) showed no genetic significance^{3,4}. Further, OPLL-specific haplotype³, constructed with 3 SNP (No. 3, 4, and 7), did not show any association with the development of AS.

Interestingly, uveitis was marginally associated with Haplotype 2. Although there have been some reports on the

association between cytokine and chemokine gene polymorphism and uveitis^{8,9}, we do not have an explanation for association of COL6A1 with uveitis.

COL6A1 region has been reported to be associated with congenital heart defects of Down syndrome¹⁰, muscle disorders such as Ullrich muscular dystrophy¹¹, Bethlem myopathy¹², and fibronectin organization¹³. The functional role of SNP of COL6A1 is still unknown, but several authors have suggested that these genetic variations of collagen genes may contribute to the membranous or enchondral ossification of the spinal ligaments^{3,14}.

In conclusion, although some SNP of COL6A1 have been reported to be strongly related to OPLL and DISH, COL6A1 gene may not be the gene responsible for development of AS. Further, SNP of COL6A1 did not seem to influence radiographic progression of AS. However, the association of Haplotype 2 (CC) of COL6A1 with uveitis in AS patients requires further investigation; the power of our study is not strong due to limited sample size and absence of non-AS uveitis controls.

REFERENCES

1. Mori K, Chano T, Ikeda T, et al. Decrease in serum nucleotide pyrophosphatase activity in ankylosing spondylitis. *Rheumatology* 2003;42:62-5.
2. Kim TJ, Kim TH, Jun JB, Joo KB, Uhm WS. Prevalence of ossification of posterior longitudinal ligament in patients with ankylosing spondylitis. *J Rheumatol* 2007;34:2460-2.
3. Tanaka T, Ikari K, Furushima K, et al. Genomewide linkage and linkage disequilibrium analyses identify COL6A1, on chromosome 21, as the locus for ossification of the posterior longitudinal ligament of the spine. *Am J Hum Genet* 2003;73:812-22.
4. Tsukahara S, Miyazawa N, Akagawa H, et al. COL6A1, the candidate gene for ossification of the posterior longitudinal ligament, is associated with diffuse idiopathic skeletal hyperostosis in Japanese. *Spine* 2005;30:2321-4.
5. van der Linden S, Valkenburg HA, Cats A. Evaluation of diagnostic criteria for ankylosing spondylitis: a proposal for modification of the New York criteria. *Arthritis Rheum* 1984;27:361-8.
6. MacKay K, Mack C, Brophy S, Calin A. The Bath ankylosing spondylitis radiology index (BASRI): a new, validated approach to disease assessment. *Arthritis Rheum* 1998;41:2263-70.
7. Tsuyama N. Ossification of the posterior longitudinal ligament of the spine. *Clin Orthop Relat Res* 1984;71-84.
8. El-Shabrawi Y, Wegscheider BJ, Weger M, et al. Polymorphisms within the tumor necrosis factor-alpha promoter region in patients with HLA-B27-associated uveitis: association with susceptibility and clinical manifestations. *Ophthalmology* 2006;113:695-700.
9. Chen Y, Vaughan RW, Kondeatis E, et al. Chemokine gene

Table 3. Haplotype analysis of SNP in COL6A1 gene in patients with AS according to uveitis.

	Dominant		Recessive		Co-dominant	
	p	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)
Haplotype 2						
Uveitis	0.075	3.280	1.000	∞	0.04978	3.331
No uveitis		(0.941, 11.439)		(0.000, ∞)		(1.001, 11.086)

∞: infinite. p value: after Bonferroni correction for 9 tests (3 haplotypes and 3 models).

- polymorphisms associate with gender in patients with uveitis. *Tissue Antigens* 2004;63:41-5.
10. Davies GE, Howard CM, Farrer MJ, et al. Genetic variation in the COL6A1 region is associated with congenital heart defects in trisomy 21 (Down's syndrome) (Pt 3). *Ann Hum Genet* 1995;59:253-69.
 11. Camacho Vanegas O, Bertini E, Zhang RZ, et al. Ullrich scleroatonic muscular dystrophy is caused by recessive mutations in collagen type VI. *Proc Natl Acad Sci USA* 2001;98:7516-21.
 12. Jobsis GJ, Keizers H, Vreijling JP, et al. Type VI collagen mutations in Bethlem myopathy, an autosomal dominant myopathy with contractures. *Nat Genet* 1996;14:113-5.
 13. Sabatelli P, Bonaldo P, Lattanzi G, et al. Collagen VI deficiency affects the organization of fibronectin in the extracellular matrix of cultured fibroblasts. *Matrix Biol* 2001;20:475-86.
 14. Maeda S, Ishidou Y, Koga H, et al. Functional impact of human collagen alpha2(XI) gene polymorphism in pathogenesis of ossification of the posterior longitudinal ligament of the spine. *J Bone Miner Res* 2001;16:948-57.