

Limited Diagnostic Value of Unilateral Sacroiliitis in Scintigraphy in Assessing Axial Spondyloarthritis

IN-HO SONG, HENNING BRANDT, MARTIN RUDWALEIT, and JOACHIM SIEPER

ABSTRACT. Objective. To assess the diagnostic value for axial spondyloarthritis (SpA) of unilateral sacroiliitis in scintigraphy in daily clinical practice.

Methods. In 207 patients with chronic back pain, the diagnostic value of scintigraphy was assessed retrospectively. The diagnosis made by the rheumatologist (axial SpA vs no axial SpA) was the standard.

Results. Sensitivities of scintigraphy for any (unilateral or bilateral), bilateral, and isolated unilateral sacroiliitis were 64.9%, 40.2%, and 24.7%, respectively. Respective specificities were 50.5%, 57.7%, and 92.8%, resulting in likelihood ratios of 1.3, 1.0, and 3.4.

Conclusion. Scintigraphy of the sacroiliac joints is of limited value for the diagnosis of axial SpA. Unilateral compared to bilateral sacroiliitis is slightly superior, but is associated with a low sensitivity. (First Release April 15 2010; J Rheumatol 2010;37:1200–2; doi:10.3899/jrheum.091216)

Key Indexing Terms:

SCINTIGRAPHY
DIAGNOSIS

ANKYLOSING SPONDYLITIS
LIKELIHOOD RATIO

SACROILIITIS
SENSITIVITY

A recent systematic literature search showed that scintigraphy of the sacroiliac joints (SIJ) is of limited value for the diagnosis of axial spondyloarthritis (SpA)¹. In daily clinical practice, an isolated unilateral sacroiliitis in bone scintigraphy is often regarded to be of better diagnostic value because the negative side acts as an internal control, while in the case of bilateral sacroiliitis the definition of positive/negative is more difficult. We retrospectively analyzed a cohort of patients who were referred because of suspected SpA to a specialized tertiary rheumatology center with special interest in SpA, and considered whether unilateral sacroiliitis would have a higher diagnostic value compared to bilateral sacroiliitis.

MATERIALS AND METHODS

Patients. We retrospectively analyzed patients with chronic low back pain who were seen in our back pain clinic² and who had undergone a bone scintigraphy examination for the evaluation of sacroiliitis. A diagnosis of axial SpA was made by experienced rheumatologists according to the modified New York criteria for ankylosing spondylitis (AS)³ or for nonradiographic axial SpA^{2,4,6}. Importantly, for the diagnosis of axial SpA, scintigraphy findings were not taken into account. Controls were defined as those patients in whom the expert excluded the diagnosis of axial SpA.

Scintigraphy. Bone scintigraphy findings were classified into unilateral or bilateral sacroiliitis, and suspicion for or exclusion of sacroiliitis. Where

available, we collected information about the exact value of the SIJ-sacrum-index (SIJ-S-index), which quantifies the uptake of the radio-tracer in the SIJ compared to the sacrum as a reference point. The decision about whether a scintigraphy result showed active sacroiliitis was based on the judgment of the radiologist who had performed the scintigraphy examination.

Statistics. Sensitivity, specificity, and likelihood ratio (LR) of scintigraphy for the diagnosis of axial SpA were calculated.

RESULTS

Out of 1202 patients who have been referred to our clinic for the evaluation of back pain, in 207 cases (17.2%) a scintigraphy examination of the SIJ had been performed before referral for the evaluation of sacroiliitis. Out of these 207 patients, we classified 97 as having axial SpA (56 as AS and 41 as nonradiographic axial SpA), 13 as suspicious for axial SpA, and 97 as controls^{2,4,5}. Demographic and clinical data are shown in Table 1.

Discriminative property of scintigraphy of the SIJ between patients and controls. All 207 scintigraphy examinations had been performed using ^{99m}Tc technetium-labeled radiotracers. The mean radiation dose was 568.9 mBq (SD 94.8). In 50.2% (104/207) of scintigraphy examinations, absolute values for the SIJ-S-index were provided.

First, we analyzed whether there was a difference between patients with axial SpA and controls regarding tracer uptake, independent of bilateral or unilateral sacroiliitis. There was no statistically significant difference between patients with axial SpA (mean uptake left SIJ/right SIJ 1.48/1.46; mean cutoff 1.33) and controls (mean uptake left SIJ/right SIJ 1.48/1.41; mean cutoff 1.30).

Diagnostic properties of scintigraphy for sacroiliitis. We next split scintigraphy findings into bilateral and unilateral

From the Charité Medical University, Campus Benjamin Franklin, Medical Clinic I, Rheumatology, Berlin, Germany.

I.H. Song, MD; H.C. Brandt, MD; M. Rudwaleit, MD; J. Sieper, MD, Charité Medical University.

Address correspondence to Dr. J. Sieper, Charité, Campus Benjamin Franklin, Medical Clinic I, Rheumatology, Hindenburgdamm 30, 12200 Berlin, Germany. E-mail: joachim.sieper@charite.de
Accepted for publication February 4, 2010.

Personal non-commercial use only. The Journal of Rheumatology Copyright © 2010. All rights reserved.

Table 1. Demographic and clinical data for the different patient groups who underwent a scintigraphy examination for the evaluation of sacroiliitis. In the patient group with axial SpA, patients with AS and patients with nonradiographic axial SpA are pooled.

Characteristic	AS	Nonradiographic Axial SpA	Axial SpA (pooled AS plus nonradiographic SpA)	Controls
No. of patients	56	41	97	97
Men, %	67.9	46.3	58.8	44.4
Mean age, yrs (SD)	39.3 (12.8)	35.1 (10.2)	37.5 (11.9)	42.8 (13.2)
Mean disease duration, yrs (SD)	11.6 (9.2)	6.7 (6.5)	9.6 (8.5)	9.6 (10.1)
% HLA-B27-positive	85.7	75.0	80.4	34.0*
BASDAI, mean (SD)	4.4 (2.3)	4.2 (2.3)	4.3 (2.3)	4.2 (1.7)
BASFI, mean (SD)	3.3 (2.3)	3.3 (2.3)	3.3 (2.4)	3.3 (2.2)

* The high percentage of HLA-B27-positive controls occurred because HLA-B27 was one of the referral measurements².




Figure 1. Sensitivity, specificity, and likelihood ratio (LR) of bone scintigraphy showing any sacroiliitis, bilateral sacroiliitis, or isolated unilateral sacroiliitis. Data are shown for the whole group of axial spondyloarthritis (SpA, n = 97) and non-SpA controls (n = 97). Specificities are calculated from 97 controls.

sacroiliitis independently of quantification. Thus, for the following analysis, qualitative and quantitative scintigraphy findings were analyzed for 97 patients with axial SpA and 97 controls. Only positive or negative findings were counted, according to the radiologist who had performed the scintigraphy examination.

Among 97 patients with axial SpA, scintigraphy showed any sacroiliitis (bilateral or unilateral) in 64.9%, bilateral sacroiliitis in 40.2%, and unilateral sacroiliitis in 24.7%. Based on the analysis of the 97 controls, specificities of 50.5%, 57.7%, and 92.8% and LR of 1.3, 1.0, and 3.4, respectively, were found (Figure 1 and group A in Table 2).

In the 2 subgroups with established AS (n = 56, Group A1 in Table 2) compared to nonradiographic axial SpA (n = 41, Group A2 in Table 2), as well as in the AS subgroups with sacroiliitis grade 2/3 (Group A1a in Table 2) and sacroiliitis grade 4 (Group A1b in Table 2), the value of scintigraphy for the diagnosis of axial SpA was very similar compared to the pooled group of axial SpA. In the subgroup of patients with axial SpA in whom absolute values of the

SIJ-S-index were given (26 patients with AS, 21 patients with nonradiographic axial SpA, and 48 controls, Group B in Table 2), it could be shown that quantitative scintigraphy was not superior to qualitative scintigraphy.

DISCUSSION

Our retrospective analysis shows that scintigraphy is still used in a proportion of patients for the evaluation of sacroiliitis. About 17% of referred patients with back pain had undergone a scintigraphic examination of the SIJ. Further, our study confirms data of a recent systematic literature search that showed that scintigraphy of the SIJ is at best of limited value for the diagnosis of axial SpA¹. This is especially true for bilateral sacroiliitis, which showed only a +LR 1.0 in our study, meaning that it is of no use at all. Unilateral sacroiliitis as shown by scintigraphy showed a slightly higher LR of 3.4. At the same time, unilateral sacroiliitis on scintigraphy was associated with a very low sensitivity of about 25%. Further, an LR of 3.4 of scintigraphic unilateral sacroiliitis is much lower than an assumed LR of around 10 for magnetic resonance imaging (MRI)⁵.

Our findings are in concordance with the few scintigraphy studies that have so far reported on unilateral sacroiliitis (6, including 104 patients with axial SpA and 36 controls)⁷⁻¹². Sensitivities of unilateral sacroiliitis in these studies were low, ranging from 6.3%¹⁰ to 33.3%¹², while specificities were high, ranging between 89.5%⁸ and 94.1%⁷, resulting in a pooled LR of 2.1.

In the diagnostic approach in our center, scintigraphy results are not included because there has been no proven value for this method for the diagnosis of axial SpA. The optimal study design would be a prospective study with patients with SpA and patients without SpA who are undergoing scintigraphy of the SIJ. Looking at all available data on the limitation of scintigraphy and the superiority of MRI, it is questionable whether such a study is still feasible.

The diagnostic value of scintigraphy of the SIJ might be better in clear cases of unilateral sacroiliitis shown by scintigraphy that correlates with the patient's symptoms of

Table 2. Sensitivity, specificity, and likelihood ratio (LR) of scintigraphy imaging for the diagnosis of axial spondyloarthritis (SpA). Scintigraphy examinations of the sacroiliac joints were performed in 207 patients. Of them, 97 patients were classified with axial SpA, 13 patients with suspicion for axial SpA, and 97 as controls.

Diagnosis	Unilateral or Bilateral Sacroiliitis, or Both	Sensitivity, %	Specificity, %	LR
Group A (total group):	Any	64.9	50.5	1.3
axial SpA (AS plus	Bilateral	40.2	57.7	1.0
Nonradiographic axial SpA, n = 97)*	Unilateral	24.7	92.8	3.4
Group A1: AS (n = 56)	Any	67.9	50.5	1.4
	Bilateral	42.9	57.7	1.0
	Unilateral	25.0	92.8	3.5
Group A1a: AS subgroup,	Any	74.4	50.5	1.5
grade 2/3 sacroiliitis (n = 43)**	Bilateral	46.5	57.7	1.1
	Unilateral	27.9	92.8	3.9
Group A1b: AS subgroup,	Any	12.5	50.5	0.3
grade 4 sacroiliitis (n = 8)**	Bilateral	12.5	57.7	0.3
	Unilateral	0	92.8	0
Group A2:	Any	61.0	50.5	1.2
Nonradiographic axial	Bilateral	36.6	57.7	0.9
SpA (n = 41)	Unilateral	24.4	92.8	3.4
Group B: Axial SpA (AS	Any	74.5	27.1	1.0
plus nonradiographic axial SpA,	Bilateral	46.8	41.7	0.8
n = 47), only quantitative sacroiliitis***	Unilateral	27.7	85.4	1.9

* Group A consists of all patients with axial SpA (pooled data) including 56 patients with AS (Group A1) and 41 patients with nonradiographic axial SpA (Group A2). ** Group A1 consists of all patients with AS according to the modified New York criteria, including patients with sacroiliitis grade 2/3 on conventional radiography (Group A1a) and patients with total ankylosis on conventional radiography (Group A1b). In 5 patients with AS, the exact grading of sacroiliitis was missing. *** Subgroup of patients with axial SpA, in whom absolute values of the sacroiliac joint sacrum index were given (26 patients with AS, 21 patients with nonradiographic axial SpA, and 48 controls).

unilateral buttock pain. This information was not available for our study, and such an analysis has not been performed in the literature.

Our analysis shows that scintigraphy of the SIJ is of quite limited value for the diagnosis of axial SpA. Unilateral sacroiliitis is slightly superior compared to scintigraphy showing a bilateral sacroiliitis, but is associated with a low sensitivity.

REFERENCES

1. Song IH, Carrasco-Fernández J, Rudwaleit M, Sieper J. The diagnostic value of scintigraphy in assessing sacroiliitis in ankylosing spondylitis: a systematic literature research. *Ann Rheum Dis* 2008;67:1535-40.
2. Brandt HC, Spiller I, Song IH, Vahldiek JL, Rudwaleit M, Sieper J. Performance of referral recommendations in patients with chronic back pain and suspected axial spondyloarthritis. *Ann Rheum Dis* 2007;66:1479-84.
3. van der Linden S, Valkenburg HA, Cats A. Evaluation of diagnostic criteria for ankylosing spondylitis. A proposal for modification of the New York criteria. *Arthritis Rheum* 1984;27:361-8.
4. Rudwaleit M, Khan MA, Sieper J. The challenge of diagnosis and classification in early ankylosing spondylitis: do we need new criteria? *Arthritis Rheum* 2005;52:1000-8.
5. Rudwaleit M, van der Heijde D, Khan MA, Braun J, Sieper J. How to diagnose axial spondyloarthritis early. *Ann Rheum Dis* 2004;63:535-43.
6. Sieper J, van der Heijde D, Landewe R, Brandt J, Burgos-Vagas R, Collantes-Estevez E, et al. New criteria for inflammatory back pain in patients with chronic back pain: a real patient exercise by experts from the Assessment of SpondyloArthritis international Society (ASAS). *Ann Rheum Dis* 2009;68:784-8.
7. Berghs H, Remans J, Driessens L, Kiebooms L, Polderman J. Diagnostic value of sacroiliac joint scintigraphy with 99m technetium pyrophosphate in sacroiliitis. *Ann Rheum Dis* 1978;37:190-4.
8. Van Laere M, Veys EM, Mielants H. Strontium 87m scanning of the sacroiliac joints in ankylosing spondylitis. *Ann Rheum Dis* 1972;31:201-6.
9. Szanto E, Axelsson B, Lindvall N. Detection of sacro-ilitis. A comparison between profile scanning with the use of 99m Tc pertechnetate and scintigraphy with 99m Tc diphosphonate. *Scand J Rheumatol* 1977;6:129-36.
10. Jee WH, McCauley TR, Lee SH, Kim SH, Im SA, Ha KY. Sacroiliitis in patients with ankylosing spondylitis: association of MR findings with disease activity. *Magn Reson Imaging* 2004;22:245-50.
11. Yildiz A, Gungor F, Tuncer T, Karayalcin B. The evaluation of sacroiliitis using 99mTc-nanocolloid and 99mTc-MDP scintigraphy. *Nucl Med Commun* 2001;22:785-94.
12. Peh WC, Ho WY, Luk KD. Applications of bone scintigraphy in ankylosing spondylitis. *Clin Imaging* 1997;21:54-62.