

Association of IgA Anti- β_2 Glycoprotein I with Clinical and Laboratory Manifestations of Systemic Lupus Erythematosus

TARANEH MEHRANI and MICHELLE PETRI

ABSTRACT. Objective. IgA isotypes of anticardiolipin and anti- β_2 glycoprotein I (anti- β_2 -GPI) are omitted from the revised antiphospholipid syndrome (APS) classification criteria. Multiple studies have found a high prevalence of IgA anti- β_2 -GPI in systemic lupus erythematosus (SLE). We determined the frequency and associations of IgA anti- β_2 -GPI in a cohort of patients with SLE.

Methods. Anti- β_2 -GPI was measured in 796 patients with SLE (93% women, 53% white, 38% African American, mean age 45 yrs). IgA anti- β_2 -GPI (> 20 phospholipid units) was found in 20%. Using a cohort database, associations with cumulative thrombotic and other manifestations were determined.

Results. Of patients with SLE who demonstrated IgA anti- β_2 -GPI positivity, about 6% had transient ischemic attack ($p = 0.070$), 4% had superficial thrombophlebitis ($p = 0.647$), 20% had deep venous thrombosis ($p = 0.003$), 4% had other venous thrombosis ($p = 0.827$), 12% had stroke ($p = 0.050$), and 1% had myocardial infarction ($p = 0.397$).

Conclusion. IgG anti- β_2 -GPI has the strongest association with thrombosis in SLE. However, IgA anti- β_2 -GPI was more strongly associated with deep venous thrombosis and with stroke than was IgM. These results indicate that assessment of IgA anti- β_2 -GPI is associated with thrombosis in SLE, and that the classification criteria for APS should be revised to include IgA anti- β_2 -GPI in patients with SLE. (First Release Oct 15 2010; J Rheumatol 2011;38:64–8; doi:10.3899/jrheum.100568)

Key Indexing Terms:

ANTI- β_2 -GLYCOPROTEIN I

SYSTEMIC LUPUS ERYTHEMATOSUS

ANTI-PHOSPHOLIPID ANTIBODIES

Studies have recognized that the IgA isotype of anti- β_2 -glycoprotein I (anti- β_2 -GPI) is very frequent in patients with systemic lupus erythematosus (SLE)^{1,2,3,4}. Fanopoulos, *et al* measured anticardiolipin antibodies (aCL) and anti- β_2 -GPI by ELISA in 48 patients with SLE and 100 control subjects, and found IgA anti- β_2 -GPI in 58% of patients with SLE versus in 2% of controls¹. Moreover, compared to aCL and other isotypes of anti- β_2 -GPI, the highest antibody levels were seen in IgA anti- β_2 -GPI¹. Tsutsumi and colleagues reported a 25% prevalence of IgA anti- β_2 -GPI in 124 Japanese patients with SLE². The presence of IgA anti- β_2 -GPI was significantly associated with a history of

thrombosis². Lee, *et al* analyzed the isotypes of anti- β_2 -GPI in 270 patients with SLE⁵. The prevalence of IgG, IgM, and IgA anti- β_2 -GPI was 38.1%, 13.7%, and 34.8%, respectively⁵. Finally, Lakos, *et al* evaluated 70 patients [30 with antiphospholipid syndrome (APS) secondary to SLE, 3 with primary APS, 37 with SLE without APS] and found anti- β_2 -GPI antibodies of the IgA isotype present in 59.3% of patients with APS⁶.

Studies have shown a high prevalence of IgA anti- β_2 -GPI in African Americans with SLE^{3,7}. Cucurull, *et al* reported that IgA anti- β_2 -GPI was the most prevalent isotype (19%) among African American patients with SLE³. In another analysis of 418 consecutive patients with SLE or APS by our group, IgM (28.9%) and IgA (22.2%) were the most frequent isotypes of anti- β_2 -GPI⁴. IgA was the most prevalent isotype (8.9%) of anti- β_2 -GPI in African Americans⁴.

Our aim was to determine the clinical significance of IgA anti- β_2 -GPI compared with the other isotypes in a large cohort of patients with SLE.

MATERIALS AND METHODS

Patients. A group of 796 patients with SLE (733 women; median age 45.0 yrs, range 19.1–85.7) were tested for all isotypes (IgG, IgM, and IgA) of anti- β_2 -GPI and aCL. There were 422 (53.5%) whites, 302 (38.3%) African Americans, 26 (3.3%) Hispanics, and 27 (3.4%) Asians. All the patients

From the Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland, and the University of California, Irvine, California, USA.

Supported by the National Institutes of Arthritis and Musculoskeletal and Skin Diseases (RO1 AR043727) and the Johns Hopkins University Institute for Clinical and Translational Research, Clinical Research Units.

T. Mehrani, MD, Rheumatology Fellow, University of California, Irvine; M. Petri, MD, MPH, Professor, Department of Medicine and Director, Johns Hopkins Lupus Center, Johns Hopkins University School of Medicine.

Address correspondence to Dr. M. Petri, Johns Hopkins University School of Medicine, 1830 East Monument Street, Suite 7500, Baltimore, MD 21205, USA. E-mail: mpetri@jhmi.edu

Accepted for publication August 17, 2010.

classified as having SLE met 4 or more of the 1982 revised American College of Rheumatology (ACR) criteria^{8,9}.

Protocol. The patients in the study were all part of the Johns Hopkins Hospital Lupus Cohort in Baltimore, which has been well described^{10,11}. The study was approved by the Johns Hopkins University School of Medicine Institutional Review Board. All patients gave written informed consent. Patients were followed quarterly, or more often if required for disease activity or complications. Disease activity was measured with the Physician Global Assessment, Lupus Activity Index, and the Safety of Estrogens in Lupus Erythematosus National Assessment–Systemic Lupus Erythematosus Disease Activity Index instrument score^{12,13}, and damage was recorded using the Systemic Lupus International Collaborating Clinics/ACR Damage Index¹⁴.

Technique and statistics. Both aCL and anti- β_2 -GPI were measured by ELISA (Inova Diagnostics, San Diego, CA, USA). Statistical analyses were done using chi-squared and Fisher's exact tests (SAS Institute, Cary, NC, USA). A p value ≤ 0.05 was considered statistically significant. The strength of the association was measured by calculating the OR with its 95% CI.

RESULTS

Prevalence of anti- β_2 -GPI. Anti- β_2 -GPI of at least 1 isotype were present in 330 patients with SLE (41.5%). IgG, IgM, and IgA anti- β_2 -GPI were positive in 43 (5.4%), 126 (15.8%), and 161 (20.2%) patients, respectively (Table 1). The prevalence of isolated IgA anti- β_2 -GPI was 104 (13.1%). Anti- β_2 -GPI was more common in whites (57.6%) than in African Americans (37.2%). IgA anti- β_2 -GPI was the most frequent isotype detected in African Americans, whites, and Hispanics (Table 2).

Prevalence of aCL. A total of 480 out of 783 patients (61.3%) with SLE were positive for aCL. However, only

196 patients (25%) had both aCL and anti- β_2 -GPI, and 19 (2.4%) had isolated IgA aCL.

Thrombosis in anti- β_2 -GPI positives and negatives. A history of venous thrombosis (17.51% of total patients with SLE) was significantly associated with IgA anti- β_2 -GPI positivity (24.2%) as compared to the IgA anti- β_2 -GPI-negative (15.8%) isotype (OR 1.7, 95% CI 1.12–2.59; p = 0.0097). There was no statistical difference in IgA anti- β_2 -GPI with arterial thrombosis (p = 0.08). IgG anti- β_2 -GPI had a stronger association with venous thrombosis (OR 3.03, 95% CI 1.58–5.79; p = 0.0014) and arterial thrombosis (OR 2.18, 95% CI 1.04–4.60; p = 0.0553) than the IgA isotype. There was no significant association of IgM anti- β_2 -GPI with arterial or venous thrombosis in patients with SLE (Table 3).

Relationship of IgA anti- β_2 -GPI or aCL to pregnancy loss, livedo, and thrombocytopenia. Of patients with SLE in the cohort who had ever been pregnant (n = 569), 231 (40.6%) had a history of pregnancy loss. Neither aCL nor anti- β_2 -GPI were associated with pregnancy loss. There was a 27.5% prevalence of livedo reticularis in our cohort. No association was found between IgA anti- β_2 -GPI and a history of livedo reticularis. However, IgA anti- β_2 -GPI positivity was associated with a history of thrombocytopenia (OR 1.6, 95% CI 1.06–2.33, p = 0.0172).

Relationship of anti- β_2 -GPI IgG, IgM, and IgA with other serological markers. The lupus anticoagulant by dilute Russell viper venom time was associated with anti- β_2 -GPI of any isotype (p < 0.0001 for all isotypes). IgG and IgA anti- β_2 -GPI correlated significantly with anti-dsDNA, false-positive rapid plasma reagin, low C3, and low C4. Of the 3 isotypes of anti- β_2 -GPI, IgA had the strongest correlation with low C3 (p < 0.0001) and was the only isotype of anti- β_2 -GPI found to have a significant association with high erythrocyte sedimentation rate (ESR; p = 0.0001, OR 2.64, 95% CI 1.56–4.45) or anti-Sm (p = 0.0373, OR 1.58, 95% CI 1.04–2.41). No relationship was detected between C3, C4, anti-dsDNA, and the IgM anti- β_2 -GPI isotype (Table 4).

For the patients with IgA anti- β_2 -GPI alone (no IgG or IgM anti- β_2 -GPI), 22.1% had venous thrombosis and 11.9% had arterial thrombosis. For IgA aCL alone (IgG or IgM aCL), 11.1% had venous thrombosis and 5.3% had arterial thrombosis.

Other clinical associations with IgA anti- β_2 -GPI. A negative correlation was seen between the presence of IgA anti- β_2 -GPI and history of Sjögren's syndrome, dry eye, and dry mouth (OR 0.58, 95% CI 0.34–1.00, p = 0.0512; OR 0.56, 95% CI 0.35–0.90, p = 0.0142; and OR 0.56, 95% CI 0.32–1.00, p = 0.0566, respectively; Table 5). Significant associations were found between IgA anti- β_2 -GPI positivity and histories of pulmonary hypertension and pulmonary fibrosis (OR 2.26, 95% CI 1.15–4.56, p = 0.0240; and OR 1.89, 95% CI 1.06–3.35, p = 0.0445, respectively).

Table 1. Frequency of anti- β_2 glycoprotein I and anticardiolipin, by isotype, in 796 patients with systemic lupus erythematosus.

	No. Patients (%)
Isotypes of anti- β_2 -GPI	
IgG+ only	12 (1.5)
IgM+ only	74 (9.3)
IgA+ only	104 (13.1)
IgG, IgM+	5 (0.6)
IgG, IgA+	10 (1.3)
IgM, , IgA+	31 (3.9)
IgG, IgM, IgA+	16 (2.0)
Any IgG	43 (5.4)
Any IgM	126 (15.8)
Any IgA	161 (20.2)
Isotypes of anticardiolipin	
IgG+ only	114 (14.3)
IgM+ only	114 (14.3)
IgA+ only	19 (2.4)
IgG, IgM+	68 (8.5)
IgG, IgA+	11 (1.4)
IgM, , IgA+	9 (1.1)
IgG, IgM, IgA+	29 (3.6)
Any IgG	222 (27.9)
Any IgM	220 (27.6)
Any IgA	68 (8.5)

Table 2. Distribution of anti- β_2 glycoprotein I (anti- β_2 -GPI) isotypes by ethnicity.

Ethnicity	Anti- β_2 -GPI, Total, n (%)	IgG Anti- β_2 -GPI, n (%)	IgM Anti- β_2 -GPI, n (%)	IgA Anti- β_2 -GPI, n (%)
Asian	7 (2.4)	2 (4.6)	1 (0.8)	4 (2.5)
African American	118 (37.2)	6 (14.0)	45 (36)	67 (41.9)
Hispanic	8 (2.8)	(0)	3 (2.4)	5 (3.1)
Caucasian	195 (57.6)	35 (81.4)	76 (60.8)	84 (52.5)

Table 3. Associations of anti- β_2 glycoprotein I (anti- β_2 -GPI), by isotype, in systemic lupus erythematosus.

Factor	IgA Anti- β_2 -GPI Positive, %	IgA Anti- β_2 -GPI Negative, %	p for IgA Anti- β_2 -GPI	OR for IgA Anti- β_2 -GPI (95% CI)	OR for IgG Anti- β_2 -GPI (95% CI)	OR for IgM Anti- β_2 -GPI (95% CI)
Transient ischemic attack	5.7	2.6	0.070	2.30 (1.00–5.31)	3.69 (1.21–11.3)	2.64 (1.11–6.27)
Superficial thrombophlebitis	4.4	3.7	0.647	1.20 (0.51–2.85)	3.78 (1.37–10.4)	1.99 (0.86–4.57)
Deep venous thrombosis	19.9	10.7	0.003	2.08 (1.31–3.30)	3.33 (1.67–6.62)	0.94 (0.52–1.69)
Other venous thrombosis	4.4	4.2	0.827	1.06 (0.45–2.49)	3.36 (1.23–9.17)	1.18 (0.48–2.92)
Stroke	11.9	7	0.050	1.79 (1.01–3.15)	2.39 (1.02–5.61)	1.27 (0.66–2.46)
Myocardial infarction	1.2	2.9	0.397	0.43 (0.10–1.87)	0.91 (0.12–6.95)	0.27 (0.04–2.06)
Total arterial thrombosis (TIA, stroke, MI, and digital gangrene)	17.2	12.5	0.083	1.44 (0.9–2.34)	2.18 (1.04–4.60)	1.42 (0.84–2.39)
Total venous thrombosis (superficial, DVT, and other venous)	24.2	15.79	0.0097	1.70 (1.12–2.59)	3.03 (1.58–5.79)	1.22 (0.75–1.97)

TIA: transient ischemic attack; MI: myocardial infarction; DVT: deep venous thrombosis.

DISCUSSION

We found that 20% of our SLE cohort had IgA anti- β_2 -GPI, less than previous prevalence studies of 58%¹, 25%², 38.1%⁵, and 59%⁶. IgA anti- β_2 -GPI was the most frequent anti- β_2 -GPI isotype detected in all ethnicities. Anti- β_2 -GPI was found more frequently in whites, yet the isotype with the highest prevalence in African Americans was IgA. There is evidence to support a high prevalence of IgA aCL and IgA anti- β_2 -GPI in African Americans with SLE as compared with other ethnic groups^{3,7}.

The association of the IgA isotype of antiphospholipid antibodies (aPL) with thrombosis and pregnancy morbidity has remained controversial^{1,5,15,16,17}. In our study, IgA anti- β_2 -GPI was significantly ($p = 0.0097$) associated with venous thrombosis, but not arterial thrombosis. Similarly, Fanopoulos, *et al* concluded that there was a significantly higher frequency ($p < 0.01$) and level ($p < 0.05$) of IgA anti- β_2 -GPI in patients with SLE who have APS than in those without APS manifestations¹. Lakos, *et al* demonstrated a strong relationship between increased IgA anti- β_2 -GPI antibody levels and a history of venous thrombosis ($p = 0.007$), thrombocytopenia ($p = 0.02$), heart valve disease ($p = 0.02$), livedo reticularis ($p = 0.01$), and epilepsy ($p = 0.01$)⁶. In the analysis of 270 patients with SLE by Lee, *et al*, stepwise multivariate logistic regression analysis was used to determine that IgA was the isotype of anti- β_2 -GPI most related to the occurrence of venous thrombosis⁵.

Other studies, in contrast, have not found a significant association of IgA anti- β_2 -GPI with thrombosis in SLE^{15,17}.

The discrepancies between clinical findings in these other published results and ours may be attributable to the overall lower prevalence of IgA anti- β_2 -GPI found in their cohorts of patients with SLE, such as 8/134 (6%)¹⁵ and 23/130 (17.7%)¹⁷. Further, lack of standardization of assays used for the measurement of aCL and anti- β_2 -GPI, as well as the ethnic composition of the populations studied, may contribute to the discrepancies seen in the frequency and isotype distribution of these aPL among studies^{1,15,17}.

Our study found, for the first time, that IgA anti- β_2 -GPI was associated with both pulmonary hypertension and pulmonary fibrosis. The Euro-Phospholipid Project found a 2.2% prevalence of pulmonary hypertension in its prospective multicenter study of 1000 consecutive patients with aPL associated with primary APS (53.1% of patients), SLE (36.2%), lupus-like syndrome (5%), or other diseases (5.9%)¹⁸. Asherson and Cervera found that aPL may activate and cause endothelial cell damage, leading to the vascular changes seen in patients with pulmonary hypertension¹⁹. The association with pulmonary fibrosis is not understood.

We found a negative association, for the first time, between IgA anti- β_2 -GPI and secondary Sjögren's syndrome. Anti- β_2 -GPI has not been measured in most studies of patients with Sjögren's syndrome^{20,21,22}. However, when anti- β_2 -GPI was measured in primary Sjögren's syndrome, only 4% of 74 patients were positive in 1 study²¹ and none of 80 patients was positive in another²². It has been reported that the frequency of aPL in secondary Sjögren's syn-

Table 4. Comparison of isotypes of anti- β_2 glycoprotein I (anti- β_2 -GPI) with serologic tests in systemic lupus erythematosus.

Factor	IgG Anti- β_2 -GPI Present*, %	IgG Anti- β_2 -GPI Absent, %	p	OR (95% CI)
Coombs positivity	36.8	14.5	0.0010	3.37 (1.68–6.75)
Lupus anticoagulant (by RVVT)	86.1	26.2	< 0.0001	17.4 (7.22–41.8)
Anticardiolipin	97.6	59.2	< 0.0001	28.2 (3.85–206)
Anti-dsDNA	74.4	56.4	0.0254	2.25 (1.12–4.53)
Anti-Sm	14.3	18.5	0.6809	0.73 (0.30–1.77)
False-positive test for syphilis	51.3	13.2	< 0.0001	6.92 (3.55–13.5)
Low CH50	28.2	12.5	0.0122	2.75 (1.32–5.73)
Low C3	69.8	51.2	0.0187	2.20 (1.13–4.28)
Low C4	67.4	45.6	0.0069	2.48 (1.29–4.76)
Anti-Ro/SSA	21.4	30.0	0.2973	0.64 (0.30–1.35)
ESR	88.4	77.0	0.0913	2.27 (0.88–5.86)
	IgM Anti- β_2 -GPI Present	IgM Anti- β_2 -GPI Absent		
Coombs positivity	25.0	14.2	0.0070	2.01 (1.24–3.28)
Lupus anticoagulant (by RVVT)	52.4	25.2	< 0.0001	3.28 (2.21–4.86)
Anticardiolipin	84.0	57.0	< 0.0001	3.96 (2.40–6.55)
Anti-dsDNA	63.2	56.3	0.1677	1.33 (0.90–1.98)
Anti-Sm	13.1	19.3	0.1254	0.63 (0.36–1.11)
False positive test for syphilis	35.1	11.5	< 0.0001	4.15 (2.62–6.57)
Low CH50	17.7	12.5	0.1405	1.50 (0.88–2.55)
Low C3	58.4	51.1	0.1434	1.35 (0.91–1.96)
Low C4	52.8	45.6	0.1438	1.33 (0.91–1.96)
Anti-Ro/SSA	21.8	31.0	0.0414	0.62 (0.39–0.98)
ESR	78.4	77.5	0.9069	1.06 (0.66–1.68)
	IgA Anti- β_2 -GPI Present	IgA Anti- β_2 -GPI Absent		
Coombs positivity	20.7	14.7	0.0932	1.51 (0.94–2.42)
Lupus anticoagulant (by RVVT)	48.7	24.6	< 0.0001	2.91 (2.03–4.18)
Anticardiolipin	77.5	57.1	< 0.0001	2.58 (1.73–3.87)
Anti-dsDNA	76.9	52.4	< 0.0001	3.02 (2.03–4.50)
Anti-Sm	24.2	16.8	0.0373	1.58 (1.04–2.41)
False positive test for syphilis	28.6	11.8	< 0.0001	2.98 (1.92–4.62)
Low CH50	22.5	10.9	0.0004	2.37 (1.49–3.77)
Low C3	70.6	47.5	< 0.0001	2.66 (1.83–3.86)
Low C4	62.5	42.7	< 0.0001	2.23 (1.56–3.19)
Anti-Ro/SSA	32.9	28.7	0.3289	1.22 (0.84–1.77)
ESR	88.7	74.8	0.0001	2.64 (1.56–4.45)

* Positive value \geq 20 phospholipid units. RVVT: Russell viper venom time; ESR: erythrocyte sedimentation rate.

Table 5. Relationship of IgA anti- β_2 glycoprotein I (anti- β_2 -GPI) to some clinical manifestations.

Factor	IgA Anti- β_2 -GPI Present	IgA Anti- β_2 -GPI Absent	p	OR (95% CI)
Sjögren's syndrome, %	10.6	17.0	0.05	0.58 (0.34–1.00)
Dry eye, %	15.0	23.9	0.01	0.56 (0.35–0.90)
Dry mouth, %	9.4	15.5	0.06	0.56 (0.32–1.00)
SLICC/ACR Damage Index				
Pulmonary hypertension, %	8.9	4.1	0.02	2.26 (1.15–4.56)
Pulmonary fibrosis, %	12.0	6.8	0.04	1.89 (1.06–3.35)
Total pulmonary damage	0.27 \pm 0.61	0.16 \pm 0.45	0.007	

SLICC/ACR: Systemic Lupus International Collaborating Clinics/American College of Rheumatology.

drome is more linked to the underlying autoimmune disease, which appears to be the case in SLE²³.

Our study supports that IgA anti- β_2 -GPI is frequent in patients with SLE of all ethnicities. IgA anti- β_2 -GPI is asso-

ciated with venous thrombosis. In contrast to the IgG or IgM isotypes, IgA anti- β_2 -GPI positivity was significantly associated with the presence of serological markers such as high ESR, low C3, and anti-Sm antibodies. IgA anti- β_2 -GPI also

differs from the other isotypes of anti- β_2 -GPI as it has significantly positive clinical correlations with pulmonary hypertension and pulmonary fibrosis, and is negatively associated with Sjögren's syndrome, dry eye, and dry mouth in patients with SLE. Given our results, we suggest that measuring IgA anti- β_2 -GPI may be important for assessing the risk of thrombosis and pulmonary disease in patients with SLE. Moreover, the classification criteria for the APS in patients with SLE should be revised to include IgA anti- β_2 -GPI.

REFERENCES

1. Fanopoulos D, Teodorescu MR, Varga J, Teodorescu M. High frequency of abnormal levels of IgA anti- β_2 -glycoprotein I antibodies in patients with systemic lupus erythematosus: relationship with antiphospholipid syndrome. *J Rheumatol* 1998;25:675-80.
2. Tsutsumi A, Matsuura E, Ichikawa K, Fujisaku A, Mukai M, Koike T. IgA class anti- β_2 -glycoprotein I in patients with systemic lupus erythematosus. *J Rheumatol* 1998;25:74-8.
3. Cucurull E, Gharavi AE, Diri E, Mendez E, Kapoor D, Espinoza LR. IgA anticardiolipin and anti- β_2 -glycoprotein I are the most prevalent isotypes in African American patients with systemic lupus erythematosus. *Am J Med Sci* 1999;318:55-60.
4. Danowski A, Kickler T, Petri M. Anti- β_2 -glycoprotein I: prevalence, clinical correlations, and importance of persistent positivity in patients with antiphospholipid syndrome and systemic lupus erythematosus. *J Rheumatol* 2006;33:1775-9.
5. Lee S, Cho M, Joo Y, Kim W, Hong Y, Min J, et al. Isotypes of anti- β_2 -glycoprotein I antibodies: association with thrombosis in patients with systemic lupus erythematosus. *J Rheumatol* 2001;28:520-4.
6. Lakos G, Kiss E, Regeczy P, Tarján P, Soltész P, Zeher M, et al. Isotype distribution and clinical relevance of anti- β_2 -glycoprotein I antibodies: importance of IgA isotype. *Clin Exp Immunol* 1999;117:574-9.
7. Diri E, Cucurull E, Gharavi AE, Kapoor D, Mendez EA, Scopelitis E, et al. Antiphospholipid (Hughes') syndrome in African-Americans: IgA aCL and anti- β_2 -glycoprotein I is the most frequent isotype. *Lupus* 1999;9:33-41.
8. Tan EM, Cohen AS, Fries JF, Masi AT, McShane DJ, Rothfield NF, et al. The 1982 revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum* 1982;25:1271-7.
9. Hochberg MC. Updating the American College of Rheumatology revised criteria for the classification of systemic lupus erythematosus [letter]. *Arthritis Rheum* 1997;40:1725.
10. Petri M, Rheinschmidt M, Whiting-O'Keefe Q, Hellmann D, Corash L. The frequency of lupus anticoagulant in systemic lupus erythematosus. A study of sixty consecutive patients by activated partial thromboplastin time, Russell viper venom time, and cardiolipin antibody level. *Ann Intern Med* 1987;106:524-31.
11. Somers E, Magder LS, Petri M. Antiphospholipid antibodies and incidence of venous thrombosis in a cohort of patients with SLE. *J Rheumatol* 2002;29:2531-6.
12. Petri M, Kim MY, Kalunian KC, Grossman J, Hahn BH, Sammaritano LR, et al. Combined oral contraceptives in women with systemic lupus erythematosus. *N Engl J Med* 2005;353:2550-8.
13. Bombardier C, Gladman DD, Chang CH, Urowitz MB, and the Committee on Prognosis Studies in SLE. Development of the Disease Activity Index: the SLEDAI. *Arthritis Rheum* 1992;35:630-40.
14. Gladman D, Ginzler E, Goldsmith C, Fortin P, Liang M, Urowitz M, et al. Systemic Lupus International Collaborative Clinics: Development of a damage index in systemic lupus erythematosus. *J Rheumatol* 1992;29:1820-1.
15. Bertolaccini ML, Atsumi T, Contreras AE, Khamashta MA, Hughes GRV. The value of IgA antiphospholipid testing for diagnosis of antiphospholipid (Hughes) syndrome in systemic lupus erythematosus. *J Rheumatol* 2001;28:2637-43.
16. Carmo-Pereira S, Bertolaccini ML, Escudero-Contreras A, Khamashta MA. Value of IgA anticardiolipin and anti- β_2 -glycoprotein I antibody testing in patients with pregnancy morbidity. *Ann Rheum Dis* 2003;62:540-3.
17. Samarkos M, Davies KA, Gordon C, Loizou S. Clinical significance of IgA anticardiolipin and anti- β_2 -gpi antibodies in patients with systemic lupus erythematosus and primary antiphospholipid syndrome. *Clin Rheumatol* 2006;25:199-204.
18. Cervera R, Piette J, Font J, Khamashta MA, Shoenfeld Y, Camps MT, et al. Antiphospholipid syndrome. Clinical and immunologic manifestations and patterns of disease expression in a cohort of 1000 patients. *Arthritis Rheum* 2002;46:1019-27.
19. Asherson RA, Cervera R. Pulmonary hypertension, antiphospholipid antibodies, and syndromes. *Clin Rev Allergy Immunol* 2007;32:153-8.
20. Ramos-Casals M, Nardi N, Brito-Zeron P, Aguilo S, Gil V, Delgado G, et al. Atypical autoantibodies in patients with primary Sjogren syndrome: clinical characteristics and follow-up of 82 cases. *Semin Arthritis Rheum* 2006;35:312-21.
21. Fauchais AL, Lambert M, Launay D, Michon-Pasturel U, Queyrel V, Nguyen N, et al. Antiphospholipid antibodies in primary Sjogren's syndrome: prevalence and clinical significance in a series of 74 patients. *Lupus* 2004;13:245-8.
22. Cervera R, Garcia-Carrasco M, Font J, Ramos M, Reverter JC, Muñoz FJ, et al. Antiphospholipid antibodies in primary Sjogren's syndrome: prevalence and clinical significance in a series of 80 patients. *Clin Exp Rheumatol* 1997;15:361-5.
23. Merkel PA, Chang Y, Pierangeli SS, Convery K, Harris EN, Polisson RP. The prevalence and clinical associations of anticardiolipin antibodies in a large inception cohort of patients with connective tissue diseases. *Am J Med* 1996;101:576-83.