

# Soluble Receptor for Advanced Glycation Endproducts Is Decreased in Patients with Juvenile Idiopathic Arthritis (ERA Category) and Inversely Correlates with Disease Activity and S100A12 Levels

ARPITA MYLES, VISHAD VISWANATH, YOGESH PREET SINGH, and AMITA AGGARWAL

**ABSTRACT. Objective.** Membrane-bound receptor for advanced glycation endproducts (mRAGE) is overexpressed in response to increasing concentrations of its ligand (e.g., S100A12) and triggers an inflammatory immune response. In contrast, soluble RAGE (sRAGE) acts as a decoy receptor and downmodulates inflammation. Decreased sRAGE levels are associated with autoimmune diseases; however, limited data are available in juvenile idiopathic arthritis (JIA). We studied sRAGE levels in patients with JIA [enthesitis-related arthritis (ERA) category].

**Methods.** sRAGE levels were estimated in the serum of patients with ERA JIA (n = 101), systemic-onset JIA and polyarticular JIA (n = 10 each), and healthy controls (n = 45). Synovial fluid (SF) sRAGE was measured in patients with ERA, rheumatoid arthritis, reactive arthritis, and osteoarthritis (n = 10). Levels of S100A12 were also measured. Twenty-four patients with ERA were followed for 4 months. Disease activity was assessed by swollen joint count (SJC), tender joint count (TJC), and erythrocyte sedimentation rate (ESR). All levels are expressed as median (range).

**Results.** The serum sRAGE (pg/ml) level was significantly lower in patients compared to healthy controls [515 (64–1887) vs 1542 (627–3159);  $p < 0.0001$ ]. In paired samples, SF had lower levels compared to corresponding plasma level [102 (51–799) vs 481 (134–1006);  $p < 0.0001$ ]. The level of S100A12 (ng/ml) was higher in SF (1042; 573–1415) than serum (638; 208–779). Serum sRAGE correlated negatively with S100A12 levels ( $r = -0.474$ ;  $p < 0.01$ ), ESR ( $r = -0.306$ ;  $p < 0.01$ ), and SJC ( $r = -0.237$ ;  $p < 0.05$ ), but not with TJC ( $r = -0.134$ ;  $p = \text{NS}$ ). The levels of sRAGE remained stable over time in patients with stable disease.

**Conclusion.** Levels of sRAGE are reduced in patients with ERA and correlate negatively with disease activity and S100A12 levels. sRAGE may be a modulator of inflammation in these patients. (First Release July 1 2011; J Rheumatol 2011;38:1994–9; doi:10.3899/jrheum.110058)

## Key Indexing Terms:

JUVENILE ARTHRITIS  
DISEASE ACTIVITY

INFLAMMATION  
RECEPTOR FOR ADVANCED GLYCATION ENDPRODUCTS

The receptor for advanced glycation endproducts (RAGE) is a cell-surface molecule belonging to the immunoglobulin superfamily and is expressed by several cells, including immune cells such as neutrophils, macrophages, and T cells<sup>1</sup>. It is encoded in the Class III region of the major histocompatibility complex. It recognizes a wide repertory of ligands<sup>2</sup> — the advanced glycation endproducts, the S100/calgranulin family

of proinflammatory cytokine-like proteins, and the high mobility group box protein-1. Receptor-ligand binding results in transcription of proinflammatory factors and in overexpression of the receptor on the cell surface. Increasing evidence suggests that membrane-bound RAGE (mRAGE) acts like a non-canonical pattern recognition receptor and its dysregulation leads to persistent inflammation in diseases such as diabetes, atherosclerosis, arthritis, and Alzheimer's disease<sup>3</sup>.

Soluble RAGE (sRAGE) lacks the short cytoplasmic domain that not only anchors the full receptor to a cell, but also transmits downstream signals. sRAGE functions as a “decoy receptor,” competing with its membrane-bound form for the same ligand and causing a reduction in overall signaling<sup>4</sup>. sRAGE may also act as a direct inhibitor of leukocyte recruitment<sup>5</sup>. Thus, it is capable of reducing the inflammation caused by mRAGE. sRAGE interacts with the integrin Mac-1 and triggers a proinflammatory cascade<sup>6</sup>, suggesting a dual function for sRAGE.

From the Department of Clinical Immunology, Sanjay Gandhi Postgraduate Institute of Medical Sciences, Lucknow, India.

Supported by a grant from the Department of Biotechnology, Government of India, to Dr. Aggarwal. Ms. Myles is supported by the Council of Scientific and Industrial Research-Junior Research Fellowship.

A. Myles, MSc; V. Viswanath, MD; Y.P. Singh, DM; A. Aggarwal, DM, Additional Professor, Department of Clinical Immunology, Sanjay Gandhi Postgraduate Institute of Medical Sciences.

Address correspondence to Dr. A. Aggarwal, Department of Clinical Immunology, Sanjay Gandhi Postgraduate Institute of Medical Sciences, Lucknow, India. E-mail: amita@sippi.ac.in; aa.amita@gmail.com  
Accepted for publication April 21, 2011.

Calgranulin C, also called S100A12 or EN-RAGE (extracellular newly identified RAGE-binding protein), is a member of a multigene family involved in calcium-dependent regulation of several cellular activities such as protein phosphorylation, cellular homeostasis, and even inflammation<sup>7</sup>. It is secreted mainly by neutrophils and activated macrophages<sup>8</sup> and is now recognized as a damage-associated molecular pattern molecule (DAMP), a group that represents endogenous ligands of pattern recognition receptors<sup>9</sup>. It binds to mRAGE and induces inflammatory cascade<sup>10</sup>.

There are several studies in which dysregulation in levels of sRAGE and S100A12 have been associated with inflammatory diseases. Studies done in different arthritides found serum levels of sRAGE to be lower in rheumatoid arthritis (RA)<sup>11</sup>, osteoarthritis (OA)<sup>12</sup>, and Sjögren's syndrome<sup>13</sup>. sRAGE levels also correlated negatively with S100A12 levels in Kawasaki disease<sup>14</sup>. In addition, serum sRAGE levels in patients with RA were found to be negatively associated with current smoking, family history of cardiovascular disease, history of vasculitis, blood pressure, presence of rheumatoid factor, and serum levels of C-reactive protein (CRP) and S100A12, again suggesting that sRAGE acts as an antiinflammatory molecule and S100A12 as a proinflammatory one<sup>15</sup>.

Juvenile idiopathic arthritis (JIA) is another disease characterized by inflammation and articular damage. Among all the categories of JIA<sup>16</sup>, enthesitis-related arthritis (ERA) is most prevalent in India. Studies have shown elevated levels of S100A12 in serum of patients with JIA<sup>8,14,17</sup>. Data on sRAGE levels in JIA are limited. Only a single study on Kawasaki disease had used patients with JIA as disease controls and shown that sRAGE levels were lower in the serum of children with systemic-onset JIA and were similar to healthy controls in oligoarticular and polyarticular JIA (polyJIA)<sup>14</sup>. That study had not included children with ERA. In our study we measured sRAGE and S100A12 levels in serum, synovial fluid (SF), and paired plasma of patients with ERA and studied its correlation with disease activity. Further, we studied the proportion of these patients longitudinally to see whether sRAGE levels change over time and whether they correlate with changes in disease activity.

## MATERIALS AND METHODS

**Patients and controls.** Patients with ERA who satisfied International League of Associations for Rheumatology criteria<sup>16</sup> and who gave or whose parents gave written informed consent were enrolled in the study. Nonrelated healthy controls matched for age and sex were enrolled from blood bank donors. Stored samples of patients with systemic-onset (SoJIA) and polyJIA were used as disease controls. The study was approved by the institutional ethics committee. We also measured sRAGE levels in stored SF of patients with RA, OA, and reactive arthritis (ReA) as disease controls.

**Separation of serum, plasma, and SF.** Blood samples and SF were centrifuged at 1000 g for 10 min within 30 min of withdrawal. Samples for plasma were collected in lithium heparin. The serum/plasma/cell-free SF thus obtained was stored at -80°C.

**ELISA.** sRAGE (DuoSet ELISA development kit; R&D Systems, Minneapolis, MN, USA) and S100A12 (Abnova Corp., Taipei, Taiwan) levels

were measured in serum/plasma/cell-free SF and determined using sandwich ELISA as per manufacturer's instructions.

**Statistical analysis.** Data were analyzed using GraphPad Prism (GraphPad Software Inc., La Jolla, CA, USA) and SPSS 13.0 software (SPSS Inc., Rainbow Technologies, Somers, NY, USA). Because we found no difference between serum and plasma levels in ERA, we analyzed both these samples together and henceforth they have been referred to as serum. The Mann-Whitney U test was used for comparing differences in serum analyte concentrations of patients and controls, while analysis of paired plasma and SF samples was done by Wilcoxon signed-rank test. The Kruskal-Wallis test was used for multiple group comparison. The relationship between sRAGE concentration and disease activity was measured by Spearman's rank correlation. All results are expressed as median (range).

## RESULTS

The study involved 101 patients with ERA (median age at sample collection 15 years and median disease duration 3 years; 98 males, 84 HLA-B27-positive; Table 1), 10 patients with SoJIA (median age at sample collection 13 years, 8 males, median disease duration 4.8 years, 9 patients taking methotrexate and 8 taking steroids; Table 1), 10 patients with polyJIA (median age at sample collection 13.5 years, 6 males, median disease duration 4 years, 9 patients taking methotrexate and 6 taking steroids; Table 1), and 45 healthy controls (median age 24 years; all males).

Median sRAGE levels were significantly lower ( $p < 0.0001$ ) in patients with ERA (515; 64–1887 pg/ml) than in healthy controls (1542; 627–3159 pg/ml). There was no difference in levels between patients with ERA, patients with SoJIA (300; 165–400 pg/ml), and patients with polyJIA (380; 180–1000 pg/ml; Figure 1A).

Median S100A12 levels were significantly higher ( $p < 0.01$ ) in patients with ERA (498; 153–783 pg/ml) than in healthy controls (144; 108–307 pg/ml). There was no difference in levels between patients with ERA, patients with SoJIA (510; 205–819.6 pg/ml), and patients with polyJIA (423; 165–710 pg/ml; Figure 1B).

SF and paired plasma were available from 10 patients with ERA (median age at onset 10.5 years and median disease duration 8 years; all males). Eight were HLA-B27-positive, 1 was negative, and data were not available for 1 patient. Five patients were taking nonsteroidal antiinflammatory drugs and 3 were taking methotrexate. Median sRAGE levels (pg/ml) in SF (102.5; 51–799) were lower than those in plasma (481.6; 134–1006;  $p < 0.0001$ ; Figure 2A). Median SF S100A12 levels (1042; 573–1415) were higher ( $p < 0.01$ ) than in plasma (638; 208–779; Figure 2B).


Median sRAGE levels in ERA SF did not differ significantly from those in RA (180; 64–430), OA (180; 64–580), and ReA SF (170; 140–290).

The serum sRAGE levels in ERA correlated negatively with S100A12 levels ( $r = -0.474$ ;  $p < 0.01$ ; Figure 3). The levels also showed inverse correlation with ESR ( $r = -0.306$ ;  $p < 0.01$ ), swollen joint count (SJC;  $r = -0.237$ ;  $p < 0.05$ ), but not with TJC ( $r = -0.134$ ;  $p =$  nonsignificant), which were taken as markers of disease activity (Figure 4). The sRAGE

Table 1. Clinical details of patients with juvenile idiopathic arthritis (JIA).

Characteristics	ERA	Polyarticular JIA	Systemic-onset JIA
No. patients	101	10	10
Median age at onset, yrs (range)	9.5 (4–16)	8.2 (2–16)	6.75 (0.8–16)
Median age at sample collection, yrs (range)	15 (12–24)	13.5 (7–24)	13 (1–30)
Median disease duration, yrs (range)	3 (2–12)	4 (0.8–13)	4.8 (0.4–18)
Male: female	98:3	6:4	8:2
DMARD usage	65	9	9
Prednisolone usage	8	6	8
Tender joint count, n (range)	0 (0–9)	0 (0–4)	0 (0–6)
Swollen joint count, n (range)	0 (0–7)	1 (0–6)	0 (0–2)
ESR, mm/h (range)	34 (4–160)	54 (16–117)	69 (10–180)
HLA-B27-positive samples	84	0	Not Done

ERA: enthesitis-related arthritis; DMARD: disease-modifying antirheumatic drugs; ESR: erythrocyte sedimentation rate.



and S100A12 levels did not have any association with the age of the patients.

In 24 paired serum samples obtained at a median difference of 4 months, there was no significant difference between SJC, TJC, ESR, and sRAGE at baseline and followup.

## DISCUSSION

Our study revealed that levels of sRAGE in the serum/plasma of patients with JIA-ERA were lower and levels of S100A12 higher compared to healthy controls, but were similar to those of patients with SoJIA and polyJIA. Further, in paired samples, the SF sRAGE level was lower and S100A12 levels were higher than the corresponding plasma level. Serum sRAGE levels had inverse correlation with disease activity measures such as ESR and SJC, and also with S100A12 concentration. However, in patients with stable disease sRAGE levels did not change significantly.

Our observation of reduced sRAGE and elevated S100A12 levels in children with ERA-JIA is similar to the data reported in children with SoJIA<sup>14</sup>. We found lower levels of S100A12 in SoJIA compared to the study by Wittkowski, *et al*<sup>14</sup>. The likely reason could be that their study enrolled drug-naive patients at the start of illness, while most of our patients were taking disease-modifying antirheumatic drugs and steroids and had a longer disease duration. Similar reduction in sRAGE and elevation in S100A12 levels is seen in RA<sup>15</sup> and OA<sup>12</sup>. On the other hand, while 1 study showed that patients with oligoarticular and polyJIA have sRAGE levels similar to those of healthy controls<sup>14</sup>, we found lower sRAGE levels in patients with polyJIA compared to healthy controls. S100A12 is known to stimulate production of proinflammatory cytokines upon binding to mRAGE<sup>10</sup>; it is likely that an

Figure 1. Levels of soluble receptor for advanced glycation endproducts (sRAGE; A) and S100A12 (B) in serum of healthy controls and patients with enthesitis-related arthritis (ERA), systemic-onset juvenile idiopathic arthritis (SoJIA), and polyarticular JIA (polyJIA). \*\*\*p < 0.0001; \*\*p < 0.01. No difference was seen between sRAGE and S100A12 levels in ERA, SoJIA, and polyJIA sera.




Figure 2. Soluble receptor for advanced glycation endproducts (sRAGE) and S100A12 levels in paired synovial fluid and serum of ERA patients (n = 10). Each dot represents an individual patient. Median is marked on the plot. p value calculated by Wilcoxon signed-rank test. A. sRAGE levels. B. S100A12 levels.




Figure 3. Correlation between levels of soluble receptor for advanced glycation endproducts (sRAGE; pg/ml) and S100A12 (ng/ml) in serum of patients with enthesitis-related arthritis (n = 30).  $r = -0.474$ .

increase in its levels contributes significantly to JIA pathology. This effect is further worsened by decrease in sRAGE concentration, which, under normal conditions, mops up mRAGE ligand, thereby modulating its signaling.

SF sRAGE levels were even lower compared to the plasma levels. This is reflective of the potent regulatory role played by sRAGE at the local site of inflammation. Our data are similar to previous reports in RA<sup>11</sup> and OA<sup>12</sup>. Lower sRAGE levels in SF may result in continued activation of RAGE pathways, leading to inflammation. Further, S100A12 levels were markedly higher in SF than in plasma. This observation is again in tandem with findings in RA<sup>11</sup>, polyJIA, and oligoar-

ticular JIA<sup>17</sup>. More recently, Baillet, *et al*<sup>18</sup> showed that S100A12 was one of the most upregulated proteins in RA SF and could distinguish RA from other joint diseases with great accuracy.

Synovial tissue samples from both RA and OA show expression of mRAGE; however, the levels were higher in RA<sup>19</sup>. It is mainly expressed by macrophages, T cells, and B cells. Cultured human synovial fibroblasts also constitutively express RAGE and on stimulation with cognate ligands produce monocyte chemoattractant proteins-1<sup>20</sup>. S100A12, after binding to RAGE, produces proinflammatory cytokines and matrix metalloprotease<sup>21</sup>. Thus, modulation of RAGE expression in the synovial compartment might be a therapeutic avenue. Indeed, Pullerits, *et al*<sup>22</sup> measured and demonstrated autoantibodies against RAGE in the SF of patients with RA. The autoantibodies' presence correlated with less erosive disease, suggesting that blocking the RAGE pathway can reduce severity of disease.

In our study, negative correlation was seen between sRAGE levels and measures of disease activity such as SJC and ESR. In RA, serum sRAGE levels correlated with CRP levels, presence of rheumatoid factor, and vasculitis, but not with age, duration of disease, or erosions<sup>15</sup>. In another study, lower urinary sRAGE level had an association with increased risk of atherothrombosis in RA<sup>23</sup>. In SoJIA and Kawasaki disease, sRAGE levels correlated negatively with CRP levels<sup>14</sup>. In our longitudinal study, we did not find any difference in sRAGE levels, suggesting that sRAGE remains stable over time, when there is no change in disease activity. Thus it can be used as a biomarker. In Kawasaki disease, sRAGE levels normalize once the inflammation subsides after intravenous immunoglobulin treatment<sup>14</sup>. In RA followup, study levels at




Figure 4. Correlation between (A) levels of soluble receptor for advanced glycation endproducts (sRAGE) and erythrocyte sedimentation rate (ESR);  $r = -0.306$ ;  $p < 0.05$ . B. sRAGE levels and tender joint count;  $r = -0.134$ ;  $p =$  nonsignificant. C. sRAGE levels and swollen joint count;  $r = -0.237$ ;  $p < 0.05$ . Each dot represents serum from individual patient with enthesitis-related arthritis (ERA;  $n = 101$ ).

1 year were significantly different from those at baseline, again suggesting inflammation-mediated regulation of sRAGE levels<sup>23</sup>. Indeed, RAGE mRNA increases in adherent macrophages on exposure to proinflammatory cytokines, especially interleukin 1 $\beta$ <sup>19</sup>.

The etiology and pathogenesis of JIA-ERA are unknown, but innate immunity is postulated to play a role in disease perpetuation. These triggers can be exogenous (microbial products)<sup>24</sup> or endogenous damage-associated molecular patterns (e.g., S100A12). They stimulate immune responses through pattern recognition receptors, and the mRAGE is one of them. sRAGE probably absorbs the ligands of the mRAGE and thus downmodulates the proinflammatory response. sRAGE also affects survival of monocytes/neutrophils and macrophage development<sup>25</sup>. In other words, a decrease in sRAGE levels probably tips the balance in favor of inflammation. However,

we were unable to establish a cause-and-effect relationship between decreased sRAGE and increased S100A12 levels.

Our study suggests that patients with ERA have low serum and SF sRAGE levels and that these levels correlate negatively with disease activity as well as concentration of S100A12 protein. Thus sRAGE may be involved with downmodulation of inflammation in patients with ERA.

## REFERENCES

- Schmidt AM, Yan SD, Yan SF, Stern DM. The multiligand receptor RAGE as a progression factor amplifying immune and inflammatory responses. *J Clin Invest* 2001;108:949-55.
- Schmidt AM, Yan SD, Yan SF, Stern DM. The biology of the receptor for advanced glycation end products and its ligands. *Biochim Biophys Acta* 2000;1498:99-111.
- Sparvero LJ, Asafu-Adjei D, Kang R, Tang D, Amin N, Im J, et al. RAGE (receptor for advanced glycation endproducts), RAGE

- ligands, and their role in cancer and inflammation. *J Transl Med* 2009;7:17-38.
4. Lin Li, Park S, Lakatta EG. RAGE signaling in inflammation and arterial aging. *Front Biosci* 2009;14:1403-13.
  5. Chavakis T, Bierhaus A, Al-Fakhri N, Schneider D, Witte S, Linn T, et al. The pattern recognition receptor (RAGE) is a counter receptor for leukocyte integrins: a novel pathway for inflammatory cell recruitment. *J Exp Med* 2003;198:1507-15.
  6. Pullerits R, Brisslert M, Jonsson IM, Tarkowski A. Soluble receptor for advanced glycation end products triggers a proinflammatory cytokine cascade via beta 2 integrin Mac-1. *Arthritis Rheum* 2006;54:3898-907.
  7. Donato R. Functional roles of S100 proteins, calcium-binding proteins of the EF-hand type. *Biochim Biophys Acta* 1999;1450:191-231.
  8. Wittkowski H, Frosch M, Wulffraat N, Goldbach-Mansky R, Kallinich T, Kuemmerle-Deschner J, et al. S100A12 is a novel molecular marker differentiating systemic-onset juvenile idiopathic arthritis from other causes of fever of unknown origin. *Arthritis Rheum* 2008;58:3924-31.
  9. Foell D, Wittkowski H, Roth J. Mechanisms of disease: a 'DAMP' view of inflammatory arthritis. *Nat Clin Pract Rheumatol* 2007;3:382-90.
  10. Hofmann MA, Drury S, Fu C, Qu W, Taguchi A, Lu Y, et al. RAGE mediates a novel proinflammatory axis: a central cell surface receptor for S100/calgranulin polypeptides. *Cell* 1999;97:889-901.
  11. Pullerits R, Bokarewa M, Dahlberg L, Tarkowski A. Decreased levels of soluble receptor for advanced glycation end products in patients with rheumatoid arthritis indicating deficient inflammatory control. *Arthritis Res Ther* 2005;7:R817-24.
  12. Chayanupatkul M, Honsawek S. Soluble receptor for advanced glycation end products (sRAGE) in plasma and synovial fluid is inversely associated with disease severity of knee osteoarthritis. *Clin Biochem* 2010;43:1133-7.
  13. Stewart C, Cha S, Caudle RM, Berg K, Katz J. Decreased levels of soluble receptor for advanced glycation end products in patients with primary Sjögren's syndrome. *Rheumatol Int* 2008;28:771-6.
  14. Wittkowski H, Hirono K, Ichida F, Vogl T, Ye F, Yanlin X, et al. Acute Kawasaki disease is associated with reverse regulation of soluble receptor for advanced glycation end products and its proinflammatory ligand S100A12. *Arthritis Rheum* 2007;56:4174-81.
  15. Chen YS, Yan W, Geczy GL, Brown MA, Thomas R. Serum levels of soluble receptor for advanced glycation end products and of S100 proteins are associated with inflammatory, autoantibody, and classical risk markers of joint and vascular damage in rheumatoid arthritis. *Arthritis Res Ther* 2009;11:R39-50.
  16. Petty RE, Southwood TR, Manners P, Baum J, Glass DN, Goldenberg J, et al. International League of Associations for Rheumatology classification of juvenile idiopathic arthritis: second revision, Edmonton, 2001. *J Rheumatol* 2004;31:390-2.
  17. Foell D, Wittkowski H, Hammerschmidt I, Wulffraat N, Schmeling H, Frosch M, et al. Monitoring neutrophil activation in juvenile rheumatoid arthritis by S100A12 serum concentrations. *Arthritis Rheum* 2004;50:1286-95.
  18. Baillet A, Trocmé C, Berthier S, Arlotto M, Grange L, Chenau J, et al. Synovial fluid proteomic fingerprint: S100A8, S100A9 and S100A12 proteins discriminate rheumatoid arthritis from other inflammatory joint diseases. *Rheumatology* 2010;49:671-82.
  19. Sunahori K, Yamamura M, Yamana J, Takasugi K, Kawashima M, Makino H. Increased expression of receptor for advanced glycation end products by synovial tissue macrophages in rheumatoid arthritis. *Arthritis Rheum* 2006;54:97-104.
  20. Hou FF, Jiang JP, Guo JQ, Wang GB, Zhang X, Stern DM, et al. Receptor for advanced glycation end products on human synovial fibroblasts: role in the pathogenesis of dialysis-related amyloidosis. *J Am Soc Nephrol* 2002;13:1296-306.
  21. Tolboom TC, Pieterman E, van der Laan WH, Toes RE, Huidekoper AL, Nelissen RG, et al. Invasive properties of fibroblast-like synoviocytes: correlation with growth characteristics and expression of MMP-1, MMP-3, and MMP-10. *Ann Rheum Dis* 2002;61:975-80.
  22. Pullerits R, Bokarewa M, Dahlberg L, Tarkowski A. Synovial fluid expression of autoantibodies specific for RAGE relates to less erosive course of rheumatoid arthritis. *Rheumatology* 2007;46:1367-71.
  23. Ferrante E, Vazzana N, Santilli F, Di Cicco M, Lauriti C, Di Battista L, et al. Determinants of thromboxane biosynthesis in rheumatoid arthritis: Role of RAGE and oxidant stress. *Free Radic Biol Med* 2010;49:857-64.
  24. Ellis JA, Munro JE, Ponsonby AE. Possible environmental determinants of juvenile idiopathic arthritis. *Rheumatology* 2010;49:411-25.
  25. Wang Y, Wang H, Piper MG, McMaken S, Mo X, Opalek J, et al. sRAGE induces human monocyte survival and differentiation. *J Immunol* 2010;185:1822-35.