

## A Single-nucleotide Polymorphism of CCR6 (rs3093024) Is Associated with Susceptibility to Rheumatoid Arthritis But Not Ankylosing Spondylitis, in a Taiwanese Population

To the Editor:

Rheumatoid arthritis (RA) is a common autoimmune disease with chronic inflammatory reactions in the synovial tissue and joint destruction. Genetic and environmental factors contribute to the development of RA<sup>1</sup>. Using genome-wide association analysis, Kochi, *et al* identified a single nucleotide polymorphism (rs3093024) in the *CCR6* gene that was associated with the risk of rheumatoid arthritis<sup>2</sup>. Further, the genotype of *CCR6* correlated with the concentration of interleukin 17 in the sera of subjects with RA<sup>2</sup>.

We investigated whether the polymorphism of *CCR6* is associated with susceptibility to RA in a Taiwanese population. A total of 1441 subjects were recruited [400 patients with RA, 361 with ankylosing spondylitis (AS), and 680 healthy subjects]. The patients with RA and the healthy subjects were from the Kaohsiung Medical University Hospital. The diagnosis of RA was according to the American College of Rheumatology 1987 revised criteria for the classification of RA. The 361 patients with AS were recruited from Chung Shan Medical University Hospital. Patients with AS who met the New York AS diagnosis criteria were asked to participate. Our study was approved by the institutional review boards of the hospitals. Informed consent was obtained before any data were collected from the subjects. DNA purification from buffy coat was done using the Genra Puregene Blood Kit (Qiagen, Valencia, CA, USA). Genotyping was performed using TaqMan allelic discrimination assays (Applied Biosystems, Foster City, CA, USA).

We found that the distribution of genotype rs3093024 was in accord with Hardy-Weinberg equilibrium for both cases and controls. Importantly, a significant association between rs3093024 genotype and the patients with RA was observed (AA/AG vs GG; OR 1.48, 95% CI 1.13–1.94,  $p = 0.0046$ ; Table 1). Because the sex distribution between RA and control groups was very different, we further divided the original study subjects by sex and tested the sex-specific genetic effects of rs3093024. A significant association between rs3093024 and the risk of RA still existed (Table 1). The minor A allele of rs3093024 was associated with susceptibility to RA in the Taiwanese female population (AA/AG vs GG; OR 1.63, 95% CI 1.17–2.28,  $p = 0.0038$ ; Table 1).

AS is also an autoimmune disease affecting mainly joints in the spine and the sacroiliac joint<sup>3</sup>. Recently, several genetic polymorphisms have

been strongly implicated in its pathogenesis<sup>4,5,6,7</sup>. However, our results indicated that rs3093024 was not associated with susceptibility to AS (data not shown). We acknowledge that the modest AS sample size in the study was underpowered to detect the genetic effect of *CCR6*. A larger sample size is needed to replicate these findings.

We found a significant association between the genotype of *CCR6* (rs3093024) and the susceptibility to RA in a Taiwanese population. In agreement with previous reports in the Japanese population, our results strongly support the importance of the *CCR6* genotype in the risk of RA.

WEI-CHIAO CHANG, <sup>DPhil</sup>, Department of Medical Genetics, Kaohsiung Medical University, Kaohsiung; PENG-YEONG WOON, <sup>PhD</sup>, Department of Molecular Biology and Human Genetics, Tzu Chi University, Hualien; JAMES CHENG-CHUNG WEI, <sup>MD</sup>, Division of Allergy, Immunology and Rheumatology, Department of Medicine and Institute of Medicine, Chung Shan Medical University Hospital, Chung Shan Medical University; Graduate Institute of Integrated Medicine, China Medical University, Taichung; CHE-MAI CHANG, <sup>BSc</sup>; YU-WEN HSU, <sup>BSc</sup>, Department of Medical Genetics, Kaohsiung Medical University; YUH-CHERNG GUO, <sup>MD</sup>, Department of Neurology, Changhua Christian Hospital, Changhua, Taiwan; DAW-YANG HWANG, <sup>MD</sup>, Department of Neurology, Kaohsiung Municipal Hsiao-Kang Hospital, Kaohsiung Medical University, Kaohsiung, Taiwan; YUTA KOCHI, <sup>MD</sup>, Laboratory for Autoimmune Diseases, Center for Genomic Medicine, Riken, Yokohama, Japan; JENG-HSIEN YEN, <sup>MD</sup>, Division of Rheumatology, Department of Internal Medicine, Kaohsiung Medical University Hospital, Taiwan. Address correspondence to Dr. J.H. Yen or Dr. W.C. Chang, 100 Tzu Chi First Road, Kaohsiung City 807, Taiwan, Kaohsiung 701. E-mail: wcc@kmu.edu.tw or jehsye@kmu.edu.tw Supported by a grant from the National Science Council, Taiwan, ROC (NSC 100-2320-B-037-002).

### REFERENCES

1. Firestein GS. Evolving concepts of rheumatoid arthritis. *Nature* 2003;423:356-61.
2. Kochi Y, Okada Y, Suzuki A, Ikari K, Terao C, Takahashi A, *et al*. A regulatory variant in *CCR6* is associated with rheumatoid arthritis susceptibility. *Nat Genet* 2010;42:515-9.
3. Appel H, Lodenkemper C, Miossec P. Rheumatoid arthritis and ankylosing spondylitis — pathology of acute inflammation. *Clin Exp Rheumatol* 2009;4 Suppl 55:S15-9.

Table 1. Genotyping and allele frequency of *CCR6* single-nucleotide polymorphism in patients with rheumatoid arthritis and in controls. Significant values are shown in bold type.

Genotype	Cases (%), n = 400	Controls (%), n = 680	Genotype, p	p and OR (95% CI)		Allelic, p
				Dominant, p	Recessive, p	
AA	83 (21.3)	107 (16.1)	<b>0.0081**</b>	<b>0.0046**</b>	<b>0.0323*</b>	<b>0.0020**</b>
AG	197 (50.5)	315 (47.2)		1.48 (1.13–1.94)	1.42 (1.03–1.95)	1.32 (1.11–1.58)
GG	110 (28.2)	245 (36.7)				
A	363 (46.5)	529 (39.7)				
G	417 (53.5)	805 (60.3)				
Female						
AA	68 (21.3)	55 (16.5)	<b>0.0122*</b>	<b>0.0038**</b>	<b>0.1173</b>	<b>0.0051**</b>
AG	166 (52.0)	154 (46.3)		1.63 (1.17–2.28)	1.37 (0.92–2.03)	1.37 (1.10–1.70)
GG	85 (26.7)	124 (37.2)				
A	302 (47.3)	264 (39.6)				
G	336 (52.7)	402 (60.4)				

\*  $0.01 \leq p < 0.05$ . \*\*  $p < 0.01$ .

4. Reveille JD, Sims AM, Danoy P, Evans DM, Leo P, Pointon JJ, et al. Genome-wide association study of ankylosing spondylitis identifies non-MHC susceptibility loci. *Nat Genet* 2010;42:123-7.
5. Australo-Anglo-American Spondyloarthritis Consortium (TASC); Wellcome Trust Case Control Consortium 2 (WTCCC2). Interaction between ERAP1 and HLA-B27 in ankylosing spondylitis implicates peptide handling in the mechanism for HLA-B27 in disease susceptibility. *Nat Genet* 2011;43:761-7.
6. Lin Z, Bei JX, Shen M, Li Q, Liao Z, Zhang Y, et al. A genome-wide association study in Han Chinese identifies new susceptibility loci for ankylosing spondylitis. *Nat Genet* 2011;44:73-7.
7. Wei JC, Yen JH, Juo SH, Chen WC, Wang YS, Chiu YC, et al. Association of ORAI1 haplotypes with the risk of HLA-B27 positive ankylosing spondylitis. *PLoS One* 2011;6:e20426.

*J Rheumatol* 2012;39:8; doi:10.3899/jrheum.120040