

Utility of Antiphosphatidylserine/Prothrombin and IgA Antiphospholipid Assays in Systemic Lupus Erythematosus

EHTISHAM AKHTER, ZAKERA SHUMS, GARY L. NORMAN, WALTER BINDER, HONG FANG, and MICHELLE PETRI

ABSTRACT. Objective. Currently, 3 antiphospholipid assays are widely used clinically [lupus anticoagulant (LAC), anticardiolipin (aCL), and anti- β_2 -glycoprotein I (anti- β_2 -GPI)]. LAC is the most specific assay, conferring the highest risk of thrombosis and pregnancy loss, but it cannot be validly performed in an anticoagulated patient. We investigated the usefulness of antiphosphatidylserine/prothrombin (anti-PS/PT) and its association with thrombosis. Anti-PS/PT is strongly associated with the presence of LAC. We also studied the association of IgA antiphospholipid isotypes and specific domains of β_2 -GPI with thrombosis in systemic lupus erythematosus (SLE).

Methods. Stored samples from patients with SLE, with and without past thrombosis, were assayed for antibodies to the whole β_2 -GPI protein (IgG/IgM/IgA), to β_2 -GPI domain 1 (IgG), to β_2 -GPI domain 4/5 (IgA), aCL (IgG/IgM/IgA), and anti-PS/PT (IgG, IgM, and IgG/M). LAC was detected using the dilute Russell's viper venom time (dRVVT) with confirmatory testing.

Results. Anti-PS/PT IgG and IgG/M and anti- β_2 -GPI IgG, IgM, and IgA were highly associated with a history of LAC by dRVVT ($p < 0.0001$). For all thrombosis, of the traditional ELISA assays, anti- β_2 -GPI IgA, IgG, and aCL IgA were most associated. Anti-PS/PT IgG and IgG/M had a similar magnitude of association to the traditional ELISA. For venous thrombosis, of the traditional ELISA, anti- β_2 -GPI (IgG and IgA), anti-PS/PT (IgG and IgG/M), and aCL IgA were associated. Again, anti-PS/PT (IgG and IgG/M) had the same magnitude of association as the traditional ELISA. For stroke, significant association was seen with anti- β_2 -GPI IgA D4/5.

Conclusion. In anticoagulated patients, where LAC testing is not valid, anti-PS/PT, either IgG or IgG/IgM, might serve as useful alternative tests to predict a higher risk of thrombosis. Anti-PS/PT antibodies were associated with all thrombosis and with venous thrombosis. IgA isotypes in secondary antiphospholipid syndrome are associated with thrombosis. Anti- β_2 -glycoprotein domain 1 was not shown to be associated with thrombosis in SLE. (First Release Feb 1 2013; J Rheumatol 2013;40:282–6; doi:10.3899/jrheum.120084)

Key Indexing Terms:

ANTI-PHOSPHATIDYLSERINE/PROTHROMBIN
DOMAIN 4/5 IgA

ANTI- β_2 -GLYCOPROTEIN I
ANTICARDIOLIPIN

The antiphospholipid syndrome (APS) is a hypercoagulable disorder defined by the association of arterial and venous thromboses and/or pregnancy morbidity (fetal loss, premature birth, or recurrent embryonic losses) occurring in the presence of antiphospholipid antibodies (aPL)¹. The commonly used diagnostic tests are the lupus anticoagulant

(LAC) functional coagulation assay, anticardiolipin (aCL), and anti- β_2 -glycoprotein I (anti- β_2 -GPI). Following a workshop in Sapporo, Japan, APS classification criteria were published in 1999¹ and updated in 2006 after another workshop in Sydney, Australia².

APL bind to phospholipid-binding proteins or phospholipid-protein complexes³. We first reported that the LAC was a better predictor of risk of venous thrombosis in SLE than was aCL⁴. In a review of 12 studies, Galli, *et al* found a significant association between thrombosis and LAC⁵.

Prothrombin, another phospholipid-binding protein, was first proposed as a possible cofactor in LAC activity by Loeliger⁶. These antibodies are detected in complex with phosphatidylserine. Bertolaccini, *et al* found antiphosphatidylserine/prothrombin (anti-PS/PT) antibodies in 31% of patients with SLE in general and in 49% of patients with SLE who also had thrombosis⁷. Galli, *et al* found anti-PS/PT in 95% of patients with thrombosis⁸.

From the Johns Hopkins University School of Medicine, Baltimore, Maryland; and Inova Diagnostics Inc., San Diego, California, USA. Supported by NIHAR 43727.

E. Akhter, MD, MPH, Senior Research Program Coordinator, Johns Hopkins University School of Medicine; Z. Shums, MS; G.L. Norman, PhD, Director, Research and Development; W. Binder, PhD, Senior VP, Research and Development, Inova Diagnostics Inc.; H. Fang, MD, MS, Senior Research Program Coordinator, Johns Hopkins University School of Medicine; M. Petri, MD, MPH, Professor of Medicine, Division of Rheumatology, Johns Hopkins University School of Medicine.

Address correspondence to Dr. M. Petri, Division of Rheumatology, Johns Hopkins University School of Medicine, 1830 East Monument Street, Ste. 7500, Baltimore, Maryland 21205, USA. E-mail: mpetri@jhmi.edu Accepted for publication November 30, 2012.

Personal non-commercial use only. The Journal of Rheumatology Copyright © 2013. All rights reserved.

The LAC test cannot be validly performed in an anticoagulated patient. The first aim of our study was to investigate whether anti-PS/PT, which detects most LAC, is associated with thrombosis and with the diluted Russell's viper venom time (dRVVT) assay for the LAC. The second aim was to investigate the association of IgA assays with thrombosis in SLE. Finally, we investigated whether antibodies to specific domains of β_2 -GPI are important in patients with SLE who have APS.

MATERIALS AND METHODS

The Hopkins Lupus Cohort was approved by the Johns Hopkins University School of Medicine Institutional Review Board. All patients gave written informed consent. The study design was cross-sectional. Stored samples from 326 SLE patients with and without past thrombosis were included, of which 164 had a history of any thrombosis, 103 had a history of venous thrombosis, and 53 had a history of stroke. Samples were assayed for anti- β_2 -GPI (IgG/IgM/IgA), anti- β_2 -GPI domain 1 (IgG), anti- β_2 -GPI domain 4/5 (IgA), aCL (IgG/IgM/IgA), and anti-PS/PT (IgG, IgM, and IgG/M). Testing for aCL, anti- β_2 -GPI, and anti-PS/PT was done on 1 sample from each patient. All LAC results were confirmed (by mixing studies and confirmatory testing) using International Society on Thrombosis and Haemostasis guidelines⁹.

Samples were tested for aCL and anti- β_2 -GPI using Quanta Lite ELISA kits (IgG, IgM, and IgA), approved by the US Food and Drug Administration. Antibodies to a complex of PS/PT of the IgG and IgM class were tested by the same kits, as well as by a prototype screening kit (investigational use only) that simultaneously detects both IgG and IgM antibodies to PS/PT. Researchers used only assays specifically detecting IgG antibodies to domain 1 of β_2 -GPI (β_2 -GPI D1) and another specifically detecting IgA antibodies to domains 4 and 5 of β_2 -GPI (β_2 -GPI D4/5).

All ELISA assays were manufactured by Inova Diagnostics Inc. These tests share a common procedure. Patient samples were diluted 1:100 and incubated in the microwell for 30 min. After washing each microwell 3 times, prediluted horseradish peroxidase-labeled conjugate was added and incubated for another 30 min. Wells were washed 3 times and liquid tetramethyl benzidine substrate was added and incubated for 30 min. Color change in microwells was read in a standard microplate reader.

Because no external, internationally recognized controls exist for PS/PT antibody, the results expressed are in arbitrary units, with a cutoff value of 30 units. Establishment of a cutoff for each assay was based on balancing sensitivity and specificity to achieve an optimal clinical utility. In the case of PS/PT, a total of 92 patients with APS, 247 healthy controls, and 43 disease controls were tested. The specificity for PS/PT was set to be high. The cutoff chosen resulted in a specificity of 99% for PS/PT IgG and 98.7% for PS/PT IgM. The determination of whether a test should be tilted toward higher sensitivity or specificity was based on the results of the clinical studies and input from clinical collaborators. In general, receiver-operating characteristics analysis provided a good estimation of an optimum cutoff. However, the final cutoff was adjusted to provide assay results with optimum usefulness for clinicians and their patients. Twenty-four LAC-positive samples were tested for the presence of PS/PT antibodies. All 24 patients were found to be positive for PS/PT IgG and/or PS/PT IgM antibodies. Similarly, the cutoff for anti-D1 IgG and anti-D4/5 IgA was 25 units. The units were derived from studies placing disease versus controls to determine what gave the best specificity and sensitivity. For D1 IgG, the calculations were based on in-house studies showing 97% specificity and 71% sensitivity, and for D4 IgA, specificity of 96%. The cutoffs for anti- β_2 -GPI domain 1 IgG and anti- β_2 -GPI D4/5 IgA were established in a similar manner. Based on clinical specimens, healthy controls, and disease control specimens, a cutoff was chosen to give a specificity of 97% for β_2 -GPI D1 IgG and 96% for β_2 -GPI D4/5 IgA. We did not test any samples with polyclonal B cell activation. The sensitivity

of anti-D4/5 IgA antibodies was 21.3% for any thrombosis, 19.4% for venous thrombosis, and 30.2% for stroke.

Statistical analysis. The p values in all tables were calculated using chi-square tests for the binary variables (for those tests in which 25% had expected counts < 5, Fisher's exact test was used).

RESULTS

The clinical characteristics of 326 patients with SLE are shown in Table 1. Ninety percent of the patients were female, 50% were white, 40% were African American, and 10% were of other ethnicity. The mean age was 45 years. The mean SLE disease duration when the samples were taken was 11 years. The average time interval was 9 ± 8 years between sample collection and all thrombosis, 9 ± 8 years in patients with venous thrombosis, and 8 ± 7 years in patients with stroke. The mean SLE Disease Activity Index and Physician Global Assessment scores were 2.28 and 0.63, respectively. Seventy-eight percent of the patients were taking hydroxychloroquine.

These assays were highly associated with a history of LAC by dRVVT ($p < 0.0001$): anti-PS/PT IgG, IgM, and IgG/M, anti- β_2 -GPI IgG, IgM, and IgA. Anti- β_2 -GPI D1 was also highly associated with a history of LAC. IgG anti- β_2 -GPI and anti-PS/PT had the highest odds of being positive when LAC was present (IgG anti- β_2 -GPI, OR 20.3, 95% CI 5.7, 72.1; IgG anti-PS/PT, OR 14.7, 95% CI 6.3, 34.7; Table 2).

Thrombosis. For all thrombosis, anti- β_2 -GPI (IgA and IgG) and aCL IgA were the most highly associated (Table 3). Out of 164 patients, 134 had aPL positivity on the followup visit. Seventy-eight were positive for anticardiolipin and 66 were positive for LAC. Fifty-four patients had both aCL and LAC.

Table 1. Demographic characteristics (n = 326). Data are mean (SD) unless otherwise indicated.

Characteristics	
Age, yrs	45.7 (14.0)
Sex, n (%)	
Female	298 (91.4)
Male	28 (8.6)
Ethnicity, n (%)	
White	163 (50.0)
African-American	131 (40.2)
Other	32 (9.8)
Measures of disease activity	
Physician global assessment (0 to 3 VAS)	0.63 (0.66)
SELENA-SLEDAI	2.28 (3.12)
Serologies	
C3, mg/dl	115.7 (35.7)
C4, mg/dl	22.1 (10.6)
Anti-dsDNA ≥ 10 , n (%)	67 (20.6)
Medications	
Prednisone, mg	4.17 (7.87)
Immunosuppressive, n (%)	117 (35.9)
Hydroxychloroquine, n (%)	254 (77.9)

VAS: visual analog scale; SELENA-SLEDAI: Safety of Estrogens in Lupus Erythematosus National Assessment-Systemic Lupus Erythematosus Disease Activity Index.

For venous thrombosis, anti- β_2 -GPI IgG, IgA, anti-PS/PT (IgG and IgG/M), and aCL IgA were significantly associated (Table 4).

Stroke. For stroke, a significant association was seen with anti- β_2 -GPI IgA D4/5 (Table 5), but not with the presence of other aPL.

DISCUSSION

Our study shows evidence for an association of anti-PS/PT assays with LAC in SLE, as expected. Four previous studies found an association of anti-PS/PT and LAC^{10,11,12,13}. One study failed to find a significant association¹⁴. The lack of correlation in the one study may exist because the APS patient population had experienced much pregnancy loss. Only 13 of the 62 patients with APS had thrombosis; the rest were patients with recurrent pregnancy loss¹⁴.

In our study, we found that anti-PS/PT was significantly associated with all thrombosis and with venous thrombosis.

A previous study found an association of anti-PS/PT (IgG and IgG/M) with all thrombosis in APS¹¹. A study of 175 patients with SLE showed that the presence of anti-PS/PT antibodies conferred an OR of 9.6 for arterial thrombosis and 7.33 for venous thrombosis¹³. One study found an association ($p = 0.009$) with anti-PT (not anti-PS/PT) and venous thrombosis in SLE¹⁵. In contrast, another study of 22 patients with APS found no association of anti-PT antibodies with thrombosis¹⁶. A study of 175 patients with SLE failed to find a significant association with venous thrombosis¹⁷. Our results are in concordance with a similar study by Bertolaccini, *et al*⁷. However, we also found that anti-PS/PT IgG and IgG/M were significantly associated with a history of LAC. We could not find any significant association with stroke in our cohort.

Of particular note was the importance of IgA APS assays for both all thrombosis and venous thrombosis. Studies of IgA aCL in SLE have been conflicting. Two studies found

Table 2. Association of antiphospholipid assays with a history of lupus anticoagulant (LAC).

Assay		LAC-positive, n = 71 No. (% positive)	LAC-negative n = 212 No. (% positive)	OR (95% CI)	p
Anti-PS/PT	IgM	35 (49)	16 (8)	11.9 (6.0, 23.7)	< 0.0001
	IgG	26 (37)	8 (4)	14.7 (6.3, 34.7)	< 0.0001
	IgG/M	34 (48)	16 (8)	11.3 (5.6, 22.4)	< 0.0001
Anticardiolipin	IgM	17 (24)	10 (5)	6.4 (2.8, 14.7)	< 0.0001
	IgG	20 (28)	9 (4)	8.8 (3.8, 20.6)	< 0.0001
	IgA	10 (14)	0 (0)		< 0.0001
Anti- β_2 -GPI	IgM	10 (14)	4 (2)	8.5 (2.6, 28.1)	< 0.0001
	IgG	16 (23)	3 (1)	20.3 (5.7, 72.1)	< 0.0001
	IgA	29 (41)	30 (14)	4.2 (2.3, 7.7)	< 0.0001
Anti- β_2 -GPI D1	IgG	9 (13)	5 (2)	6.0 (1.9, 18.6)	0.0005
Anti- β_2 -GPI D4/5	IgA	21 (30)	35 (17)	2.1 (1.1, 4.0)	0.0167

Anti-PS/PT: antiphosphatidylserine/prothrombin; anti- β_2 -GPI: anti- β_2 -glycoprotein I; anti- β_2 -GPI D1: anti- β_2 -glycoprotein I domain 1; anti- β_2 -GPI D4/5: anti- β_2 -glycoprotein I domain 4/5.

Table 3. Association of antiphospholipid assays with all thrombosis (venous or arterial).

Assay		Thrombosis, n = 164 No. (% positive)	No Thrombosis, n = 162 No. (% positive)	OR 95% CI*	p (adjusted)*
IgG	Anti- β_2 -GPI	18 (11)	6 (4)	3.3 (1.2, 8.9)	0.021
	Anti- β_2 -GPI D1	11 (7)	9 (6)	1.1 (0.4, 2.9)	0.79
	aCL	22 (13)	16 (10)	1.4 (0.7, 2.9)	0.34
IgM	Anti-PS/PT	26 (16)	14 (9)	2.2 (1.0, 4.8)	0.045
	Anti- β_2 -GPI	8 (5)	9 (6)	0.8 (0.3, 2.3)	0.69
	aCL	19 (12)	14 (9)	1.3 (0.6, 2.8)	0.47
IgA	Anti-PS/PT	43 (26)	25 (15)	1.9 (1.1, 3.3)	0.033
	Anti- β_2 -GPI	48 (29)	24 (15)	2.4 (1.3, 4.2)	0.0028
	aCL	11 (7)	1 (1)	9.5 (1.2, 75.8)	0.034
IgG/M	Anti- β_2 -GPI D4/5	35 (21)	28 (17)	1.3 (0.7, 2.3)	0.38
	Anti-PS/PT	38 (23)	24 (15)	1.7 (0.9, 3.1)	0.087
LAC**	dRVVT	52 (37)	19 (13)	3.8 (2.0, 7.3)	< 0.0001

* Adjusted for age, sex, and ethnicity. ** Forty-three patients did not have dilute Russell viper venom test (dRVVT) results available. Thus, the total number of patients for the thrombosis group was 44, and the no-thrombosis group, 239. Anti- β_2 -GPI: anti- β_2 -glycoprotein I; anti- β_2 -GPI D1: anti- β_2 -glycoprotein I domain 1; aCL: anticardiolipin; anti-PS/PT: antiphosphatidylserine/prothrombin; LAC: lupus anticoagulant; anti- β_2 -GPI D4/5: anti- β_2 -glycoprotein I domain 4/5.

Table 4. Association of antiphospholipid assays with venous thrombosis.

Assay		Thrombosis, n = 103 No. (% positive)	No Thrombosis, n = 223 No. (% positive)	OR 95% CI*	p (adjusted)*
IgG	Anti- β_2 -GPI	15 (15)	9 (4)	3.7 (1.5, 9.1)	0.0048
	Anti- β_2 -GPI D1	9 (9)	11 (5)	1.7 (0.7, 4.2)	0.29
	aCL	17 (17)	21 (9)	1.7 (0.8, 3.5)	0.14
IgM	Anti-PS/PT	21 (20)	19 (9)	2.7 (1.3, 5.8)	0.0082
	Anti- β_2 -GPI	6 (6)	11 (5)	1.1 (0.4, 3.1)	0.88
	aCL	13 (13)	20 (9)	1.4 (0.7, 3.1)	0.36
IgA	Anti-PS/PT	27 (26)	41 (18)	1.5 (0.8, 2.6)	0.19
	Anti- β_2 -GPI	31 (30)	41 (18)	1.9 (1.1, 3.3)	0.024
	aCL	8 (8)	4 (2)	4.3 (1.2, 14.8)	0.023
IgG/M	Anti- β_2 -GPI D4/5	20 (19)	43 (19)	1.0 (0.6, 1.9)	0.97
IgG/M	Anti-PS/PT	29 (28)	33 (15)	2.1 (1.2, 3.8)	0.015
LAC**	dRVVT	36 (41)	35 (18)	3.3 (1.8, 6.0)	< 0.0001

* Adjusted for age, sex, and ethnicity. ** Forty-three patients did not have dilute Russell viper venom test (dRVVT) results available. Thus, the total number of patients for the thrombosis group was 88, and the no-thrombosis group, 195. Anti- β_2 -GPI: anti- β_2 -glycoprotein I; anti- β_2 -GPI D1: anti- β_2 -glycoprotein I domain 1; aCL: anticardiolipin; anti-PS/PT: antiphosphatidylserine/prothrombin; LAC: lupus anticoagulant; anti- β_2 -GPI D4/5: anti- β_2 -glycoprotein I domain 4/5.

Table 5. Association of antiphospholipid assays with stroke.

Assay		Stroke, n = 53 No. (% positive)	No Stroke, n = 273 No. (% positive)	OR 95% CI*	p (adjusted)*
IgG	Anti- β_2 -GPI	5 (9)	19 (7)	2.0 (0.7, 6.2)	0.21
	Anti- β_2 -GPI D1	3 (6)	17 (6)	1.1 (0.3, 4.1)	0.89
	aCL	7 (13)	31 (11)	1.6 (0.6, 4.1)	0.33
IgM	Anti-PS/PT	8 (15)	32 (12)	2.5 (1.0, 6.3)	0.062
	Anti- β_2 -GPI	2 (4)	15 (5)	0.7 (0.2, 3.3)	0.66
	aCL	8 (15)	25 (9)	1.6 (0.7, 4.1)	0.28
IgA	Anti-PS/PT	14 (26)	54 (20)	1.6 (0.8, 3.2)	0.21
	Anti- β_2 -GPI	17 (32)	55 (20)	1.9 (1.0, 3.7)	0.070
	aCL	4 (8)	8 (3)	2.0 (0.6, 7.4)	0.28
IgG/M	Anti- β_2 -GPI D4/5	16 (30)	47 (17)	2.2 (1.1, 4.4)	0.033
IgG/M	Anti-PS/PT	11 (21)	51 (19)	1.5 (0.7, 3.2)	0.35
LAC**	dRVVT	17 (39)	54 (23)	2.3 (1.1, 4.8)	0.022

* Adjusted for age, sex, and ethnicity. ** Forty-three patients did not have dilute Russell viper venom test (dRVVT) results available. Thus, the total number of patients for the thrombosis group was 44, and the no-thrombosis group, 239. Anti- β_2 -GPI: anti- β_2 -glycoprotein I; anti- β_2 -GPI D1: anti- β_2 -glycoprotein I domain 1; aCL: anticardiolipin; anti-PS/PT: antiphosphatidylserine/prothrombin; LAC: lupus anticoagulant; anti- β_2 -GPI D4/5: anti- β_2 -glycoprotein I domain 4/5.

an association with all thrombosis^{18,19}, whereas 3 studies did not^{20,21,22}. One study found a statistically significant association of IgA aCL with thrombosis and thrombocytopenia ($p = 0.02$)¹⁹. Similarly, no consensus has been reached about the association of IgA anti- β_2 -GPI and thrombosis in SLE. Several studies, including our own previous investigations^{23,24,25}, have found an association with venous thrombosis, but others have not^{26,27}. In one study, a borderline association was seen in patients with arterial thrombosis ($p = 0.06$)²².

To our knowledge, ours is the first study of stroke in SLE to show an association with D4/5 IgA. Two past non-SLE studies have suggested such an association^{28,29}. IgA anti- β_2 -GPI targets D4 of the β_2 -GPI molecule in non-SLE patients and was associated with atherosclerosis (acute

myocardial infarction, acute coronary syndrome, and peripheral artery disease)²⁸. IgA anti- β_2 -GPI was present in 22% of non-SLE patients with ischemic stroke²⁹. Interestingly, D4/5 IgA was not associated with venous thrombosis.

In contrast to a study by de Laat, *et al*³⁰, our study showed no association of IgG anti- β_2 -GPI D1 with thrombosis. In the de Laat study, anti-D1 IgG was significantly correlated with a history of thrombosis. However, that study did not differentiate between patients with APS and those with SLE (i.e., results were based on any thrombosis in either population). Our study specifically included only patients with SLE. Future studies will seek to clarify the differing results observed between these 2 studies.

Anti-PS/PT, either IgG or IgG/IgM, offers clinicians an

alternative test to predict increased risk of thrombosis in anticoagulated patients where LAC measurement is not reliable. IgA isotype aPL antibodies are highly associated with thrombosis in SLE. For stroke, only D4/5 IgA was associated. These novel assays will require validation in additional cohorts.

REFERENCES

- Wilson WA, Gharavi AE, Koike T, Lockshin MD, Branch DW, Piette JC, et al. International consensus statement on preliminary classification criteria for definite antiphospholipid syndrome: Report of an international workshop. *Arthritis Rheum* 1999;42:1309–11.
- Miyakis S, Lockshin MD, Atsumi T, Branch DW, Brey RL, Cervera R, et al. International consensus statement on an update of the classification criteria for definite antiphospholipid syndrome (APS). *J Thromb Haemost* 2006;4:295–306.
- Roubey RA. Autoantibodies to phospholipid-binding plasma proteins: A new view of lupus anticoagulants and other “antiphospholipid” antibodies. *Blood* 1994;84:2854–67.
- Petri M, Rheinschmidt M, Whiting-O’Keefe Q, Hellmann D, Corash L. The frequency of lupus anticoagulant in systemic lupus erythematosus. A study of sixty consecutive patients by activated partial thromboplastin time, Russell viper venom time, and anticardiolipin antibody level. *Ann Intern Med* 1987;106:524–31.
- Galli M, Luciani D, Bertolini G, Barbui T. Lupus anticoagulants are stronger risk factors for thrombosis than anticardiolipin antibodies in the antiphospholipid syndrome: A systematic review of the literature. *Blood* 2003;101:1827–32.
- Loeliger A. Prothrombin as a cofactor of the circulating anticoagulant in systemic lupus erythematosus. *Thromb Diath Haemorrh* 1959;3:237–56.
- Bertolaccini ML, Atsumi T, Koike T, Hughes GR, Khamashta MA. Antiprothrombin antibodies detected in two different assay systems. Prevalence and clinical significance in systemic lupus erythematosus. *Thromb Haemost* 2005;93:289–97.
- Galli M, Beretta G, Daldossi M, Bevers EM, Barbui T. Different anticoagulant and immunological properties of antiprothrombin antibodies in patients with antiphospholipid antibodies. *Thromb Haemost* 1997;77:486–91.
- Petri M, Nelson L, Weimer F, Anderson D, Darlington T, Corash L. The automated modified Russell viper venom time test for lupus anticoagulant. *J Rheumatol* 1991;18:1823–5.
- Matsuda J, Saitoh N, Gotoh M, Kawasugi K, Gohchi K, Tsukamoto M. Phosphatidyl serine-dependent antiprothrombin antibody is exclusive to patients with lupus anticoagulant. *Br J Rheumatol* 1996;35:589–91.
- Atsumi T, Ieko M, Bertolaccini ML, Ichikawa K, Tsutsumi A, Matsuura E, et al. Association of autoantibodies against phosphatidylserine-prothrombin complex with manifestations of the antiphospholipid syndrome and with the presence of lupus anticoagulant. *Arthritis Rheum* 2000;43:1982–93.
- Ieko M, Nakabayashi T, Tarumi T, Yoshida M, Naito S, Atsumi T, et al. [Phosphatidylserine-dependent anti-prothrombin antibody as a new marker for the diagnosis of antiphospholipid syndrome. Article in Japanese.] *Rinsho Byori* 2006;54:256–62.
- Nojima J, Iwatani Y, Suehisa E, Kuratsune H, Kanakura Y. The presence of anti-phosphatidylserine/prothrombin antibodies as risk factor for both arterial and venous thrombosis in patients with systemic lupus erythematosus. *Haematologica* 2006;91:699–702.
- Jaskowski TD, Wilson AR, Hill HR, Branch WD, Tebo AE. Autoantibodies against phosphatidylserine, prothrombin and phosphatidylserine-prothrombin complex: Identical or distinct diagnostic tools for antiphospholipid syndrome? *Clin Chim Acta* 2009;410:19–24.
- Puurunen M, Vaarala O, Julkunen H, Aho K, Palosuo T. Antibodies to phospholipid-binding plasma proteins and occurrence of thrombosis in patients with systemic lupus erythematosus. *Clin Immunol Immunopathol* 1996;80:16–22.
- Pengo V, Biasiolo A, Brocco T, Tonetto S, Ruffatti A. Autoantibodies to phospholipid-binding plasma proteins in patients with thrombosis and phospholipid-reactive antibodies. *Thromb Haemost* 1996;75:721–4.
- Horbach DA, van Oort E, Donders RC, Derksen RH, de Groot PG. Lupus anticoagulant is the strongest risk factor for both venous and arterial thrombosis in patients with systemic lupus erythematosus. Comparison between different assays for the detection of antiphospholipid antibodies. *Thromb Haemost* 1996;76:916–24.
- Kalunian KC, Peter JB, Middlekauff HR, Sayre J, Ando DG, Mangotich M, et al. Clinical significance of a single test for anti-cardiolipin antibodies in patients with systemic lupus erythematosus. *Am J Med* 1988;85:602–8.
- Lopez LR, Santos ME, Espinoza LR, La Rosa FG. Clinical significance of immunoglobulin A versus immunoglobulins G and M anti-cardiolipin antibodies in patients with systemic lupus erythematosus. Correlation with thrombosis, thrombocytopenia, and recurrent abortion. *Am J Clin Pathol* 1992;98:449–54.
- Weidmann CE, Wallace DJ, Peter JB, Knight PJ, Bear MB, Klinenberg JR. Studies of IgG, IgM and IgA antiphospholipid antibody isotypes in systemic lupus erythematosus. *J Rheumatol* 1988;15:74–9.
- Merkel PA, Chang Y, Pierangeli SS, Convery K, Harris EN, Polissone RP. The prevalence and clinical associations of anticardiolipin antibodies in a large inception cohort of patients with connective tissue diseases. *Am J Med* 1996;101:576–83.
- Escalante A, Brey RL, Mitchell BD Jr, Dreiner U. Accuracy of anticardiolipin antibodies in identifying a history of thrombosis among patients with systemic lupus erythematosus. *Am J Med* 1995;98:559–65.
- Lakos G, Kiss E, Regéczy N, Tarján P, Soltész P, Zeher M, et al. Isotype distribution and clinical relevance of anti-beta 2-glycoprotein I (beta 2-GPI) antibodies: Importance of IgA isotype. *Clin Exp Immunol* 1999;117:574–9.
- Lee SS, Cho ML, Joo YS, Kim WU, Hong YS, Min JK, et al. Isotypes of anti-beta 2-glycoprotein I antibodies: Association with thrombosis in patients with systemic lupus erythematosus. *J Rheumatol* 2001;28:520–4.
- Mehrani T, Petri M. Association of IgA anti-beta 2 glycoprotein I with clinical and laboratory manifestations of systemic lupus erythematosus. *J Rheumatol* 2011;38:64–8.
- Samarkos M, Davies KA, Gordon C, Loizou S. Clinical significance of IgA anticardiolipin and anti-beta 2-GPI antibodies in patients with systemic lupus erythematosus and primary antiphospholipid syndrome. *Clin Rheumatol* 2006;25:199–204.
- Bertolaccini ML, Atsumi T, Escudero Contreras A, Khamashta MA, Hughes GR. The value of IgA antiphospholipid testing for diagnosis of antiphospholipid (Hughes) syndrome in systemic lupus erythematosus. *J Rheumatol* 2001;28:2637–43.
- Iverson GM, von Mühlen CA, Staub HL, Lassen AJ, Binder W, Norman GL. Patients with atherosclerotic syndrome, negative in anti-cardiolipin assays, make IgA autoantibodies that preferentially target domain 4 of beta 2-GPI. *J Autoimmun* 2006;27:266–71.
- Staub HL, Norman GL, Crowther T, da Cunha VR, Polanczyk A, Bohn JM, et al. Antibodies to the atherosclerotic plaque components beta 2-glycoprotein I and heat-shock proteins as risk factors for acute cerebral ischemia. *Arq Neuropsiquiatr* 2003;61:757–63.
- de Laat B, Pengo V, Pabinger I, Musial J, Voskuyl AE, Bultink IE, et al. The association between circulating antibodies against domain 1 of beta 2-glycoprotein and thrombosis: An international multicenter study. *J Thromb Haemost* 2009;7:1767–73.