

# Fibromyalgia and the Risk of a Subsequent Motor Vehicle Crash

Donald A. Redelmeier, Jeremy D. Zung, Deva Thiruchelvam, and Robert J. Tibshirani

**ABSTRACT. Objective.** Motor vehicle crashes are a widespread contributor to mortality and morbidity, sometimes related to medically unfit motorists. We tested whether patients diagnosed with fibromyalgia (FM) have an increased risk of a subsequent serious motor vehicle crash.

**Methods.** We conducted a population-based self-matched longitudinal cohort analysis to estimate the incidence rate ratio of crashes among patients diagnosed with FM relative to the population norm in Ontario, Canada. We included adults diagnosed from April 1, 2006, to March 31, 2012, excluding individuals younger than 18 years, living outside Ontario, lacking valid identifiers, or having only a single visit for the diagnosis. The primary outcome was an emergency department visit as a driver involved in a motor vehicle crash.

**Results.** The patients (n = 137,631) accounted for 738 crashes during the first year of followup after diagnosis, equal to an incidence rate ratio of 2.44 compared with the population norm (95% CI 2.27–2.63, p < 0.001). The crash rate was more than twice the population norm for those with a new or a persistent diagnosis. The increased risk included patients with diverse characteristics, approached the rate observed among other patients diagnosed with alcoholism, and was mitigated among those who received dedicated FM care or a physician warning for driving safety.

**Conclusion.** A diagnosis of FM is associated with an increased risk of a subsequent motor vehicle crash that might justify medical interventions for traffic safety. (First Release May 15 2015; J Rheumatol 2015;42:1502–10; doi:10.3899/jrheum.141315)

*Key Indexing Terms:*

TRAFFIC ACCIDENT  
DRIVING ASSESSMENT

MOTOR VEHICLE CRASH  
NONMALIGNANT PAIN

FITNESS TO DRIVE  
DRIVER COMPENSATION

Fibromyalgia (FM) affects more than 1 million Americans and is characterized by chronic pain, tender points, and fluctuating severity<sup>1</sup>. The condition is associated with fatigue, nonrestorative sleep, cognitive dysfunction, headaches, anxiety, depression, joint stiffness, dizziness, atypical chest

pain, and other symptoms that might interfere with the safe driving of a motor vehicle<sup>2</sup>. Treatment may include exercise, behavioral therapy, analgesia, antidepressants, diet change, sedatives, and anticonvulsants<sup>3</sup>. No blood test or imaging scan can routinely diagnose the condition, gauge case severity, or monitor a patient's course<sup>4</sup>. FM is an example of an imperfectly understood condition associated with substantial patient suffering and medical uncertainty<sup>5</sup>.

Motor vehicle crashes can be common complications for patients with chronic medical conditions. The blatant damage of a serious crash provides the rationale for health prevention, including warnings by physicians to patients regarding motor vehicle safety<sup>6</sup>. Specific diseases can contribute to crash risk by causing sudden unconsciousness (e.g., narcolepsy), temporary impairment (e.g., chemotherapy), or chronic disability (e.g., dementia). In turn, medical authorities in the United States and elsewhere have developed guidelines for assessing a patient's fitness to drive<sup>7,8</sup>. No guidelines, however, address patients with FM although some cautions appear for patients with pain or impaired functional status.

Prior research has investigated whether FM can be a consequence of a past motor vehicle crash; however, the literature contains no data on whether FM could be a contributor to a future motor vehicle crash<sup>9</sup>. We hypothesized that FM might be associated with limitations that compromise a patient's driving. Neck pain or opioid medications, for

*From the Department of Medicine, University of Toronto; Evaluative Clinical Sciences Platform, Sunnybrook Research Institute; Institute for Clinical Evaluative Sciences in Ontario; Division of General Internal Medicine, Sunnybrook Health Sciences Centre; Center for Leading Injury Prevention Practice Education and Research, Toronto, Ontario, Canada; Department of Statistics, Stanford University, Stanford, California, USA.*

*Supported by a Canada Research Chair in Medical Decision Sciences, the Canadian Institutes of Health Research, the Physicians' Services Incorporated Foundation, and a D+H SRI Summer Student award.*

*D.A. Redelmeier, MD, FRCPC, MSHSR, FACP, Department of Medicine, University of Toronto, and the Evaluative Clinical Sciences Platform, Sunnybrook Research Institute, and the Institute for Clinical Evaluative Sciences in Ontario, and Division of General Internal Medicine, Sunnybrook Health Sciences Centre, and the Center for Leading Injury Prevention Practice Education and Research; J.D. Zung, BSc, Department of Medicine, University of Toronto, and the Evaluative Clinical Sciences Platform, Sunnybrook Research Institute, and the Institute for Clinical Evaluative Sciences in Ontario; D. Thiruchelvam, MSc, Evaluative Clinical Sciences Platform, Sunnybrook Research Institute, and the Institute for Clinical Evaluative Sciences in Ontario; R.J. Tibshirani, PhD, Department of Statistics, Stanford University.*

*Address correspondence to Dr. D.A. Redelmeier, Sunnybrook Health Sciences Centre, G-151, 2075 Bayview Ave., Toronto, Ontario M4N 3M5, Canada. E-mail: dar@ices.on.ca*

*Accepted for publication March 24, 2015.*

example, may interfere with a motorist's ability to check blind spots or react quickly<sup>10,11</sup>. Moreover, the chronic subjective varying nature of FM might provide insufficient objective feedback for self-regulation of driving<sup>12</sup>. Our study question was: "Are drivers diagnosed with FM more prone to motor vehicle crashes than the population norm?"

## MATERIALS AND METHODS

**Population setting.** Ontario in 2006 had a population of about 9,949,480 adults, of whom 8,867,965 were licensed drivers involved in a total of 216,247 police-reported crashes<sup>13,14</sup>. We defined a serious motor vehicle crash as an incident resulting in the driver going to a hospital emergency department, thereby excluding crashes that resulted in minor injury or vehicle damage only. This definition also excluded cases where the patient was a passenger or a pedestrian. For the population of Ontario in 2006, these criteria amounted to 19,507 crashes that sent the driver to a hospital emergency department, equal to an annual event rate of 2.20 serious motor vehicle crashes per 1000 drivers in the general population ( $19,507 \div 8,867,965 \times 1000$ ).

Ontario on April 1, 2006, was distinctive in introducing an additional physician fee (code K037) for dedicated special care of patients diagnosed with FM<sup>15,16,17</sup>. This was a comprehensive fee code encompassing all common and specific elements of medical care (\$51.70) for this diagnosis<sup>18</sup>. The intent was to provide priority attention for special situations, analogous to physician fee K039 for dedicated smoking cessation counseling<sup>19,20</sup>. Regardless of dedicated special care for FM, patients had free access to emergency care under universal health insurance and could be tracked over time through validated population-based databases covering all 174 full-service emergency departments in the region<sup>21</sup>.

**Patient selection and diagnosis date.** We identified adults diagnosed with FM between April 1, 2006, and March 31, 2012, and excluded individuals younger than age 18 years, living outside Ontario, or lacking valid identifiers under universal health insurance. The 6-year interval was selected to provide all data following the introduction of the dedicated care program and to ensure a minimum 1-year followup for all patients. FM was defined as a physician diagnosis coded by the International Classification of Disease 9th Revision (integer code: 729)<sup>22</sup>. To reduce false-positive errors, we excluded patients with only a single physician visit for the diagnosis, in accord with past research<sup>23</sup>. These codes had a sensitivity of 62% to 85% and a specificity of 94% to 99% when validated for other disorders<sup>24,25,26,27</sup>. The diagnosis of FM, therefore, is prone to false positives, false negatives, and inconsistencies over time<sup>28</sup>.

FM often has a gradual onset and delayed diagnosis so that the onset is also uncertain. We defined the start of followup for each patient (time-zero) according to the date of a second (confirmed) diagnosis of FM following April 1, 2006. Patients who had 2 or more visits for FM during the prior 5 years were defined as having an ongoing persistent diagnosis (prevalent cases). The remaining patients who had no other visits for FM during earlier years were defined as having an initial new diagnosis (incident cases). The distinction between a persistent and a new diagnosis served as our pre-specified subgroup analysis to examine patients with differing disease durations. The available databases lacked data on diagnostic certainty and severity.

**Patient characteristics.** Additional diagnoses that are often considered and sometimes misdiagnosed as FM were ascertained from physician billing data for the full year prior to diagnosis<sup>29</sup>. Psychiatric conditions included depression, anxiety, alcoholism, substance abuse, and posttraumatic stress disorder (codes 296, 300, 303, 304, 305, 308, 309). Rheumatic disorders included systemic lupus erythematosus, rheumatoid arthritis, ankylosing spondylitis, and polymyalgia rheumatica (codes 710, 712, 714, 720, 725, 737). General medical comorbidities included migraine headaches, irritable bowel disorder, hypothyroidism, multiple sclerosis, breast cancer, and endometriosis (codes 042, 043, 044, 070, 174, 244, 340, 346, 564, 617).

Acute injuries included fractures, dislocations, sprains, concussions, open wounds, lacerations, contusions, abrasions, and burns (codes 800 through 859).

Patient age was obtained from the population registry, as were data on sex and home location (rural, urban). Prior hospitalizations, emergency visits, and outpatient encounters for any reason were ascertained for the year prior to diagnosis based on validated identifiers. We also identified a clinical intervention likely to change subsequent traffic risks; namely, a medical warning by a physician for fitness to drive (code K035)<sup>30</sup>. Similarly, we identified a diagnostic procedure unlikely to change subsequent traffic risks; namely, an electrocardiogram (EKG; code G313). The databases did not contain information on driving record, distance traveled, traffic infractions, vehicle ownership, use of medications, or functional status.

**Motor vehicle crashes.** We identified motor vehicle crashes resulting in an emergency visit to any hospital in the region, representing all data available for 1 year of observation before and after the diagnosis date for each patient. We focused on emergencies characterized as a crash using the International Classification of Diseases diagnostic codes (codes V20 to V69)<sup>31</sup>. These included individuals involved as a driver in a motor vehicle crash and excluded emergencies where the patient was a passenger, pedestrian, or other road user. These codes were an exact match with medical records (3 character precision) in about 94% of cases when validated in a past study of diagnostic accuracy<sup>32</sup>. These methods have been further validated in prior research focusing on specific subgroups of motorists<sup>33,34</sup>.

**Primary and secondary comparisons.** The high frequency of motor vehicle crashes allowed for several comparison analyses. Our primary comparison evaluated the risk of a crash among patients with FM relative to the entire driving population of the region. In secondary analyses, we also repeated the primary analysis by selecting patients diagnosed with alcoholism (code 303) instead of patients diagnosed with FM. Similarly, we also repeated the analysis by selecting patients diagnosed with an arm fracture (code 812) instead of patients diagnosed with FM. The intent of these 2 secondary analyses was to assess crash risks for patients in whom a major increase in risk was anticipated (alcoholism) and for patients where no major increase was anticipated (arm fracture)<sup>35</sup>.

**Statistical analysis.** The primary analysis evaluated emergency department visits for drivers in motor vehicle crashes and focused on the year following diagnosis. The absolute risks following diagnosis were summarized using incidence rate ratios relative to the general population of drivers in Ontario, along with 95% CI<sup>36</sup>. Graphical displays of the distribution of motor vehicle crashes were computed for the full year preceding and following diagnosis. The entire analysis was then replicated for patients diagnosed with alcoholism (positive control) and patients diagnosed with an arm fracture (negative control). Together, these analyses assessed the patients' driving risk compared with population norms, their own baselines, and positive or negative controls.

Prespecified subgroup analyses examined patients with an incident diagnosis of FM separately from patients with a persistent diagnosis of FM. An additional prespecified subgroup analysis evaluated patients accrued during the first 2 years of the study (to allow an extended 5-yr followup interval after diagnosis). All other patient characteristics were tested by subgroup analyses to check robustness. The estimated effectiveness of each of the 3 clinical interventions (dedicated special care for FM, physician warning about fitness to drive, diagnostic EKG) were each evaluated using nested exposure-crossover analyses<sup>37</sup>. The study protocol was approved by the Sunnybrook Research Ethics Board, including a waiver for individual consent.

## RESULTS

**Descriptive overview.** A total of 137,631 patients were identified, of whom about two-thirds were new cases and one-third were persistent cases. Patients in both groups spanned a broad range of ages and socioeconomic status

(Table 1). The median patient was a 52-year-old woman (a demographic profile that would otherwise predict crash risks below the population norm). Few patients had a history of an additional rheumatic, psychiatric, or general medical comorbidity (yet more than the population norm)<sup>38</sup>. Patients with persistent FM tended to be older, more likely to have a psychiatric diagnosis, and less likely to have an acute injury than patients with a new diagnosis of FM.

**Motor vehicle crashes.** The patients accounted for 1566 serious motor vehicle crashes during the year before diagnosis. This was equal to 11.38 events per 1000 individuals annually and demonstrated a strong reciprocal association where many crashes preceded the diagnosis of FM (Figure 1). The patients also accounted for an additional 738 serious motor vehicle crashes during the subsequent year following diagnosis. This subsequent rate was equal to 5.36 events per 1000 patients annually and was more than twice the population norm. The observed risk during followup equaled an incidence rate ratio of 2.44 (95% CI 2.27–2.63,  $p < 0.001$ ) and a net increase of 438 crashes compared with the population norm.

**Control patients.** During the same interval, 75,617 other patients were diagnosed with alcoholism and 40,753 other patients were diagnosed with an arm fracture. Those with alcoholism accounted for 395 crashes during the first year following their diagnosis (5.22 events per 1000 patients

annually). The observed risk with alcoholism equaled an incidence rate ratio of 2.38 (95% CI 2.15–2.63) and a net increase of 229 crashes compared with the population norm. Those with an arm fracture accounted for 98 crashes during the first year following their diagnosis (2.40 events per 1000 patients annually). The observed risk with an arm fracture equaled an incidence rate ratio of 1.10 (95% CI 0.90–1.34) and no net increase compared with the population norm.

**Individual characteristics.** The increased risk associated with FM was consistent for patients with different characteristics. Each subgroup had a risk more than twice the population norm, except for those who were older than age 60 years, free of comorbid diagnoses, having few outpatient clinic visits, or devoid of prior emergency visits (Table 2). No subgroup had a risk within the population norm. Patients with an emergency visit in the prior year had risks exceeding 3 times the population norm. The increased risk was observed regardless of whether FM was a new or a persistent diagnosis, was sustained for 5 years in the subgroup of patients with extended followup (Figure 2), and was broadly relevant with fewer than 6 patients contributing more than 1 crash in 1 year.

**Dedicated FM care.** A large subgroup ( $n = 29,007$ ) received dedicated special care for FM. These patients had a high baseline risk amounting to 854 crashes during the 3-year interval preceding special care (9.81 events per 1000 patients annually). Similarly, the patients accounted for 198 crashes

Table 1. Patient characteristics ( $n = 137,631$ ). Values are  $n$  (%).

Characteristics		New Diagnosis, $n = 89,349$	Persistent Diagnosis, $n = 48,282$
Age, yrs	≤ 39	19,435 (22)	6375 (13)
	40–49	21,847 (24)	11,382 (24)
	50–59	21,516 (24)	14,155 (29)
	≥ 60	26,551 (30)	16,370 (34)
Sex	Female	60,405 (68)	35,541 (74)
	Male	28,944 (32)	12,741 (26)
Home location	Urban	77,878 (87)	43,145 (89)
	Rural	11,441 (13)	5110 (11)
	Missing	30 (0)	27 (0)
Socioeconomic status	Highest	15,579 (17)	8509 (18)
	Next to highest	17,367 (19)	9694 (20)
	Middle	17,228 (19)	9491 (20)
	Next to lowest	18,740 (21)	10,092 (21)
	Lowest	20,102 (22)	10,377 (21)
Comorbidity*	Missing	333 (0)	119 (0)
	Rheumatic	5614 (6)	2819 (6)
	Psychiatric	28,182 (32)	16,811 (35)
	General medical	11,824 (13)	7096 (15)
Total care*	Acute trauma	27,805 (31)	12,885 (27)
	Outpatient clinic ≥ 7	71,753 (80)	39,913 (83)
	Emergency department ≥ 1	34,971 (39)	16,742 (35)
Cohort entry**	Hospital admission ≥ 1	9712 (11)	4054 (8)
	Initial yrs	22,755 (25)	24,923 (52)
	Subsequent yrs	66,594 (75)	23,359 (48)

\* Physician diagnosis during prior year. \*\* Initial years for patients identified 2006–2007, subsequent years for patients identified 2008–2012.








Figure 3. Crashes before and after dedicated care. Bar graph shows count of motor vehicle crashes as a driver for each patient over 5 years for subgroup who received dedicated special care following fibromyalgia diagnosis (n = 29,007). X axis shows time divided into segments of 28 days with time-zero defined as the day of first dedicated special care treatment (K037). Y axis shows total count of crashes involving the patient as a driver that result in an emergency department visit. Rate for general population shown by horizontal line. Time line grouped as baseline interval (3 yrs) and subsequent interval (1 yr). Analysis includes all data so 1 patient might have more than 1 crash over the 5-year timespan. Results show relative reduction in motor vehicle crash counts comparing baseline to subsequent interval following dedicated special care.




Figure 4. Crashes before and after physician warning. Bar graph shows count of motor vehicle crashes as a driver for each patient over 5 years for subgroup who received a medical warning about fitness to drive following fibromyalgia diagnosis (n = 3691). X axis shows time divided into segments of 28 days with time-zero defined as the day of first medical warning about fitness to drive by a physician (K035). Y axis shows total count of crashes involving the patient as a driver that result in an emergency department visit. Rate for general population shown by horizontal line. Time line grouped as baseline interval (3 yrs) and subsequent interval (1 yr). Analysis includes all data so 1 patient might have more than 1 crash over the 5-year timespan. Results show reduction in motor vehicle crash counts comparing baseline to subsequent interval following warning.

risk of about 4.8% among drivers with FM ( $2.4 \times 2$ ). A one-third relative risk reduction following dedicated special care, therefore, might reduce this absolute annual risk to about 3.2% ( $4.8 \times$  two-thirds), mathematically equal to a number-needed-to-treat of about 63 to prevent 1 police-reported crash within 1 year. These estimates of net benefit are based on testable assumptions that could be explored in future research.

Several other limitations also remain as sources of uncertainty. We relied on ICD-9 codes (that are imprecise and overlap other muscular rheumatism disorders), included only patients with a physician diagnosis (which is fallible), and could not ascertain cases that were never diagnosed. Our study did not include passengers or pedestrians involved in the crash. We lacked data on odometers, destinations, diaries, and other indicators of the amount of active driving<sup>40</sup>. The diagnosis of FM is often unclear, diagnostic definitions change over time, and health services data tend to add further vagaries in coding comorbid diagnoses<sup>41,42</sup>. We lacked data for at-fault analyses and cannot distinguish whether FM might compromise a patient's ability to avoid a crash caused by someone else.

One strength of our study is to demonstrate a method that has advantages beyond traditional ways for estimating traffic risks. The most widely cited study on the dangers of drunk driving<sup>43</sup>, for example, would fail current standards of evidence-based medicine<sup>44</sup>. Simulator studies tend to yield ambiguous results based on healthy volunteers engaged in artificial tasks with hypothetical risks. Randomized trials are rarely attempted in traffic because of methodological and ethical obstacles<sup>45</sup>. Here we provide a methodological design that is feasible, reasonable, and relevant to an array of diseases that might predispose patients to a motor vehicle crash. Further, we illustrate the methods with a novel finding (linking FM with fitness to drive) that was not previously discovered.

Patients with FM commonly report frustration with physicians, dissatisfaction with care, and exasperation with standard biomedical paradigms for their situation<sup>46</sup>. An overly patronizing approach to motor vehicle traffic safety, therefore, might further compromise a doctor-patient relationship and lessen patient well-being<sup>47</sup>. At present, physician warnings for medically unfit motorists can be effective for improving driving safety in older patients and are mandatory in several regions of North America<sup>48,49</sup>. Clinical judgment is needed in the remaining locations when deciding whether the crash risks associated with FM merit a tactful warning in a compassionate manner to improve driving safety and maintain patient health.

## ACKNOWLEDGMENT

We thank the following for helpful comments: David Alter, Mary Bell, Gregory Choy, Edward Etchells, Leonard Evans, David Juurlink, Sharon May, Harvey Moldofsky, Jay Moss, Mark Rapoport, Matthew Schlenker, Steven Shumak, John Staples, and Christopher Yarnell.

## REFERENCES

1. Wolfe F, Smythe HA, Yunus MB, Bennett RM, Bombardier C, Goldenberg DL, et al. The American College of Rheumatology 1990 criteria for the classification of fibromyalgia. Report of the Multicenter Criteria Committee. *Arthritis Rheum* 1990;33:160-72.
2. Fan A, Wilson KG, Acharya M, Cranney A, Buenger U, Marshall S. Self-reported issues with driving in patients with chronic pain. *PM R* 2012;4:87-95.
3. Fitzcharles MA, Ste-Marie PA, Goldenberg DL, Pereira JX, Abbey S, Choinière M, et al. Canadian Pain Society and Canadian Rheumatology Association recommendations for rational care of persons with fibromyalgia: a summary report. *J Rheumatol* 2013;40:1388-93.
4. Wolfe F, Clauw DJ, Fitzcharles MA, Goldenberg DL, Katz RS, Mease P, et al. The American College of Rheumatology preliminary diagnostic criteria for fibromyalgia and measurement of symptom severity. *Arthritis Care Res* 2010;62:600-10.
5. Abeles AM, Pillinger MH, Solitar BM, Abeles M. Narrative review: the pathophysiology of fibromyalgia. *Ann Intern Med* 2007;146:726-34.
6. Redelmeier DA, Tien HC. Medical interventions to reduce motor vehicle collisions. *CMAJ* 2014;186:118-24.
7. Carr DB, Schwartzberg JG, Manning L, Sempek J. Physician's guide to assessing and counseling older drivers, 2nd ed. Chicago: American Medical Association; 2010.
8. Canadian Medical Association. CMA driver's guide: determining medical fitness to operate motor vehicles, 8th ed. Ottawa: Canadian Medical Association; 2012.
9. Wolfe F, Häuser W, Walitt BT, Katz RS, Rasker JJ, Russell AS. Fibromyalgia and physical trauma: the concepts we invent. *J Rheumatol* 2014;41:1737-45.
10. Veldhuijzen DS, van Wijck AJ, Wille F, Verster JC, Kenemans JL, Kalkman CJ, et al. Effect of chronic nonmalignant pain on highway driving performance. *Pain* 2006;122:28-35.
11. Nilsen H, Landrø NI, Kaasa S, Jenssen GD, Fayers P, Borchgrevink PC. Driving functions in a video simulator in chronic non-malignant pain patients using and not using codeine. *Eur J Pain* 2011;15:409-15.
12. Horswill MS, Sullivan K, Lurie-Beck JK, Smith S. How realistic are older drivers' ratings of their driving ability? *Accid Anal Prev* 2013;50:130-7.
13. Statistics Canada. Canada year book 2010. [Internet. Accessed April 17, 2015.] Available from: [www.statcan.gc.ca/pub/11-402-x/2010000/pdf/population-eng.pdf](http://www.statcan.gc.ca/pub/11-402-x/2010000/pdf/population-eng.pdf)
14. Ontario Ministry of Transportation. Ontario road safety annual reports (ORSAR). [Internet. Accessed April 20, 2015.] Available from: [www.mto.gov.on.ca/english/publications/ontario-road-safety-annual-report.shtml](http://www.mto.gov.on.ca/english/publications/ontario-road-safety-annual-report.shtml)
15. Ministry of Health and Long-Term Care. Schedule of benefits and fees. [Internet. Accessed April 17, 2015.] Available from: [www.health.gov.on.ca/english/providers/program/ohip/sob/physerv/physerv\\_mn.html](http://www.health.gov.on.ca/english/providers/program/ohip/sob/physerv/physerv_mn.html)
16. Ontario Medical Association. Explaining: individual counseling (K013/K033). Education and Prevention Committee interpretive bulletin. [Internet. Accessed April 17, 2015.] Available from: [www.oma.org/Resources/Documents/0301EPC\\_bulletin.pdf](http://www.oma.org/Resources/Documents/0301EPC_bulletin.pdf)
17. Ontario Medical Association. Questions and answers relevant to recent EPC bulletins. Education and Prevention Committee interpretive bulletin. [Internet. Accessed April 17, 2015.] Available from: [www.oma.org/Resources/Documents/0807EPC\\_Bulletin.pdf](http://www.oma.org/Resources/Documents/0807EPC_Bulletin.pdf)
18. Ontario Ministry of Health and Long-Term Care. Schedule of benefits for physician services amendments – April 1, 2006. [Internet. Accessed April 17, 2015.] Available from: [www.health.gov.on.ca/en/pro/programs/ohip/bulletins/4000/bul4435.pdf](http://www.health.gov.on.ca/en/pro/programs/ohip/bulletins/4000/bul4435.pdf)
19. Ontario Medical Association. Specific neurocognitive assessment

- (K032) and smoking cessation services (E079 and K039). Education and Prevention Committee interpretive bulletin. [Internet. Accessed April 17, 2015.] Available from: [www.oma.org/Resources/Documents/0802EPC\\_Bulletin.pdf](http://www.oma.org/Resources/Documents/0802EPC_Bulletin.pdf)
20. Ontario Medical Association. Billing for time-based services. Education and Prevention Committee interpretive bulletin. [Internet. Accessed April 17, 2015.] Available from: [www.oma.org/Resources/Documents/0905EPC\\_Bulletin.pdf](http://www.oma.org/Resources/Documents/0905EPC_Bulletin.pdf)
  21. Chan BT, Schull MJ, Schultz SE. Atlas Report: Emergency department services in Ontario 1993 – 2000. Toronto: Institute for Clinical Evaluative Sciences; 2001.
  22. Centers for Disease Control and Prevention. Fibromyalgia. [Internet. Accessed April 17, 2015.] Available from: [www.cdc.gov/arthritis/basics/fibromyalgia.htm](http://www.cdc.gov/arthritis/basics/fibromyalgia.htm)
  23. Muggah E, Graves E, Bennett C, Manuel DG. Ascertainment of chronic diseases using population health data: a comparison of health administrative data and patient self-report. *BMC Public Health* 2013;13:16.
  24. Hux JE, Ivis F, Flintoft V, Bica A. Diabetes in Ontario: determination of prevalence and incidence using a validated administrative data algorithm. *Diabetes Care* 2002;25:512-6.
  25. Tu K, Mitiku T, Lee DS, Guo H, Tu JV. Validation of physician billing and hospitalization data to identify patients with ischemic heart disease using data from the Electronic Medical Record Administrative data Linked Database (EMRALD). *Can J Cardiol* 2010;26:e225-8.
  26. Tu K, Wang M, Young J, Green D, Ivers NM, Butt D, et al. Validity of administrative data for identifying patients who have had a stroke or transient ischemic attack using EMRALD as a reference standard. *Can J Cardiol* 2013;29:1388-94.
  27. Schultz SE, Rothwell DM, Chen Z, Tu K. Identifying cases of congestive heart failure from administrative data: a validation study using primary care patient records. *Chronic Dis Inj Can* 2013;33:160-6.
  28. Masters ET, Mardekian J, Emir B, Clair A, Kuhn M, Silverman SL. Electronic medical record data to identify variables associated with a fibromyalgia diagnosis: importance of health care resource utilization. *J Pain Res* 2015;8:131-8.
  29. Fitzcharles MA, Ste-Marie PA, Goldenberg DL, Pereira JX, Abbey S, Choinière M, et al; the National Fibromyalgia Guideline Advisory Panel. 2012 Canadian guidelines for the diagnosis and management of fibromyalgia syndrome: executive summary. *Pain Res Manag* 2013;18:119-26.
  30. Redelmeier DA, Yarnell CJ, Thiruchelvam D, Tibshirani RJ. Physicians' warnings for unfit drivers and the risk of trauma from road crashes. *N Engl J Med* 2012;367:1228-36.
  31. World Health Organization. International statistical classification of diseases and related health problems, 10th revision. [Internet. Accessed April 20, 2015.] Available from: [apps.who.int/classifications/icd10/browse/2015/en](http://apps.who.int/classifications/icd10/browse/2015/en)
  32. Schull MJ, Hatcher CM, Guttmann A, Leaver CA, Vermeulen M, Rowe BH, et al. Development of a consensus on evidence-based quality of care indicators for Canadian emergency departments. Toronto: Institute for Clinical Evaluative Sciences; 2010.
  33. Macpherson A, Schull M, Manuel D, Cernat G, Redelmeier D, Laupacis A. Injuries in Ontario. Toronto: Institute for Clinical Evaluative Sciences; 2005.
  34. Redelmeier DA, Katz D, Lu H, Saposnik G. Roadway crash risks in recent immigrants. *Accid Anal Prev* 2011;43:2128-33.
  35. Gregory JJ, Stephens AN, Steele NA, Groeger JA. Effects of upper-limb immobilisation on driving safety. *Injury* 2009;40:253-6.
  36. Sedgwick P. Incidence rate ratio. *BMJ* 2010;341:c4804.
  37. Redelmeier DA. The exposure-crossover design is a new method for studying sustained changes in recurrent events. *J Clin Epidemiol* 2013;66:955-63.
  38. Jaakkimainen L, Upshur R, Klein-Geltink JE, Leong A, Maaten S, Schultz SE, et al, editors. Primary care in Ontario: ICES Atlas. Toronto: Institute for Clinical Evaluative Sciences; 2006.
  39. McLean SA, Williams DA, Clauw DJ. Fibromyalgia after motor vehicle collision: evidence and implications. *Traffic Inj Prev* 2005;6:97-104.
  40. Braitman KA, Williams AF. Changes in self-regulatory driving among older drivers over time. *Traffic Inj Prev* 2011;12:568-75.
  41. Ferrari R, Russell AS. A questionnaire using the modified 2010 American College of Rheumatology criteria for fibromyalgia: specificity and sensitivity in clinical practice. *J Rheumatol* 2013;40:1590-5.
  42. Ratnasingham S, Cairney J, Rehm J, Manson H, Kurdyak PA. Opening eyes, opening minds: the Ontario burden of mental illness and addictions report. Toronto: Institute for Clinical Evaluative Sciences and Public Health Ontario; 2012.
  43. Borkenstein RF, Crowther RF, Shumate RP, Ziel WB, Zylman R. The role of the drinking driver in traffic accidents. Bloomington, Indiana: Department of Police Administration, Indiana University; 1964.
  44. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP; STROBE Initiative. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *J Clin Epidemiol* 2008;61:344-9.
  45. Redelmeier DA, Yarnell CJ. Lethal misconceptions: interpretation and bias in studies of traffic deaths. *J Clin Epidemiol* 2012; 65:467-73.
  46. Kool MB, van Middendorp H, Boeije HR, Geenen R. Understanding the lack of understanding: invalidation from the perspective of the patient with fibromyalgia. *Arthritis Rheum* 2009;61:1650-6.
  47. Betz ME, Jones J, Petroff E, Schwartz R. "I wish we could normalize driving health:" a qualitative study of clinician discussions with older drivers. *J Gen Intern Med* 2013;28:1573-80.
  48. Redelmeier DA, Vinkatesh V, Stanbrook MB. Mandatory reporting by physicians of patients potentially unfit to drive. *Open Med* 2008;2:e8-e17.
  49. Wang CC, Kosinski CJ, Schwartzberg JG, Shanklin AV. Physician's guide to assessing and counseling older drivers. Washington, DC: National Highway Traffic Safety Administration; 2003.



**APPENDIX 1.** Crashes before and after electrocardiogram (EKG). Bar graph of count of motor vehicle crashes as a driver for each patient over 5 years for subgroup who received an EKG following fibromyalgia diagnosis (n = 108,089). X axis shows time divided into segments of 28 days with time-zero defined as the day of first EKG (G313). Y axis shows total count of crashes involving the patient as a driver that result in an emergency department visit. Rate for general population shown by horizontal grey line. Timeline grouped as baseline interval (3 yrs) and subsequent interval (1 yr). Analysis includes all data, so 1 patient might have more than 1 crash over the 5-year timespan. Results show no reduction in motor vehicle crash counts comparing baseline with subsequent interval following EKG.

