

Extensive Lipoatrophy on a Buttock Revealing Systemic Lupus Erythematosus Panniculitis

YA-CHI LI, MD, Division of Rheumatology/Immunology/Allergy, Department of Internal Medicine, Tri-Service General Hospital, National Defense Medical Center, Taipei, Taiwan; FU-CHIANG YEH, MD, Division of Rheumatology/Immunology/Allergy, Department of Internal Medicine, Tri-Service General Hospital, National Defense Medical Center, Taipei, Taiwan; HSIANG-CHENG CHEN, MD, PhD, Chief, Division of Rheumatology/Immunology/Allergy, Department of Internal Medicine, Tri-Service General Hospital, National Defense Medical Center, Taipei, Taiwan. Address correspondence to Dr. Hsiang-Cheng Chen, Chief, Division of Rheumatology/Immunology/Allergy, Department of Internal Medicine, Tri-Service General Hospital, National Defense Medical Center, No. 325, Sec. 2, Cheng-Gong Rd., Neihu 114, Taipei, Taiwan. E-mail: hccheng@ndmctsgh.edu.tw. *J Rheumatol* 2016;43:180–1; doi:10.3899/jrheum.150604

Lupus erythematosus panniculitis (LEP) has been reported to be the first manifestation of systemic lupus erythematosus (SLE)¹. Face and upper limbs are most commonly involved; however, extensive LEP in the buttock region or breasts is relatively rare, but also reported^{2,3}. Differential diagnosis such as subcutaneous panniculitis-like T cell lymphoma should always be considered⁴.

A 41-year-old woman presented with an area of tender swelling on her right buttock, which subsequently shrank forming an extensive depression after 3 months (Figure 1). Arthritis, malar rash, and oral ulcers were also noted. Serological studies showed positive antinuclear antibodies (1:160, nucleolar pattern) and positive anti-dsDNA (14.3 IU/ml, normal < 10 IU/ml). She was diagnosed with SLE since she fulfilled more than 4 criteria of the 1997-revised criteria for the classification of SLE⁵. Her serum complement levels were within normal range (C3: 113 mg/dl, normal 90–180 mg/dl and C4: 29.6 mg/dl, normal 10–40 mg/dl). Magnetic resonance imaging revealed a markedly decreased volume of the subcutaneous fatty layer and thickening over the covering cutis with edema along the lateroposterior site of the right gluteus maximus. A skin biopsy disclosed lobular

panniculitis with infiltration of lymphoplasmacytic cells, focal hyaline fat necrosis, and fibrin deposition on the vascular walls (Figure 2). Intravenous methylprednisolone (1 mg/kg) was prescribed for 5 days, followed by oral prednisolone (40 mg/day), hydroxychloroquine (200 mg/day), and azathioprine (100 mg/day) for more than 3 months. The extensive induration extending from her buttock to hip region indicated a poor response to treatment.

REFERENCES

1. Díaz-Jouanen E, DeHoratius RJ, Alarcón-Segovia D, Messner RP. Systemic lupus erythematosus presenting as panniculitis (lupus profundus). *Ann Intern Med* 1975;82:376-9.
2. Arai S, Katsuoka K. Clinical entity of lupus erythematosus panniculitis/lupus erythematosus profundus. *Autoimmun Rev* 2009;8:449-52.
3. Martens PB, Moder KG, Ahmed I. Lupus panniculitis: clinical perspectives from a case series. *J Rheumatol* 1999;26:68-72.
4. Ma L, Bandarchi B, Glusac EJ. Fatal subcutaneous panniculitis-like T-cell lymphoma with interface change and dermal mucin, a dead ringer for lupus erythematosus. *J Cutan Pathol* 2005;32:360-5.
5. Hochberg MC. Updating the American College of Rheumatology revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum* 1997;40:1725.



Figure 1. An extensive lipoatrophy presented on the patient's right buttock within 3 months.




Figure 2. Skin biopsy disclosed lobular panniculitis with infiltration of lymphoplasmacytic cells.