Cardiac and Pulmonary Artery Involvement Detected by ¹⁸F-fluorodeoxyglucose Positron Emission Tomography/Computed Tomography in Large-vessel Giant Cell Arteritis

RAFFAELLA UGOLINI, MD, Rheumatology Unit, Azienda Unità Sanitaria Locale IRCCS di Reggio Emilia; FRANCESCO MURATORE, MD, Rheumatology Unit, Azienda Unità Sanitaria Locale IRCCS di Reggio Emilia, and University of Modena and Reggio Emilia; CARLO SALVARANI, MD, Rheumatology Unit, Azienda Unità Sanitaria Locale IRCCS di Reggio Emilia, Italy, and University of Modena and Reggio Emilia, Reggio Emilia, Italy. Address correspondence to Dr. F. Muratore, Department of Rheumatology, Arcispedale Santa Maria Nuova IRCCS, Viale Risorgimento 80, 42123 Reggio Emilia, Italy. E-mail: francesco.muratore@ausl.re.it. Ethics approval was not required by the authors' institutions for case reports. The patient's written informed consent was obtained. *J Rheumatol* 2018;45:1603–4; doi:10.3899/jrheum.180061

Cardiac and pulmonary artery involvement have been only rarely described in giant cell arteritis^{1,2}. Further, the role of ¹⁸F-fluorodeoxyglucose positron emission tomography/computed tomography (FDG-PET/CT) for evaluating pulmonary artery involvement in large-vessel vasculitis has not been investigated.

The patient, a 58-year-old woman with a 4-month history of low-grade fever, asthenia, and weight loss presented to the emergency department of our hospital for syncope. Electrocardiogram showed a third-degree atrioventricular block requiring the implantation of a permanent pacemaker. Blood test revealed elevation of acute-phase reactants

Figure 1. FDG-PET/CT scanning shows (arrows) increased FDG uptake at the origin of the brachiocephalic trunk and left carotid artery (a), the central pulmonary artery (b), and 2 cardiac areas between aortic valve and left atrium (c). Chest angio-CT shows (arrows) wall thickening of the central pulmonary artery (d) and aortic arch (e). FDG-PET/CT: ¹⁸F-fluorodeoxyglucose positron emission tomography/computed tomography.

(C-reactive protein 100 mg/l, erythrocyte sedimentation rate 113 mm/1st h) and anemia of chronic disease (hemoglobin 10 g/dl). An initial diagnostic workup for fever of unknown origin including serologic tests, blood and urine cultures, QuantiFERON-TB gold, chest radiograph, abdominal and pelvic ultrasound, and echocardiogram were negative. A whole-body FDG-PET/CT scanning and a chest angio-CT showed inflammatory involvement of the aortic arch, the origin of the epiaortic vessels, the pulmonary artery, and 2 cardiac areas (Figure 1). These findings were consistent with large-vessel giant cell arteritis with cardiac and pulmonary artery involvement³. Cardiac magnetic resonance imaging was not performed because of recent permanent pacemaker implantation. High-dose glucocorticoid was started (prednisone 1 mg/kg/day) along with interleukin 6 inhibitor (tocilizumab 162 mg/week), with remission of symptoms and normalization of blood test.

This case demonstrates the utility of FDG-PET/CT to detect cardiac and pulmonary artery inflammation in patients with large-vessel vasculitis.

REFERENCES

1. Marten K, Schnyder P, Schirg E, Prokop M, Rummeny EJ, Engelke C. Pattern-based differential diagnosis in pulmonary vasculitis using volumetric CT. *AJR Am J of Roentgenol* 2005;184:720-33.
2. Mielniczuk L, DeKemp RA, Dennie C, Yoshinaga K, Burwash IG, Bénard F, et al. Images in cardiovascular medicine. Fluorine-18-labeled deoxyglucose positron emission tomography in the diagnosis and management of aortitis with pulmonary artery involvement. *Circulation* 2005;111:e375-6.
3. Muratore F, Kermani TA, Crowson CS, Green AB, Salvarani C, Matteson EL, et al. Large-vessel giant cell arteritis: a cohort study. *Rheumatology* 2015;54:463-70.