

Validity of Self-reported Diagnosis of Rheumatoid Arthritis

To the Editor:

I have read the article entitled "Identifying Rheumatoid Arthritis Cases within the Quebec Health Administrative Database" by Zeinab F. Slim, Cristiano Soares de Moura, Sasha Bernatsky, and Elham Rahme (*The Journal of Rheumatology*, December 2019)¹ with interest and I would like to add some comments.

In this article, rheumatoid arthritis (RA) prevalence based on a self-reported survey (CARTaGENE) was 2.9% (564 out of 19,704 patients) with a 95% CI of 2.6–3.1 before misclassification adjustment. This estimate would constitute the highest RA prevalence reported in the world since 1980, followed by Cuba (2.7%) in 2000², Finland in 1989³, and Australia in the National Health Survey (1.9%) in 2014–2015⁴. Despite a large sample size that closely resembled the general population in Canada⁵, it is much higher than the database-estimated prevalence by Lacaille, *et al* (0.8%)⁶, Widdifield, *et al* (0.9%)⁷, or Jean, *et al* (0.8%)⁸.

This estimate is highly biased for the reason that self-reported diagnosis of RA is not sufficient for the ascertainment of RA cases without clinical verification. It is an unreliable method for research or clinical purposes in various populations, because of the low percentage (21–34%) of people who will have confirmed RA^{9,10}. Our concerns are about the validity of self-reported data in the CARTaGENE study to estimate the RA prevalence in Quebec (Canada), without clinical verification by a doctor or by using any RA classification criteria.

KHALID B. AL MUTAIRI , MCLinPharm, MHA, PhD Student, King Fahd Specialist Hospital, Burydah, Saudi Arabia, and School of Medicine, The University of Western Australia, Perth, Australia; JOHANNES C. NOSSENT , FRACP, MD, PhD, Professor of Rheumatology, School of Medicine, The University of Western Australia, and Sir Charles Gairdner and Osborne Park Health Care Group, Perth; CHARLES A. INDERJEETH, MBChB, MPH, MRACP, Professor of Rheumatology and Gerontology, School of Medicine, The University of Western Australia, and Sir Charles Gairdner and Osborne Park Health Care Group. Address correspondence C/O Prof. Charles Inderjeeth, The University of Western Australia, Faculty of Health and Medical Sciences, School of Medicine, 35 Stirling Highway, Perth, Western Australia, 6009 Australia. E-mail: Charles.Inderjeeth@health.wa.gov.au

REFERENCES

1. Slim ZF, Soares de Moura C, Bernatsky S, Rahme E. Identifying

rheumatoid arthritis cases within the Quebec Health Administrative Database. *J Rheumatol* 2019;46:1570–6.

2. Reyes Llerena GA, Guibert Toledano M, Hernandez Martinez AA, Gonzalez Otero ZA, Alcocer Varela J, Cardiel MH. Prevalence of musculoskeletal complaints and disability in Cuba. A community-based study using the COPCORD core questionnaire. *Clin Exp Rheumatol* 2000;18:739–42.
3. Aho K, Heliovaara M, Sievers K, Maatela J, Isomaki H. Clinical arthritis associated with positive radiological and serological findings in Finnish adults. *Rheumatol Int* 1989;9:7–11.
4. Australian Institute of Health and Welfare. Rheumatoid arthritis [2019]. Canberra: Australian Institute of Health and Welfare. [Internet. Accessed January 9, 2020.] Available from: www.aihw.gov.au/reports/chronic-musculoskeletal-conditions/rheumatoid-arthritis
5. Awadalla P, Boileau C, Payette Y, Idaghmour Y, Goulet JP, Knoppers B, et al. Cohort profile of the CARTaGENE study: Quebec's population-based biobank for public health and personalized genomics. *Int J Epidemiol* 2013;42:1285–99.
6. Lacaille D, Anis AH, Guh DP, Esdaile JM. Gaps in care for rheumatoid arthritis: a population study. *Arthritis Rheum* 2005;53:241–8.
7. Widdifield J, Paterson JM, Bernatsky S, Tu K, Thorne JC, Ahluwalia V, et al. The rising burden of rheumatoid arthritis surpasses rheumatology supply in Ontario. *Can J Public Health* 2013;104:e450–e5.
8. Jean S, Hudson M, Gamache P, Bessette L, Fortin PR, Boire G, et al. Temporal trends in prevalence, incidence, and mortality for rheumatoid arthritis in Quebec, Canada: a population-based study. *Clin Rheumatol* 2017;36:2667–71.
9. Kvien T, Glennäs A, Knudsrød O, Smedstad L. The validity of self-reported diagnosis of rheumatoid arthritis: results from a population survey followed by clinical examinations. *J Rheumatol* 1996;23:1866–71.
10. Ling SM, Fried LP, Garrett E, Hirsch R, Guralnik JM, Hochberg MC. The accuracy of self-report of physician diagnosed rheumatoid arthritis in moderately to severely disabled older women. Women's Health and Aging Collaborative Research Group. *J Rheumatol* 2000;27:1390–4.

First Release February 15 2020; *J Rheumatol* 2020;47:5; doi:10.3899/jrheum.191401