

Predictive Utility of Cardiovascular Risk Prediction Algorithms in Inflammatory Rheumatic Diseases: A Systematic Review

Keith Colaco, Vanessa Ocampo, Ana Patricia Ayala^{ID}, Paula Harvey, Dafna D. Gladman^{ID}, Vincent Piguet, and Lihi Eder

ABSTRACT. Objective. We performed a systematic review of the literature to describe current knowledge of cardiovascular (CV) risk prediction algorithms in rheumatic diseases.

Methods. A systematic search of MEDLINE, EMBASE, and Cochrane Central databases was performed. The search was restricted to original publications in English, had to include clinical CV events as study outcomes, assess the predictive properties of at least 1 CV risk prediction algorithm, and include patients with rheumatoid arthritis (RA), ankylosing spondylitis (AS), systemic lupus erythematosus (SLE), psoriatic arthritis (PsA), or psoriasis. By design, only cohort studies that followed participants for CV events were selected.

Results. Eleven of 146 identified manuscripts were included. Studies evaluated the predictive performance of the Framingham Risk Score, QRISK2, Systematic Coronary Risk Evaluation (SCORE), Reynolds Risk Score, American College of Cardiology/American Heart Association Pooled Cohort Equations (PCE), Expanded Cardiovascular Risk Prediction Score for Rheumatoid Arthritis (ERS-RA), and the Italian Progetto CUORE score. Approaches to improve predictive performance of general risk algorithms in patients with RA included the use of multipliers, biomarkers, disease-specific variables, or a combination of these to modify or develop an algorithm. In both SLE and PsA patients, multipliers were applied to general risk algorithms. In studies of RA and SLE patients, efforts to include nontraditional risk factors, disease-related variables, multipliers, and biomarkers largely failed to substantially improve risk estimates.

Conclusion. Our study confirmed that general risk algorithms mostly underestimate and at times overestimate CV risk in rheumatic patients. We did not find studies that evaluated models for psoriasis or AS, which further demonstrates a need for research in these populations. (First Release February 15 2020; J Rheumatol 2020;47:928–38; doi:10.3899/jrheum.190261)

Key Indexing Terms:

CARDIOVASCULAR DISEASE
ALGORITHMS

INFLAMMATION

RHEUMATIC DISEASES
RISK ASSESSMENT

From Women's College Research Institute, Women's College Hospital; Centre for Prognosis Studies in the Rheumatic Diseases, Toronto Western Hospital; Institute of Medical Science, Department of Medicine, and Gerstein Science Information Centre, University of Toronto, Toronto, Ontario, Canada.

K. Colaco was supported by an award from the Enid Walker Estate and Women's College Research Institute. L. Eder was supported by a Young Investigator Award from the Arthritis Society.

K. Colaco, MSc, Women's College Research Institute, Women's College Hospital, and Centre for Prognosis Studies in the Rheumatic Diseases, Toronto Western Hospital, and Institute of Medical Science, University of Toronto; V. Ocampo, MD, Centre for Prognosis Studies in the Rheumatic Diseases, Toronto Western Hospital, and Department of Medicine, University of Toronto; A.P. Ayala, MSc, Gerstein Science Information Centre, University of Toronto; P. Harvey, BMBS, PhD, Women's College Research Institute, Women's College Hospital, and Department of Medicine, University of Toronto; D.D. Gladman, MD, Centre for Prognosis Studies in the Rheumatic Diseases, Toronto Western Hospital, and Institute of Medical Science, and Department of Medicine, University of Toronto; V. Piguet, MD, PhD, Women's College Research Institute, Women's College Hospital, and Centre for Prognosis Studies in the Rheumatic Diseases, Toronto Western Hospital, and Department of Medicine, University of Toronto; L. Eder, MD, PhD, Women's College Research Institute, Women's College Hospital, and Department of Medicine, University of Toronto.

Address correspondence to Dr. L. Eder, Women's College Research Institute, Room 6326, Women's College Hospital, 76 Grenville St., Toronto, Ontario M5S 1B2, Canada. E-mail: lihi.eder@wchospital.ca
Accepted for publication August 2, 2019.

Chronic inflammatory rheumatic diseases (IRD) are associated with significant cardiovascular (CV) morbidity and mortality^{1,2}. Patients with rheumatoid arthritis (RA)^{3,4,5}, systemic lupus erythematosus (SLE)^{6,7,8,9,10}, ankylosing spondylitis (AS)^{11,12,13,14}, psoriasis^{15,16}, and psoriatic arthritis (PsA)^{17,18,19,20,21} have an increased CV risk compared to the general population; this increased risk is attributed to a combination of systemic inflammation and high prevalence of traditional risk factors.

CV risk prediction algorithms are an important tool for clinicians to estimate patients' risk of developing future CV events. Based on the estimated risk, patients are stratified into risk groups, thereby allowing for preventive interventions to be appropriately targeted to those patients who are most likely to benefit. Therefore, precise estimates of CV risk are

desirable and could lead to more effective healthcare delivery, ultimately resulting in decreased CV morbidity and mortality. Several CV risk prediction algorithms have been developed for use in the general population. These algorithms estimate the expected CV risk using various combinations of traditional CV risk factors. The Framingham Risk Score (FRS), one of the most widely used algorithms, was developed and validated in an American cohort to calculate the 10-year risk of CV disease and was most recently updated in 2008^{22,23}. The Systematic Coronary Risk Evaluation (SCORE) algorithm was developed and validated in 12 European cohorts to predict the 10-year risk of CV mortality²⁴. In 2013, the American College of Cardiology and American Heart Association released the Pooled Cohort Equations (PCE)²⁵. The PCE was derived from large racially and geographically diverse cohort studies to predict 10-year risk of atherosclerotic CV disease events. Unlike these scores, which are based solely on traditional risk factors, the QRISK2 algorithm includes RA as an independent risk factor²⁶. Similarly, the Reynolds Risk Score (RRS) incorporates the inflammatory marker C-reactive protein (CRP) in addition to traditional risk factors^{27,28}.

The performance of these algorithms in IRD is suboptimal because traditional CV risk factors do not fully explain the increased CV risk in rheumatic patients, and current risk algorithms do not represent other contributing factors, thereby underestimating the actual CV risk²⁹. In an attempt to address these limitations, the 2016 European League Against Rheumatism (EULAR) recommendations for CV risk management proposed to apply a 1.5 multiplier to any calculated CV risk score to accommodate the risk³⁰.

The accuracy of these risk algorithms in predicting future CV events has not been summarized in IRD. Therefore, the aims of this systematic review were (1) to describe current knowledge of CV risk prediction algorithms in patients with IRD, and (2) to identify approaches to improve CV risk stratification. The results of this review could identify current knowledge gaps and inform the development of novel risk prediction algorithms.

MATERIALS AND METHODS

Study protocol. We followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines to conduct our review and analysis. We searched OVID Medline (1946 to October 30, 2017), OVID Embase (1947 to October 30, 2017), and Cochrane Central Register databases using a search strategy developed by an experienced academic health sciences librarian (APA), with input from the study leads. The search strategy (Supplementary Data 1, available with the online version of this article) was limited to English publications in humans.

Eligibility criteria and study selection. To be included in the systematic review, original studies needed to fulfill the following inclusion criteria: study design (retrospective or prospective cohort), population (psoriasis, PsA, AS, RA, or SLE), study outcome (myocardial infarction, stroke, transient ischemic attack, angina, ischemic heart disease, heart failure, CV death), and predictive ability (evaluated predictive performance of a CV risk prediction algorithm using relevant statistics).

Titles and abstracts were initially screened by 2 reviewers (KC and VO) for potential inclusion. Selected publications were retrieved in full, and 2 reviewers (KC and VO) independently assessed them for eligibility; upon disagreement, a final decision was reached through discussion with a third reviewer (LE).

Data were independently extracted by 2 reviewers (KC and VO) according to a standardized form and summarized in tables. For each study, the following information was recorded: year of publication, disease, study location, study duration, mean age, sex, data source, sample size, incidence rate of CV events, evaluated predictors, type of CV outcomes, case definition, and performance of risk score.

Risk of bias in individual studies. For assessing methodological quality and risk of bias in cohort studies, the Newcastle-Ottawa Scale was used. This tool uses a star system to judge studies on 3 broad perspectives: the selection of study groups, the comparability of the groups, and the ascertainment of the outcome of interest. The highest quality studies are awarded up to 9 stars. We decided to rate studies as low risk of bias if they received 9 stars, moderate risk of bias if they received 7 or 8 stars, and high risk of bias if they received < 7 stars, because no explicit guidance exists.

RESULTS

An initial search identified 12,099 entries, of which 146 articles were retrieved for review (Figure 1). After reviewing the manuscripts, 137 were excluded for the following reasons: 121 used at least 1 CV risk prediction algorithm but did not evaluate its predictive performance, 10 were the wrong study type (e.g., case-control, cross-sectional), and 6 did not report clinical CV events. During the preparation of the manuscript we identified 2 additional articles^{31,32} that met the inclusion criteria; however, because they were published in 2018, they were not included in the initial search results. We added these articles to the list of publications included in this review. Thus, a total of 11 studies (9 RA, 1 SLE, 1 PsA) were included in this review. The study characteristics and major findings are summarized in Table 1 and Table 2.

RA. The performance of existing risk scores in predicting CV risk varies in different studies. Crowson, *et al*³³ found that the observed CV risk was 1.8 times higher than the predicted risk by FRS. The discrepancy was particularly high in women, seropositive patients, and those with persistently elevated inflammatory markers. The RRS, which includes CRP in addition to traditional CV risk factors, showed similar deficits³³. In contrast, a more recent publication from a large international cohort showed a tendency of existing risk scores, including FRS and QRISK2, to overestimate CV risk, while the RRS underestimated CV risk³⁴. QRISK2 also overestimated risk in a Dutch cohort, whereas application of the FRS, RRS, and SCORE led to underestimations³⁵.

Several approaches for improving CV risk prediction in patients with RA were assessed. The first approach included applying a multiplication factor to existing risk scores or recalibration of these scores by applying different weights to their components. This approach was evaluated in 3 studies^{33,34,36}. In a population-based study of the Rochester Epidemiology Project, a multiplication factor of 1.8 was applied to the FRS in an attempt to improve model performance³³. Although this adjustment improved calibra-




Figure 1. PRISMA diagram. PRISMA: Preferred Reporting Items for Systematic reviews and Meta-Analyses. From: Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. PLoS Med 2009;6:e1000097. Distributed under the terms of the Creative Commons Attribution License.

tion (the agreement between observed and predicted CV risk) for patients with moderate CV risk, it had no effect on overall discrimination (correct classification of patients into the event and the non-event groups). The application of the EULAR multiplier to the FRS and PCE in an international multicenter study resulted in overestimation of future CV risk and did not improve discrimination, as measured by C-statistics, compared to the existing risk scores³⁴. Arts, *et al*³⁶ evaluated the performance of a recalibrated version of SCORE by adjusting the weights of predictors originally included in SCORE. After the recalibrated SCORE was analyzed in their Dutch cohort, there was no improvement in discrimination. In fact, the reweighted traditional risk factors underestimated CV risk across all risk groups.

A second approach, performed in a Swiss cohort³⁷, included addition of autoantibodies and biomarkers of

inflammation to the FRS. The predictive ability of these was modest: CRP, immunoglobulin M rheumatoid factor, anticyclic citrullinated peptide, oxidized low-density lipoprotein, and N-terminal pro-brain natriuretic peptide (NT-proBNP). Only anti-apolipoprotein A-I (anti-apoA-I) substantially enhanced the discrimination of the FRS. This led to a significant increase in the area under the curve (AUC) from 0.72 for FRS alone to 0.81 for the FRS and anti-apoA-I combined, corresponding to a relative increase in integrated discrimination improvement of 175%. Combining all biomarkers did not result in improvement, compared to the combination of FRS and anti-apoA-I alone. However, it should be noted that the assessments of predictive ability of the combined models were not adjusted to the time at risk and that the duration of followup varied across study patients (interquartile range 5–15 yrs).

Table 1. Characteristics of the studies included in the systematic review.

Author, Year	Dx	Country	Mean Followup, yrs (inclusion yrs)	Mean Age, yrs	Males, %	Data Source	Sample Size (no. events)	Incidence Rate of CV Events (per 100 patient-yrs)	Evaluated Predictors	Outcomes	Outcome Source	Model Properties Statistics
Alamao 2017 ³⁸	RA	UK	6.0 (1997–2010)	58.5	29 (RA), 31.6 (non-RA)	Patient registry	12,747	• RA: 4.29 (FRS), 1.78 (QRISK2); • Non-RA: 3.1 (FRS), 1.3 (QRISK2)	• FRS: All traditional risk factors, except dyslipidemia; • QRISK2: All traditional risk factors, except dyslipidemia, obesity, atrial fibrillation, renal disease	• FRS: MI, stroke, HF, aortic aneurysm, TIA, unstable angina, IC; • QRISK2: MI, CHD, stroke, TIA	ICD code	C-statistic, NRI
Arts 2015 ³⁵	RA	Netherlands	(1985/1990–2011)	54 (without CVD event = 53; with CVD event = 61)	34	Patient registry	1157 (149)	1.14	FRS, SCORE, RRS (excluding hsCRP), QRISK2	ACS, angina, CVA, TIA, PVD, HF	Chart review	C-statistic, sensitivity, specificity, PPV, NPV
Arts 2016 ³⁶	RA	Netherlands	N/A (1985–2011)	54 (without CVD event = 53; with CVD event = 62)	33.7	Patient registry	1016 (103)	N/A	SCORE vs mSCORE (smoking status, systolic BP, TC:HDL ratio, BMI, diabetes mellitus, hypertension, DAS28)	ACS, CVA, HF, CV death	Chart review	C-statistic
Crowson 2012 ³³	RA	USA	8.4 (1988–2008)	57	31	Population administrative data	525 (84)	N/A	mFRS (1.8 multiplication factor), FRS, RRS	MI, CV death, angina, stroke, IC, HF	ICD code, chart review	C-statistic
Crowson 2017 ⁴⁰	RA	UK, Norway, Netherlands, USA, Sweden, Greece, South Africa, Spain, Canada, Mexico	5.8 (1985–2014)	55	24	Several int'l patient registries	5638 (389)	1.3%/yr	• Model A (with DAS28-ESR), Model B (with HAQ), FRS, PCE, QRISK2, SCORE; • Age, sex, hypertension, current smoker, TC:HDL ratio, DAS28-ESR, HAQ	ACS, chronic ischemic heart disease, coronary revascularization, coronary death, other CV death, cerebrovascular events, peripheral vascular events	N/A	C-statistic
Crowson 2017 ³⁴	RA	UK, Norway, Netherlands, USA, South Africa, Canada, Mexico	6.9 (1985–2013, varies based on cohort)	54	26	Several int'l patient registries	1796 (100)	0.8	QRISK2, EULAR, multiplier, and ERS-RA, vs PCE, FRS-ATP, and RRS	MI, ischemic stroke, CV death	N/A	C-statistic, NRI
Finckh 2012 ³⁷	RA	Switzerland	9	77 (MACE), 64 (no MACE)	42 (MACE), 23 (no MACE)	Patient registry	118 (19)	1.70	FRS, mFRS (CRP, RF, anti-CCP, ox-LDL, NT-proBNP, anti-apoA-I)	ACS, stroke	Chart review	C-statistic, IDI
Ljung 2018 ³²	RA	Sweden	2.4–7.6 (2006–2015); varies based on cohort	54.9–61.2	25.8–28.3	Patient registry	Cohort 1: 20,822 (2017); Cohort 2: 20,477 (136); Cohort 3: 15,575 (427)	N/A	ERS-RA	MI, stroke, CV death	ICD code	C-statistic
Navarini 2018 ³¹	PsA	Italy	N/A	48	39	Patient registry	155 (21)	1.35	FRS, QRISK2, RRS, SCORE, CUORE, EULAR multiplier	CV death, CAD (stable and unstable angina, MI), CVA, TIA, PAD, HF	Chart review	C-statistic, sensitivity, specificity, PPV, NPV

Table 1. Continued.

Author, Year	Dx	Country	Mean Followup, yrs (inclusion yrs)	Mean Age, yrs	Males, %	Data Source	Sample Size (no. events)	Incidence Rate of CV Events (per 100 patient-yrs)	Evaluated Predictors	Outcomes	Outcome Source	Model Properties Statistics
Solomon 2015 ³⁹	RA	USA	2.9 (2001–2011)	57	22	Patient registry	23,605 (161)	2.5 (MI), 3.0 (stroke), 1.0 (CV death)	FRS, PCE, ERS-RA (CDAL, mHAQ-DI, prednisone use, disease duration)	MI, stroke, CV death	Chart review	C-statistic, NRI
Urowitz 2016 ⁴¹	SLE	Canada	9.0 (1970–present)	43.7 (part 1) 42.4 (part 2)	10.1 (part 1) 10.9 (part 2)	Patient registry	1013 (95)	N/A	FRS (age, sex, disease duration, hsCRP, smoking, blood pressure, cholesterol, HDL), SLEDAL-2K	CAD = MI, angina, sudden death	N/A	Sensitivity, specificity

ACS: acute coronary syndrome; anti-CCP: anticyclic citrullinated peptide; anti-apoA-I: anti-apolipoprotein A-I; BMI: body mass index; BP: blood pressure; CAD: coronary artery disease; CHD: coronary heart disease; CDAI: Clinical Disease Activity Index; hsCRP: high-sensitivity C-reactive protein; CV: cardiovascular; CVD: CV disease; CVA: cerebrovascular accident; DAS28: 28-joint count Disease Activity Score for RA; DAS28-ESR: DAS28 using erythrocyte sedimentation rate; ERS-RA: Expanded Cardiovascular Risk Prediction Score for RA; EULAR: European League Against Rheumatism; FRS: Framingham Risk Score; FRS-ATP: FRS in Adult Treatment Panel; HAQ: Health Assessment Questionnaire; HAQ-DI: HAQ-Disability Index; HDL: high-density lipoprotein; HF: heart failure; IC: intermittent claudication; ICD: International Classification of Diseases; IDI: integrated discrimination improvement; MACE: major adverse CV event; mFRS: modified FRS; mHAQ-DI: modified HAQ-DI; MI: myocardial infarction; N/A: not applicable; NPV: negative predictive value; NRI: net reclassification improvement; NT-proBNP: N-terminal pro-brain natriuretic peptide; ox-LDL: oxidized low-density lipoprotein; PAD: peripheral artery disease; PCE: American College of Cardiology/American Heart Association Pooled Cohort Equation; PsA: psoriatic arthritis; PPV: positive predictive value; PVD: peripheral vascular disease; RA: rheumatoid arthritis; RF: rheumatoid factor; RRS: Reynolds Risk Score; SCORE: Systematic Coronary Risk Evaluation; mSCORE: modified SCORE; SLE: systemic lupus erythematosus; SLEDAL-2K: SLE Disease Activity Index 2000; TC: total cholesterol; TIA: transient ischemic attack.

In a third approach, 2 studies added disease-specific variables to general risk scores^{36,38}. Alemao, *et al*³⁸ evaluated the addition of CRP to 2 existing risk scores, FRS and QRISK2, in a population-based cohort of patients with RA from the UK. Although CRP was associated with an increased CV risk when added to the FRS (12% increase in HR), the addition of CRP as a predictor to both models resulted in subtle improvements in discrimination that were clinically insignificant. In addition, reclassification using the FRS was characterized by a nonsignificant improvement and a worsening of reclassification by QRISK2. In the second study using a Dutch cohort, the original SCORE was adapted with the addition of both traditional and disease-specific risk factors³⁶. The adapted SCORE showed a subtle improvement in discriminatory ability compared to the original SCORE, which was not significant. Further, it did not lead to a significant improvement in reclassification of patients into risk groups that better matched their actual CV risk. When the adapted SCORE was evaluated in external cohorts from the UK and Norway, the discriminatory ability of the adapted model was worse than the original SCORE.

Two studies attempted to derive new RA-specific risk algorithms using traditional CV risk factors and RA characteristics^{39,40}. The Expanded Cardiovascular Risk Prediction Score for Rheumatoid Arthritis (ERS-RA) was developed and internally validated using a large patient registry in the United States³⁹. The score was derived from a base model that included only traditional CV risk factors and an expanded model that evaluated RA- and non-RA related variables. The addition of measures of RA disease activity [Clinical Disease Activity Index (CDAI)], disability [modified Health Assessment Questionnaire (HAQ)–Disability Index], daily prednisone use, and disease duration (> 10 yrs) contributed to a model that demonstrated a significant improvement in discrimination with adequate model calibration (improvement in C-statistic from 0.73 in the base model to 0.76 in the expanded model). The ERS-RA significantly improved the net reclassification of patients using both the FRS (17% of the patients) and PCE (10% of the patients) to reclassify to the correct risk categories in the expanded model. However, in a more recent study, Crowson, *et al*³⁴ found that the ERS-RA overestimated CV risk in a large international cohort and that its discriminatory ability was inferior to that of general risk scores including QRISK2, FRS, PCE, and RRS. External validation of the ERS-RA in Swedish cohorts demonstrated good discriminatory capability, and underestimation of the 10-year CV risk in high-risk groups was observed. However, no comparisons were made to general risk scores³².

The second study attempting to derive an RA-specific risk score included several international longitudinal cohorts. Crowson, *et al*⁴⁰ assessed 2 models that included traditional risk factors along with either HAQ or 28-joint count Disease Activity Score using erythrocyte sedimentation rate. Neither of these models demonstrated improved discrimination

Table 2. Summary of results of studies included in the systematic review.

Author, Year	Objective(s)	Evaluated Algorithms	C-statistics	Other Statistics	Major Findings
Alemao 2017 ³⁸	To compare the performance of FRS and QRISK2 in RA and matched non-RA patients, and to evaluate whether their performance could be enhanced by the addition of CRP	FRS FRS + CRP QRISK2 QRISK2 + CRP	0.764 0.767 0.764 0.765	FRS + CRP: NRI = 3.2% (95% CI: -2.8, 5.7) QRISK2 + CRP: NRI = -2.0% (95% CI: -5.8, 4.5)	<ul style="list-style-type: none"> •The FRS and QRISK2 underestimate CV risk. •Discrimination of both the FRS and QRISK2 was lower in the RA population compared to the general population. •The addition of CRP to both risk scores was not associated with a significant improvement in reclassification of CV risk.
Arts 2015 ³⁵	To assess the predictive ability of 4 established CV risk models for the 10-year risk of fatal and non-fatal CV diseases in European patients with RA	FRS QRISK2 RRS SCORE	0.80 0.79 0.78 0.78		The FRS, RRS, and SCORE underestimated risk of future CV events, while QRISK2 overestimated risk.
Arts 2016 ³⁶	To adapt SCORE with determinants of CV risk in RA patients and to compare the performance of the modified SCORE to the original SCORE regarding CV risk prediction in patients with RA	SCORE Recalibrated SCORE Adapted SCORE	0.78 0.78 0.80		<ul style="list-style-type: none"> •The original and adapted SCORE underestimated risk in low- and moderate-risk groups, and overestimated risk in high-risk groups •The recalibrated SCORE underestimated risk in all risk groups •Recalibrated and adapted SCORE models do not provide sufficient improvement in risk estimates compared to the original SCORE.
Crowson 2012 ³³	To assess the accuracy of the FRS and RRS for predicting CV events in patients with RA	FRS (overall) FRS (low risk) FRS (intermediate risk)	0.786 0.562 0.505		<ul style="list-style-type: none"> •The FRS significantly underestimated CV risk (especially in older ages, patients with positive RF, and those with persistently elevated ESR). •To improve calibration, FRS was multiplied by 1.8, but that had no effect on discrimination. The RRS underestimated risk in women, despite inclusion of CRP.
Crowson 2017 ⁴⁰	To develop a CV risk calculator for patients with RA	Model A (DAS28-ESR) Model B (HAQ) FRS PCE SCORE QRISK2	0.70 0.71 0.71 0.72 0.70 0.72		<ul style="list-style-type: none"> •The developed models, SCORE and QRISK2, overestimated CV risk, while the FRS and PCE underestimated risk in the highest risk groups. •Neither developed model (with the addition of HAQ and DAS28-ESR) demonstrated improved performance compared to general calculators (FRS, PCE, SCORE, QRISK2).
Crowson 2017 ³⁴	To externally validate risk algorithms recommended for use in patients with RA including the EULAR 1.5 multiplier, the ERS-RA, and QRISK2	ERS-RA QRISK2 RRS FRS-ATP FRS-ATP + EULAR multiplier PCE PCE + EULAR multiplier	0.69 0.72 0.72 0.75 0.75 0.72 0.72	ERS-RA vs PCE: NRI = -0.8% (95% CI: -8.2, 7.1) ERS-RA vs FRS: NRI = 2.3% (95% CI: -8.3, 26.6) QRISK2 vs PCE: NRI = -2.4% (95% CI: -10.9, 6.5) QRISK2 vs FRS: NRI = 25% (95% CI: -9.4, 34.7)	<ul style="list-style-type: none"> •RRS underestimated CV risk. •QRISK2, FRS, and PCE significantly overestimated CV risk. •ERS-RA overestimated CV risk, but it was less pronounced than the other risk algorithms. •RA-specific risk calculators (EULAR multiplier, ERS-RA, QRISK2) did not predict CV disease more accurately than general population risk calculators (FRS-ATP, PCE, RRS).

Table 2. Continued.

Author, Year	Objective(s)	Evaluated Algorithms	C-statistics	Other Statistics	Major Findings
Finckh 2012 ³⁷	To determine whether including CV biomarkers offers added predictive ability over the established FRS for CV risk prediction in patients with RA	FRS + CRP FRS + RF FRS + anti-CCP FRS + ox-LDL FRS + NT-proBNP FRS + anti-apoA-I	0.73 0.73 0.76 0.73 0.76 0.81	FRS + anti-apoA-I: IDI = +175.4% (p = 0.01)	NT-proBNP was moderately predictive of subsequent MACE but did not substantially improve predictive ability of traditional risk factors. Only anti-apoA-I substantially enhanced the discrimination of the FRS (improvement in AUC +0.09).
Ljung 2018 ³²	To perform an external validation of the ERS-RA in a Swedish cohort of patients with RA	ERS-RA (Cohort 1) ERS-RA (Cohort 2 – including smoking data) ERS-RA (Cohort 2 – excluding smoking data) ERS-RA (Cohort 3)	0.78 0.77 0.75 0.76		The ERS-RA had good discriminatory capability but underestimated the 10-year CV risk in high-risk groups and in the absence of data on smoking.
Navarini 2018 ³¹	To evaluate the performance of FRS, SCORE, QRISK2, RRS, and CUORE, and adapt them to EULAR guidelines in patients with PsA	SCORE SCORE + EULAR multiplier CUORE CUORE + EULAR multiplier FRS FRS + EULAR multiplier QRISK2 QRISK2 + EULAR multiplier RRS RRS + EULAR multiplier	0.7679 0.7679 0.864 0.8648 0.7575 0.7584 0.8660 0.8664 0.7183 0.7183		<ul style="list-style-type: none"> •All evaluated algorithms underestimated CV risk. •The EULAR multiplier did not increase the discriminative ability or calibration of any of the evaluated algorithms.
Solomon 2015 ³⁹	To develop and internally validate an expanded CV risk prediction score for RA	Base algorithm Developed algorithm (ERS-RA)	0.7261 0.7609	Base model vs ERS-RA (FRS): NRI = 40% (95% CI: 37, 44) Base model vs ERS-RA (PCE): NRI = 7% (95% CI: 6, 8)	<ul style="list-style-type: none"> •Model discrimination improved significantly from the base model to the expanded model (ERS-RA). •RA disease activity, disability, daily prednisone use and disease duration contributed to a significantly improved model.
Urowitz 2016 ⁴¹	To determine whether an adjustment to the FRS would more accurately reflect the higher prevalence of coronary artery disease among patients with SLE.	FRS 1.5 FRS 2 FRS 3 FRS 4 FRS	N/A	Sensitivity: 13.0, Specificity: 98.2 Sensitivity: 19.7, Specificity: 89.4 Sensitivity: 31.5, Specificity: 80.9 Sensitivity: 45.5, Specificity: 72.0 Sensitivity: 46.1, Specificity: 68.8	Applying a multiplication factor of 2 to the FRS more accurately identified patients at moderate/high risk of coronary artery disease and more accurately predicts subsequent coronary artery disease.

Anti-CCP: anticyclic citrullinated peptide; anti-apoA-I: anti-apolipoprotein A-I; AUC: area under the curve; CRP: C-reactive protein; CV: cardiovascular; ESR: erythrocyte sedimentation rate; DAS28-ESR: 28-joint count Disease Activity Score for RA using ESR; ERS-RA: Expanded Cardiovascular Risk Prediction Score for RA; EULAR: European League Against Rheumatism; FRS: Framingham Risk Score; FRS-ATP: FRS in Adult Treatment Panel; HAQ: Health Assessment Questionnaire; IDI: integrated discrimination improvement; MACE: major adverse CV event; N/A: not applicable; NRI: net reclassification improvement; NT-proBNP: N-terminal pro-brain natriuretic peptide; ox-LDL: oxidized low-density lipoprotein; PCE: American College of Cardiology/American Heart Association Pooled Cohort Equation; PsA: psoriatic arthritis; RA: rheumatoid arthritis; RF: rheumatoid factor; RRS: Reynolds Risk Score; SCORE: Systematic Coronary Risk Evaluation; SLE: systemic lupus erythematosus.

compared to general risk scores including FRS, PCE, SCORE, or QRISK2 (C-statistic ranged from 0.70 to 0.72). Although the RA-specific models showed better calibration than the general risk scores, this may have occurred because

calibration is expected to be better in the cohort used to develop the new risk score than the general scores developed in other models. The developed models also significantly overestimated CV events.

SLE. In the study that followed patients seen at the University of Toronto Lupus Clinic since 1970, the FRS was compared to a modified FRS with 4 multiplication factors (range 1.5–4)⁴¹. A multiplier of 2 predicted CV risk more accurately than the original FRS (20% vs 2.9% risk), and most appropriately categorized patients with moderate/high risk with sufficient sensitivity and specificity. The sensitivity and specificity for FRS (moderate/high risk vs low risk) to predict CV events were 6.8 and 98.1, respectively, whereas the sensitivity for the FRS with a multiplication factor of 2 increased to 34.5 with a modest reduction in specificity to 84.4, respectively. The study did not compare global measures of model predictive ability between the models (e.g., AUC).

PsA. In a small Italian cohort of patients with PsA, 5 general risk scores (FRS, SCORE, QRISK2, RRS, and CUORE) were adapted to EULAR recommendations by adding a 1.5 multiplier or including weight adaptation for RA in QRISK2³¹. The 5 algorithms underestimated CV risk, and the adaptation suggested by EULAR did not increase the discriminative ability or calibration of any of the evaluated algorithms. Overall, the original risk scores demonstrated relatively good discrimination between patients with or without CV events, with a range of AUC between 0.718 (for RRS) and 0.866 (for QRISK2).

Quality assessment. Two studies had low risk of bias, 7 studies had moderate risk of bias, and 2 studies had high risk of bias (Table 3). In one study, it was unclear whether participants with a history of CV disease were excluded from analysis³⁸. There was limited information on how CV events were ascertained in 3 studies^{34,40,41}. Owing to the limited number of studies included in this review, those with lower scores were not excluded from the quality assessment.

DISCUSSION

There are several challenges associated with deriving and validating disease-specific algorithms or modifying existing

general risk scores to improve risk prediction and stratification. This study identified potential predictors of future CV events that warrant further investigation.

Most studies evaluated performance using clinical variables rather than novel laboratory biomarkers, which may be more difficult to implement in a clinical setting. Findings by Finckh, *et al*³⁷ that anti-apoA-I significantly improved the predictive accuracy of the FRS demonstrates its potential as a clinically useful CV biomarker, because it is easily measurable and may assist in identification of high-risk RA patients⁴². Anti-apoA-I antibodies and NT-proBNP have been found to be associated with increased atherosclerotic plaque vulnerability and cardiac ischemia, respectively^{43,44,45}. Other risk markers have shown promise in improving risk discrimination in rheumatic patients. Several noninvasive imaging techniques, including carotid ultrasound (US)^{29,46,47} and coronary artery calcium (CAC) quantification by computed tomography, have identified markers for determining subclinical atherosclerosis. In RA, carotid atherosclerosis as assessed by US was found to predict CV events in patients with greater carotid intima-media thickness and in those with bilateral plaques⁴⁸. In patients with RA stratified according to a modified SCORE, carotid US was sensitive to detect patients at moderate risk (1–5%)⁴⁹. When compared to CAC, carotid US was found to be more sensitive in the stratification of CV risk⁵⁰, and similar results were reported in an axial spondyloarthritis group⁵¹. These results highlight the potential use of carotid US for improving CV risk stratification in rheumatic patients and encourage further research of this tool in combination with traditional risk scores.

Multipliers have been widely applied to general risk calculators so that they more accurately reflect the effect of each variable in the algorithm while retaining their relative value. Despite EULAR's recommendation³⁰, other studies show that applying the multiplication factor does not significantly improve risk prediction^{31,36}. In addition, application of the

Table 3. Newcastle-Ottawa Quality Assessment Scale for cohort studies.

Author, Year	Selection (4 stars)	Comparability (2 stars)	Outcome (3 stars)	Score (9 stars)	Risk of Bias
Alemao 2017 ³⁸	★★★	★★	★★★	8	Moderate
Arts 2015 ³⁵	★★★	★★	★★	7	Moderate
Arts 2016 ³⁶	★★★	★★	★★	7	Moderate
Crowson 2012 ³³	★★★★	★★	★★★	9	Low
Crowson 2017 ⁴⁰	★★★	★★	★	6	High
Crowson 2017 ³⁴	★★★	★★	★	6	High
Finckh 2012 ³⁷	★★★	★★	★★	7	Moderate
Ljung 2018 ³²	★★★★	★★	★★★	9	Low
Navarini 2018 ³¹	★★★	★★	★★	7	Moderate
Solomon 2015 ³⁹	★★★	★★	★★★	8	Moderate
Urowitz 2016 ⁴¹	★★★★	★★	★★	8	Moderate

Risk of bias was assessed using the Newcastle-Ottawa scale. Studies were judged on 3 broad perspectives: the selection of study groups, the comparability of the groups, and the ascertainment of the outcome of interest for cohort studies. A study is awarded stars for items within each category for a maximum of 9 stars. We rated studies as low risk of bias if they received 9 stars, moderate risk of bias if they received 7 or 8 stars, and high risk of bias if they received < 7 stars.

multiplier reduced calibration without improving discrimination or reclassification to the correct CV risk category in patients with RA³⁴. Similar results were found in patients with PsA, where the EULAR multiplier failed to demonstrate improvement in both discrimination and calibration for any of the 5 evaluated general risk scores³¹. On the other hand, Urowitz, *et al*⁴¹ applied a factor of 2 to the FRS, which improved the accuracy of classified SLE patients at moderate/high risk. It should be noted that even after this adaptation, the sensitivity to identify high-risk individuals was only about 30% and the study did not assess measures of discrimination and calibration of the suggested adaptation. Whether these results have a similar effect in other SLE populations is questionable, given that the FRS variables were not recalibrated to SLE and retained the same weights derived from the general population. Although multipliers can be used as a means to improve risk prediction, they would improve calibration, but not discrimination, resulting in a missed opportunity to intervene early³³.

Four RA studies included either inflammatory biomarkers, disease-specific variables, or a combination of both to modify or develop an algorithm^{36,38,39,40}. The only algorithm to significantly improve predictive performance was the ERS-RA, which could be readily applied to a clinical setting. The ERS-RA does not incorporate novel serum biomarkers, but its use of the CDAI, a composite measure of disease activity, may be representative of systemic inflammation underlying the excess CV risk seen in RA. However, the ERS-RA may lead to inaccuracies in estimation of risk. The score was developed using registry data that did not contain lipid levels or blood pressure measurements, and included a population with a mean followup rate of less than 3 years. When the ERS-RA was evaluated in an international multicenter cohort study, it overestimated risk and produced lower risk estimates than current risk algorithms³⁴. However, it performed well in a Swedish cohort and showed excellent calibration for patients with 5–10% 10-year CV risk, but analyses included partial data on smoking status³². The CV risk profile of the American cohort in which the ERS-RA was derived and internally validated may be generalizable to other non-American cohorts.

This study identified 1 disease-specific algorithm (ERS-RA), and its performance varied after being externally validated in Swedish and international cohorts. Further validation and tailoring of the ERS-RA to specific populations is needed before recommendations can be made. Our review also affirmed that general risk algorithms do not perform well in rheumatic patients. These models were largely derived in cohorts established in the late 20th century when participants were less socioeconomically and ethnically diverse, and CV event rates were more than double the current rates⁵². Only the QRISK equations are regularly updated in modern cohorts and include several predictors, such as deprivation measures, but their applicability outside

the UK is limited⁵². Though it is possible to update existing algorithms, this approach has limitations. Yadlowsky, *et al* evaluated 2 approaches for improving the PCE: using the PCE with updated cohort data, and using both updated data and new derivation methods⁵³. The first approach modestly improved discrimination, whereas the second approach improved both calibration and discrimination. Most general risk algorithms are also likely to be out of date because of major changes in preventive treatments over recent decades^{52,54}. Among the risk factor-modifying drugs, statins have been recently studied in rheumatic patients owing to their lipid-lowering effects and antiinflammatory properties. A randomized trial of patients with RA found that the addition of statins to disease-modifying antirheumatic drug treatment improved disease variables such as swollen joint count and inflammatory markers⁵⁵. However, a separate trial in RA showed that statins had no effect on disease activity⁵⁶. A previous randomized trial was initiated to examine the effect of atorvastatin in preventing CV events in patients with RA, but the trial was terminated early as a result of a low event rate⁵⁷. Other previously published studies also highlight the role of statins in carotid plaque regression⁵⁸ and mortality reduction^{59,60} in RA, AS, and PsA. The lack of account for treatment effect can cause difficulties in the use of CV risk algorithms and underestimation of CV risk. Ideally, risk algorithms should be derived from populations free of treatment. Regarding outcome definitions, most algorithms predicted the risk of fatal or nonfatal coronary heart disease or the combined outcome of CV disease. Because different definitions of CV outcomes lead to different estimated predictor effects, international consensus on a more uniform definition is necessary to aid comparison of developed risk algorithms. Given the challenges associated with using outdated cohort data, increased use of preventive therapy for CV events, and variation in outcome definitions, it is not possible to recommend a general risk algorithm for rheumatic patients.

It appears that subclinical vascular disease is not accurately reflected in risk algorithms, leading to underestimated CV risk and preventable CV events^{61,62,63}. We suggest that measures of subclinical vascular disease be used to improve risk estimates beyond models that use traditional CV risk factors alone. Carotid US or CAC may optimize CV risk estimation and aid in more accurate CV risk stratification. Additional predictors, including measures of ethnicity and socioeconomic status, are also needed to avoid undertreatment of high-risk groups. Although recalibration is likely to reduce overtreatment, general risk algorithms will continue to underperform in rheumatic patients. Unless risk of CV disease is estimated using algorithms derived or recalibrated in present-day populations that represent the patients they are applied to, under- or overestimation of risk is likely to persist.

Our study confirmed that general risk algorithms mostly underestimate and at times overestimate CV risk in rheumatic patients and the excess CV risk in these patients cannot be

explained by traditional risk factors alone. Efforts to include nontraditional risk factors, disease-related variables, multipliers, and biomarkers largely failed at substantially improving risk estimates. Rather than recalibrating general risk algorithms, future research should place more emphasis on developing new models and identifying new disease-specific predictors. Further validation and recalibration of the ERS-RA to target populations is needed before recommendations can be made for use in patients with RA.

ONLINE SUPPLEMENT

Supplementary material accompanies the online version of this article.

REFERENCES

1. Symmons DP, Gabriel SE. Epidemiology of CVD in rheumatic disease, with a focus on RA and SLE. *Nat Rev Rheumatol* 2011;7:399-408.
2. Mok CC, Ko GT, Ho LY, Yu KL, Chan PT, To CH. Prevalence of atherosclerotic risk factors and the metabolic syndrome in patients with chronic inflammatory arthritis. *Arthritis Care Res* 2011; 63:195-202.
3. Avina-Zubieta JA, Thomas J, Sadatsafavi M, Lehman AJ, Lacaille D. Risk of incident cardiovascular events in patients with rheumatoid arthritis: a meta-analysis of observational studies. *Ann Rheum Dis* 2012;71:1524-9.
4. Gabriel SE. Cardiovascular morbidity and mortality in rheumatoid arthritis. *Am J Med* 2008;10 Suppl 1:S9-14.
5. Meune C, Touze E, Trinquart L, Allanore Y. Trends in cardiovascular mortality in patients with rheumatoid arthritis over 50 years: a systematic review and meta-analysis of cohort studies. *Rheumatology* 2009;48:1309-13.
6. Goldberg RJ, Urowitz MB, Ibanez D, Nikpour M, Gladman DD. Risk factors for development of coronary artery disease in women with systemic lupus erythematosus. *J Rheumatol* 2009;36:2454-61.
7. Ahmad Y, Shelmerdine J, Bodill H, Lunt M, Patrick MG, Teh LS, et al. Subclinical atherosclerosis in systemic lupus erythematosus (SLE): the relative contribution of classic risk factors and the lupus phenotype. *Rheumatology* 2007;46:983-8.
8. Manzi S, Selzer F, Sutton-Tyrrell K, Fitzgerald SG, Rairie JE, Tracy RP, et al. Prevalence and risk factors of carotid plaque in women with systemic lupus erythematosus. *Arthritis Rheum* 1999;42:51-60.
9. Roman MJ, Shanker BA, Davis A, Lockshin MD, Sammaritano L, Simantov R, et al. Prevalence and correlates of accelerated atherosclerosis in systemic lupus erythematosus. *N Engl J Med* 2003;349:2399-406.
10. de Leeuw K, Freire B, Smit AJ, Bootsma H, Kallenberg CG, Bijl M. Traditional and non-traditional risk factors contribute to the development of accelerated atherosclerosis in patients with systemic lupus erythematosus. *Lupus* 2006;15:675-82.
11. Peters MJ, van der Horst-Bruinsma IE, Dijkmans BA, Nurmohamed MT. Cardiovascular risk profile of patients with spondylarthropathies, particularly ankylosing spondylitis and psoriatic arthritis. *Semin Arthritis Rheum* 2004;34:585-92.
12. Rueda-Gotor J, Corrales A, Blanco R, Fuentevilla P, Portilla V, Exposito R, et al. Atherosclerotic disease in axial spondyloarthritis: increased frequency of carotid plaques. *Clin Exp Rheumatol* 2015;33:315-20.
13. Mathieu S, Pereira B, Soubrier M. Cardiovascular events in ankylosing spondylitis: an updated meta-analysis. *Semin Arthritis Rheum* 2015;44:551-5.
14. Haroon NN, Paterson JM, Li P, Inman RD, Haroon N. Patients with ankylosing spondylitis have increased cardiovascular and cerebrovascular mortality: a population-based study. *Ann Intern Med* 2015;163:409-16.
15. Gelfand JM, Neimann AL, Shin DB, Wang X, Margolis DJ, Troxel AB. Risk of myocardial infarction in patients with psoriasis. *JAMA* 2006;296:1735-41.
16. Mehta NN, Yu Y, Pinnelas R, Krishnamoorthy P, Shin DB, Troxel AB, et al. Attributable risk estimate of severe psoriasis on major cardiovascular events. *Am J Med* 2011;124:775.e1-6.
17. Eder L, Thavaneswaran A, Chandran V, Cook R, Gladman DD. Increased burden of inflammation over time is associated with the extent of atherosclerotic plaques in patients with psoriatic arthritis. *Ann Rheum Dis* 2015;74:1830-5.
18. Sobchak C, Eder L. Cardiometabolic disorders in psoriatic disease. *Curr Rheumatol Rep* 2017;19:63.
19. Tam LS, Tomlinson B, Chu TT, Li M, Leung YY, Kwok LW, et al. Cardiovascular risk profile of patients with psoriatic arthritis compared to controls—the role of inflammation. *Rheumatology* 2008;47:718-23.
20. Horreau C, Pouplard C, Brenaut E, Barnette T, Misery L, Cribier B, et al. Cardiovascular morbidity and mortality in psoriasis and psoriatic arthritis: a systematic literature review. *J Eur Acad Dermatol Venerol* 2013;27 Suppl 3:12-29.
21. Gladman DD. Mortality in psoriatic arthritis. *Clin Exp Rheumatol* 2008;5 Suppl 51:S62-5.
22. D'Agostino RB Sr, Vasani RS, Pencina MJ, Wolf PA, Cobain M, Massaro JM, et al. General cardiovascular risk profile for use in primary care: the Framingham Heart Study. *Circulation* 2008;117:743-53.
23. Wilson PW, D'Agostino RB, Levy D, Belanger AM, Silbershatz H, Kannel WB. Prediction of coronary heart disease using risk factor categories. *Circulation* 1998;97:1837-47.
24. Conroy RM, Pyorala K, Fitzgerald AP, Sans S, Menotti A, De Backer G, et al; SCORE project group. Estimation of ten-year risk of fatal cardiovascular disease in Europe: the SCORE project. *Eur Heart J* 2003;24:987-1003.
25. Goff DC Jr, Lloyd-Jones DM, Bennett G, Coady S, D'Agostino RB Sr, Gibbons R, et al. 2013 ACC/AHA guideline on the assessment of cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2014;25 Pt B:2935-59.
26. Hippisley-Cox J, Coupland C, Vinogradova Y, Robson J, Minhas R, Sheikh A, et al. Predicting cardiovascular risk in England and Wales: prospective derivation and validation of QRISK2. *BMJ* 2008;336:1475-82.
27. Ridker PM, Buring JE, Rifai N, Cook NR. Development and validation of improved algorithms for the assessment of global cardiovascular risk in women: the Reynolds Risk Score. *JAMA* 2007;297:611-9.
28. Ridker PM, Paynter NP, Rifai N, Gaziano JM, Cook NR. C-reactive protein and parental history improve global cardiovascular risk prediction: the Reynolds Risk Score for men. *Circulation* 2008;118:2243-51, 4p following 51.
29. Eder L, Chandran V, Gladman DD. The Framingham Risk Score underestimates the extent of subclinical atherosclerosis in patients with psoriatic disease. *Ann Rheum Dis* 2014;73:1990-6.
30. Agca R, Heslinga SC, Rollefstad S, Heslinga M, McInnes IB, Peters MJ, et al. EULAR recommendations for cardiovascular disease risk management in patients with rheumatoid arthritis and other forms of inflammatory joint disorders: 2015/2016 update. *Ann Rheum Dis* 2017;76:17-28.
31. Navarini L, Margiotta DPE, Caso F, Currado D, Tasso M, Angeletti S, et al. Performances of five risk algorithms in predicting cardiovascular events in patients with psoriatic arthritis: an Italian bicentric study. *PLoS ONE* 2018;13:e0205506.
32. Ljung L, Ueda P, Liao KP, Greenberg JD, Etzel CJ, Solomon DH, et

- al. Performance of the Expanded Cardiovascular Risk Prediction Score for Rheumatoid Arthritis in a geographically distant national register-based cohort: an external validation. *RMD Open* 2018;4:e000771.
33. Crowson CS, Matteson EL, Roger VL, Thernau TM, Gabriel SE. Usefulness of risk scores to estimate the risk of cardiovascular disease in patients with rheumatoid arthritis. *Am J Cardiol* 2012;110:420-4.
34. Crowson CS, Gabriel SE, Semb AG, van Riel P, Karpouzias G, Dessein PH, et al; Trans-Atlantic Cardiovascular Consortium for Rheumatoid Arthritis. Rheumatoid arthritis-specific cardiovascular risk scores are not superior to general risk scores: a validation analysis of patients from seven countries. *Rheumatology* 2017;56:1102-10.
35. Arts EE, Popa C, Den Broeder AA, Semb AG, Toms T, Kitas GD, et al. Performance of four current risk algorithms in predicting cardiovascular events in patients with early rheumatoid arthritis. *Ann Rheum Dis* 2015;74:668-74.
36. Arts EE, Popa CD, Den Broeder AA, Donders R, Sandoo A, Toms T, et al. Prediction of cardiovascular risk in rheumatoid arthritis: performance of original and adapted SCORE algorithms. *Ann Rheum Dis* 2016;75:674-80.
37. Finckh A, Courvoisier DS, Pagano S, Bas S, Chevallier-Ruggeri P, Hochstrasser D, et al. Evaluation of cardiovascular risk in patients with rheumatoid arthritis: do cardiovascular biomarkers offer added predictive ability over established clinical risk scores? *Arthritis Care Res* 2012;64:817-25.
38. Alemao E, Cawston H, Bourhis F, Al M, Rutten-van Molken M, Liao KP, et al. Comparison of cardiovascular risk algorithms in patients with vs without rheumatoid arthritis and the role of C-reactive protein in predicting cardiovascular outcomes in rheumatoid arthritis. *Rheumatology* 2017;56:777-86.
39. Solomon DH, Greenberg J, Curtis JR, Liu M, Farkouh ME, Tsao P, et al. Derivation and internal validation of an expanded cardiovascular risk prediction score for rheumatoid arthritis: a Consortium of Rheumatology Researchers of North America Registry Study. *Arthritis Rheumatol* 2015;67:1995-2003.
40. Crowson CS, Rollefstad S, Kitas GD, van Riel PL, Gabriel SE, Semb AG; A Trans-Atlantic Cardiovascular Risk Consortium for Rheumatoid Arthritis (ATACC-RA). Challenges of developing a cardiovascular risk calculator for patients with rheumatoid arthritis. *PLoS ONE* 2017;12:e0174656.
41. Urowitz MB, Ibañez D, Su J, Gladman DD. Modified Framingham Risk Factor Score for systemic lupus erythematosus. *J Rheumatol* 2016;43:875-9.
42. Morrow DA, de Lemos JA. Benchmarks for the assessment of novel cardiovascular biomarkers. *Circulation* 2007;115:949-52.
43. Siriwardena M, Kleffmann T, Ruygrok P, Cameron VA, Yandle TG, Nicholls MG, et al. B-type natriuretic peptide signal peptide circulates in human blood: evaluation as a potential biomarker of cardiac ischemia. *Circulation* 2010;122:255-64.
44. Vuilleumier N, Rossier MF, Pagano S, Python M, Charbonney E, Nkoulou R, et al. Anti-apolipoprotein A-1 IgG as an independent cardiovascular prognostic marker affecting basal heart rate in myocardial infarction. *Eur Heart J* 2010;31:815-23.
45. Montecucco F, Vuilleumier N, Pagano S, Lenglet S, Bertolotto M, Braunersreuther V, et al. Anti-apolipoprotein A-1 auto-antibodies are active mediators of atherosclerotic plaque vulnerability. *Eur Heart J* 2011;32:412-21.
46. Eder L, Jayakar J, Shanmugarajah S, Thavaneswaran A, Pereira D, Chandran V, et al. The burden of carotid artery plaques is higher in patients with psoriatic arthritis compared with those with psoriasis alone. *Ann Rheum Dis* 2013;72:715-20.
47. Eder L, Gladman DD, Ibañez D, Urowitz MB. The correlation between carotid artery atherosclerosis and clinical ischemic heart disease in lupus patients. *Lupus* 2014;23:1142-8.
48. Evans MR, Escalante A, Battafarano DF, Freeman GL, O'Leary DH, del Rincon I. Carotid atherosclerosis predicts incident acute coronary syndromes in rheumatoid arthritis. *Arthritis Rheum* 2011;63:1211-20.
49. Corrales A, Gonzalez-Juanatey C, Peiro ME, Blanco R, Llorca J, Gonzalez-Gay MA. Carotid ultrasound is useful for the cardiovascular risk stratification of patients with rheumatoid arthritis: results of a population-based study. *Ann Rheum Dis* 2014;73:722-7.
50. Corrales A, Parra JA, Gonzalez-Juanatey C, Rueda-Gotor J, Blanco R, Llorca J, et al. Cardiovascular risk stratification in rheumatic diseases: carotid ultrasound is more sensitive than Coronary Artery Calcification Score to detect subclinical atherosclerosis in patients with rheumatoid arthritis. *Ann Rheum Dis* 2013;72:1764-70.
51. Rueda-Gotor J, Llorca J, Corrales A, Parra JA, Portilla V, Genre F, et al. Cardiovascular risk stratification in axial spondyloarthritis: carotid ultrasound is more sensitive than coronary artery calcification score to detect high-cardiovascular risk axial spondyloarthritis patients. *Clin Exp Rheumatol* 2018;36:73-80.
52. Pylypchuk R, Wells S, Kerr A, Poppe K, Riddell T, Harwood M, et al. Cardiovascular disease risk prediction equations in 400 000 primary care patients in New Zealand: a derivation and validation study. *Lancet* 2018;391:1897-907.
53. Yadlowsky S, Hayward RA, Sussman JB, McClelland RL, Min YI, Basu S. Clinical implications of revised pooled cohort equations for estimating atherosclerotic cardiovascular disease risk. *Ann Intern Med* 2018;169:20-9.
54. Liew SM, Doust J, Glasziou P. Cardiovascular risk scores do not account for the effect of treatment: a review. *Heart* 2011;97:689-97.
55. McCarey DW, McInnes IB, Madhok R, Hampson R, Scherbakov O, Ford I, et al. Trial of atorvastatin in rheumatoid arthritis (TARA): double-blind, randomised placebo-controlled trial. *Lancet* 2004;363:2015-21.
56. Kumar P, Kennedy G, Khan F, Pullar T, Belch JJ. Rosuvastatin might have an effect on C-reactive protein but not on rheumatoid disease activity: Tayside randomized controlled study. *Scott Med J* 2012;57:80-3.
57. Kitas GD, Nightingale P, Armitage J, Sattar N, Belch JJ, Symmons DP; TRACE RA Consortium. A multicenter, randomized, placebo-controlled trial of atorvastatin for the primary prevention of cardiovascular events in patients with rheumatoid arthritis. *Arthritis Rheumatol* 2019; 71:1437-49.
58. Rollefstad S, Ikdahl E, Hisdal J, Olsen IC, Holme I, Hammer HB, et al. Rosuvastatin-induced carotid plaque regression in patients with inflammatory joint diseases: the rosuvastatin in rheumatoid arthritis, ankylosing spondylitis and other inflammatory joint diseases study. *Arthritis Rheumatol* 2015;67:1718-28.
59. Oza A, Lu N, Schoenfeld SR, Fisher MC, Dubreuil M, Rai SK, et al. Survival benefit of statin use in ankylosing spondylitis: a general population-based cohort study. *Ann Rheum Dis* 2017;76:1737-42.
60. Schoenfeld SR, Lu L, Rai SK, Seeger JD, Zhang Y, Choi HK. Statin use and mortality in rheumatoid arthritis: a general population-based cohort study. *Ann Rheum Dis* 2016;75:1315-20.
61. Nikpour M, Urowitz MB, Ibanez D, Harvey PJ, Gladman DD. Importance of cumulative exposure to elevated cholesterol and blood pressure in development of atherosclerotic coronary artery disease in systemic lupus erythematosus: a prospective proof-of-concept cohort study. *Arthritis Res Ther* 2011;13:R156.
62. Nikpour M, Harvey PJ, Ibanez D, Gladman DD, Urowitz MB. High-sensitivity C-reactive protein as a marker of cardiovascular risk in systemic lupus erythematosus. *Arthritis Rheum* 2012;64:3052-3.
63. Nikpour M, Gladman DD, Ibañez D, Urowitz MB. Variability and correlates of high sensitivity C-reactive protein in systemic lupus erythematosus. *Lupus* 2009;18:966-73.