

Copy Number Variation of Multiple Genes in SAPHO Syndrome

Changlong Guo^{ID}, Xin Tian, Feifei Han, Lihong Liu, Jianen Gao^{ID}, and Xu Ma

ABSTRACT. *Objective.* SAPHO (synovitis, acne, pustulosis, hyperostosis, osteitis) syndrome is a type of rare chronic aseptic inflammation of unknown etiology. To date, no research to our knowledge has reported copy number variation (CNV) of genes that could affect predisposition to SAPHO syndrome. We investigated the association between CNV profile and SAPHO syndrome.

Methods. We used array comparative genomic hybridization (CGH) to screen for CNV in a nuclear family including 2 patients and a healthy control. We then validated the copy numbers of candidate genes found in the array CGH assay and other candidate genes by TaqMan real-time PCR in 360 case and control samples.

Results. Ten regions from 8 chromosomes were found to have abnormal gene copies in the nuclear family, so the CNV of candidate genes (*ADAM5*, *CSF2RA*, *IL3RA*, and 9 other genes) were tested by TaqMan PCR. Significant copy number loss of *CSF2RA* ($p = 0.000$) and *NOD2* ($p = 0.005$), and significant copy number gain of *MEGF6* ($p = 0.002$) and *ADAM5* ($p = 0.000$) were seen in patients with SAPHO compared with controls at the $\alpha = 0.05$ level. There were no differences in the other 8 candidate genes between patient and control samples ($p > 0.05$).

Conclusion. Our study established the first association between CNV in *CSF2RA*, *NOD2*, *MEGF6*, and *ADAM5* and SAPHO syndrome. These findings may offer insight into the pathogenesis of SAPHO and provide the basis for improved diagnosis and treatment. (First Release April 15 2020; J Rheumatol 2020;47:1323–9; doi:10.3899/jrheum.181393)

Key Indexing Terms:

SAPHO SYNDROME

ARRAY COMPARATIVE GENOMIC HYBRIDIZATION

COPY NUMBER VARIATION

TAQMAN PCR

SAPHO (synovitis, acne, pustulosis, hyperostosis, osteitis) syndrome, a rare disease with an estimated prevalence of less than 1 in 10,000¹, was first identified in 1987². SAPHO syndrome can occur at any age³, and its exact etiology is unknown. However, a large body of literature supports a pathogenesis involving a combination of genetic, infectious, and immunological components⁴. Experiments in mice indicate that SAPHO is associated with mutations in *PSTPIP1*,

*PSTPIP2*⁵, and *NOD2* genes^{6,7}, but the genetic basis of human SAPHO syndrome remains poorly understood.

Copy number variation (CNV) is widespread in human populations, with 5%–10% of the human reference genome showing CNV in healthy individuals^{8,9}. Several studies have concluded that variation in DNA copy number may influence the expression of genes or even alter their structure, thus contributing to substantial phenotypic variation. Moreover, CNV of genes causes a variety of human genetic diseases¹⁰.

Little is known about CNV as a potential risk factor for SAPHO syndrome; we screened the CNV profile in a nuclear family using array comparative genomic hybridization (CGH). We then further confirmed the candidate CNV genes in a larger sample size (156 SAPHO cases and 204 healthy controls) using TaqMan real-time PCR. We hope to propose the potential application of CNV evaluation in individuals at high risk of developing SAPHO syndrome, and the use of CNV genes as candidates for gene therapy.

MATERIALS AND METHODS

In the first stage, a core family with SAPHO syndrome was identified [mother: age 51 yrs, with SAPHO syndrome, with palmoplantar pustulosis and osteoarticular symptoms (anterior chest wall, spine, peripheral skeleton), duration of disease 8 yrs; daughter: age 22 yrs, with SAPHO syndrome, with osteoarticular symptoms (anterior chest wall, spine), duration of disease 1.5 yrs; father: age 53 yrs, healthy control]. The second group comprised 156 patients with SAPHO syndrome (92 women, 64

From the National Research Institute for Family Planning, Beijing; The No. 1 Hospital of Shijiazhuang, Shijiazhuang; Shijiazhuang Maternity Hospital, Shijiazhuang; and Beijing Chao-Yang Hospital, Capital Medical University, Beijing, China.

Supported by grants 0102012DFB30130 from the International Science & Technology Cooperation Program of China; 2016YFC1000300, 20161000307, 2016YFC1000803, and 2017YFC1002000 from the National Key Research and Development Program; and 81501418 from the National Natural Science Fund.

C. Guo, MD, PhD, National Research Institute for Family Planning; X. Tian, BS, The No. 1 Hospital of Shijiazhuang, Shijiazhuang Maternity Hospital; F. Han, MD, PhD, Beijing Chao-Yang Hospital, Capital Medical University; L. Liu, MD, PhD, Beijing Chao-Yang Hospital, Capital Medical University; J. Gao, MD, PhD, National Research Institute for Family Planning; X. Ma, MD, PhD, National Research Institute for Family Planning. Dr. Guo and Xin Tian contributed equally to this report.

Address correspondence to Prof. J. Gao, Center for Genetics, National Research Institute for Family Planning, 12 Dahuisi Road, Haidian, Beijing, 100081, China. E-mail: gaojianen@nrifp.org.cn

Accepted for publication September 27, 2019.

men; average age 41.4 ± 0.08 yrs) and 204 controls (122 women, 82 men; average age 59.1 ± 0.08 yrs; Table 1). All individuals were enrolled from Beijing Chaoyang Hospital and were matched for ethnic and geographic characteristics. SAPHO syndrome was diagnosed according to the criteria of Kahn, *et al*¹¹.

The study was approved by the Ethics Committee of the National Research Institute for Family Planning (registration no. 2201512), and all participants provided written informed consent.

DNA isolation. Participants' peripheral blood samples were collected into tubes containing EDTA. DNA extraction was carried out using the RelaxGene Blood DNA System (Tiangen Biotech) according to the manufacturer's instructions. Array CGH analysis was performed using $4 \times 180K$ commercial arrays (Capitalbio Technology Corp.). DNA labeling, hybridization, and washing were performed according to the manufacturer's instructions. After hybridization, arrays were scanned in a dual-laser scanner, and images were extracted and analyzed using Agilent CytoGenomics software. Probes with fewer than 2 valid replicate values or probes that showed SD above 0.1 between replicate values were excluded. Chromosomal regions with a ratio between 1.11 and 1.41 were scored as "gained" and those with a ratio of 1.41 or greater were scored as "amplified." Chromosomal regions with a ratio of between 0.84 and 0.73 were scored as "loss," whereas a second threshold for loss was set for regions showing a ratio < 0.73 .

TaqMan quantitative PCR (qPCR). TaqMan qPCR was used to identify candidate genes. Primers and fluorescence-labeled probes are shown in Table 2. qPCR was performed in a final volume of $25 \mu\text{l}$ including 20 ng of extracted DNA and $20 \mu\text{l}$ of PCR mixture. All qPCR reactions were run for 45 cycles. The Ct value of each gene was recorded and $2^{-\Delta\text{Ct}}$ ($\Delta\text{Ct} = \text{Ct target gene} - \text{Ct RNase P}$) was used to determine the CNV of each sample.

Statistical analysis. Clinical data were described as mean \pm SD. Comparison of cases and controls was by Pearson chi-square test or Fisher's exact test. Statistical analyses were performed using SPSS version 13.0 (SPSS Inc.); $p < 0.05$ was considered statistically significant.

RESULTS

We conducted a 2-stage candidate gene study to test the association of CNV and SAPHO syndrome. First, through array CGH analysis, we identified several aberrant chromosome regions shared by 2 of the core family patients with SAPHO syndrome: q11 region of chromosome 11 was deemed as

amplification, including *OR4S2*, *OR4C6*; 5 regions (q31.3 of chromosome 1, p11.22 of chromosome 8, p12 of chromosome 19, q11.23–q12.1 of chromosome 22, and q13.1 of chromosome 22) were deemed as gain, including *ADAM5P*, *APOBEC3A*, *CFHR3*, etc. Three regions were deemed as loss (q13 of chromosome 2, p22.3 of chromosome 9, p22.33 of chromosome X), including *CSF2RA*, *MIR3690*, *IL3RA*, etc. Detailed results are shown in Table 3.

In the validation stage, we chose *ADAM5*, *CSF2RA*, *IL3RA*, and 9 other genes associated with inflammation or autoimmune (*AAGAB*, *HRVR*, *IGSF9*, *PSTPIP1*, *PSTPIP2*, *LPIN2*, *IRAK3*, *NOD2*, *MEGF6*) as candidate genes. TaqMan qPCR was used to determine the variations of candidate gene copy numbers in 156 patients with SAPHO syndrome and 204 controls. Finally, 4 of 12 chosen genes were found to have significantly different copy numbers between patients with SAPHO syndrome and controls at the $\alpha = 0.05$ level, of which *CSF2RA* (ratio $\text{SAPHO/HC} = 0.597$, $p = 0.000$) and *NOD2* (ratio $\text{SAPHO/HC} = 0.470$, $p = 0.002$) had fewer gene copies, and *MEGF6* (ratio $\text{SAPHO/HC} = 1.407$, $p = 0.001$) and *ADAM5* (ratio $\text{SAPHO/HC} = 1.713$, $p = 0.000$) had more copies, compared with healthy controls. There were no differences in the other 8 candidate genes between patients with SAPHO syndrome and control samples ($p > 0.05$; Tables 4A and 4B; Figures 1 and 2). In subsequent analysis, patients with SAPHO syndrome were subgrouped according to osteoarticular and skin symptoms (Tables 4A and 4B). The results showed the gene copies of *CSF2RA*, *NOD2*, *ADAM5*, and *MEGF6* were significantly different in palmoplantar pustulosis; meanwhile, *CSF2RA* was found to be strongly correlated with osteoarticular symptoms; the results achieved statistical differences in all subgroups. As for other symptoms, perhaps because of small sample sizes, the results showed inconsistencies among different subgroups, especially in the no-skin-manifestation samples, so the results should be treated cautiously.

DISCUSSION

SAPHO syndrome is a type of rare chronic aseptic inflammation, and its complete pathogenesis is unknown. Previous investigations showed that *PSTPIP1*, *PSTPIP2*, and *LPIN2* were associated with SAPHO syndrome in mice^{5,12,13}. Other research showed that dysregulation of interleukin 1 (IL-1) signaling caused sterile osteomyelitis in *PSTPIP2*-deficient mice^{10,14}. However, genetic screening found no specific variants in *PSTPIP1*, *PSTPIP2*, *NOD2*, or *LPIN2* in patients with SAPHO syndrome⁷. We used array CGH and TaqMan qPCR to determine the role of genetic factors in the development of SAPHO. For the first time, we demonstrated that copy number loss of *CSF2RA* and *NOD2*, and copy number gain of *MEGF6* and *ADAM5*, contribute to SAPHO syndrome.

CSF2RA is located on Xp22.33 and Yp11.2, and is inherited in an autosomal recessive manner¹⁵. The *CSF2RA* protein functions as the alpha subunit of the heterodimeric

Table 1. General characteristics of patients with SAPHO syndrome (n = 156) in this study.

Characteristic	N (%)
Sex	
Female	92 (59)
Male	64 (41)
Skin manifestations	
None	7 (4.5)
Palmoplantar pustulosis	102 (65.4)
Severe acne	17 (10.9)
Palmoplantar pustulosis + psoriasis vulgaris	25 (16.0)
Osteoarticular symptoms	
None	2 (1.3)
Anterior chest wall	35 (22.4)
Anterior chest wall + spine	57 (36.5)
Anterior chest wall + peripheral skeleton	26 (16.7)
Anterior chest wall + spine + peripheral skeleton	36 (23.1)

SAPHO: synovitis, acne, pustulosis, hyperostosis, osteitis syndrome.

Table 2. Candidate genes and primers used in this study.

Gene		Forward Primer	Probe	Reverse Primer
AAGAB	170684	TGC AGG GTC CTT TGA AAT CC	5'-FAM-TGA CAA CAT TACCATGTAAAGTGCAAAACC-3'BHQ	CTC ACA GCA TCA TTG GAA GTC AC
HRNR	705321	TTC GTT AGC CAG ATG TAT TGT CC	5'-FAM-AAT CCA CTT TCT TGT TAT GGT CTC GAT CC-3'BHQ	TGA TTT TAT TAC GAG CCT GAA CC
IGSF9	2060483	CTA ATG GTG GGG TTT CTT CTG G	5'-FAM-CTC GGG TGT TAC ATG AGC CAA GCC-3'BHQ	GGG ATG TCC TTC TGA TCA GCT C
PSTPIP2	1782873	CAA CGC AAA ACT GGA CAG ATT C	5'-FAM-AAG AAA CAG GAC ATG AAG AGG CAG GC-3'BHQ	CTG CAT TCT TCT GGG AGG AGT AG
LPIN2	149540	TGA AAT CTG GCA TAC AAG TGA GTA G	5'-FAM-CCT TCA CAG TCA CAA TCA CCT GCC C-3'BHQ	GGT TAA TGC CCT TGT AGA GTT CC
IL3RA	2139684	GCT CAG GGA ACA CGT ATC GG	5'-FAM-CGA CTC TCC AGC GGT TCT CAA AGT TC-3'BHQ	CTA AAG CCG TGG TAT CAC AGA AC
CSF2RA	911684	AAG GGG TGT GTA AAG GAC AGG	5'-FAM-CTG GGT TCC ACG ACT CTG TTC TTC TTG T-3'BHQ	CAC CAG CCT CTA GTC TCA ATG TG
IRAK3	197978	TCT GTG GAA TGG TGG GAA CTAG	5'-FAM-ACA TGG TGG TCT TCG GCT TTG GTT-3'BHQ	AAG GAA AGT AGG GGG AAG ATA AG
NCR2	161264	GGC CTT GAC CTA TGC GTT AC	5'-FAM-CAG GCA CAA TCC AAG GCT CAG GTA CT-3'BHQ	GAA GCC TCC TTA CAC CAG CC
Rnase P	150941	GTT CAG AGG TGG TGC TAG AAA TG	5'-FAM-AAC GTT CTA GTG GCA GCA GTG GA-3'BHQ	TTT CGT CTT CGT GCT GAT GAG
NOD2	2061083	CCC TTT GAG CTC TGACAT TCT G	5'-FAM-ATT TGG GAA GAC ATG TTG GTT GGA TAT AC-3'BHQ	TGT CCC TAA CCT GCA ATC AAT G
PSTPIP1	148556	TCC TGG GCT GTC TAC TAA ATG G	5'-FAM-AAA GAG ACT GCC CAA AGC CAC TCA AT-3'BHQ	GAC CAC AAC ACA GAT CTC GGG
MEGF6	2062656	TGT CCT TCC TAA ATA TCA GTC TTC AC	5'-FAM-CCA GCA AAT AAC TAA GAC AGT GAG GGG T-3'BHQ	AGT CTC CTC CTT CTG AAC CAG C
ADAM5	198273	GCA GGC AGA AGA ACA TTG AAA G	5'-FAM-TCC TGC CCT TGA ACA TCG AAC TCC-3'BHQ	CAG TCC CAA AAT CTC AAA AGT AGG

receptor for colony-stimulating factor 2, a cytokine that controls the production, differentiation, and function of granulocytes and macrophages^{16,17}. We demonstrated that the *CSF2RA* copy number was significantly decreased in patients with SAPHO syndrome, which would likely decrease its interaction with granulocyte-macrophage colony-stimulating factor (GM-CSF) and its receptor, resulting in an accumulation of GM-CSF in the peripheral blood of patients. Previous research showed GM-CSF played a pivotal role in regulating inflammatory networks, and aberrant expression of GM-CSF was found in some autoimmune diseases, including rheumatoid arthritis^{18,19} and synovitis^{20,21,22}. In synovitis, abundant GM-CSF can promote joint inflammation by secreting large amounts of tumor necrosis factor- α (TNF- α) and IL-6²³. Indeed, elevated IL-6 and TNF concentrations have been observed in the serum of patients with SAPHO syndrome²⁴, while the use of anti-TNF agents has proven a valid alternative for patients unresponsive to conventional treatments, including nonsteroidal antiinflammatory drugs, corticosteroids, disease-modifying antirheumatic drugs, and bisphosphonates²⁵. GM-CSF also enhances IL-1 β production by macrophages by regulating lipopolysaccharide (LPS)-mediated pro-IL-1 β expression²⁶. Excessive IL-1 β can lead to osteitis, a major feature of SAPHO syndrome, and enhance mesenchymal cell differentiation in osteoblasts²⁷. SAPHO syndrome has been shown to partly depend on genetically encoded overproduction of IL-1 β ²⁷. Additionally, excessive IL-1 β triggers an innate immune response in the skin²⁸, including *in vivo* inflammatory responses to *Propionibacterium acnes*, another major phenotype of SAPHO syndrome, which is in accord with the acne phenotype of SAPHO syndrome. We therefore speculate that the excess GM-CSF induced by decreased CSF2RA is a major cause of SAPHO syndrome development.

NOD2 is located in chromosome 16, a region highly associated with some systemic autoinflammatory diseases. *NOD2* was the first susceptibility gene identified in inflammatory bowel disease²⁹. Previous results described genetic variants of *NOD2* in a series of systemic autoinflammatory diseases, including Crohn disease, Blau syndrome, and Yao syndrome³⁰. *NOD2* is also associated with skin diseases³¹, and it plays a pivotal role in the immune response to intracellular bacterial LPS by recognizing the muramyl dipeptide and activating the NF κ B protein³¹; and delayed and ineffective recognition of a localized bacterial infection in the absence of *NOD2* protein can lead to increased bacterial load and a delayed, more severe local inflammatory response³². Given its important role in autoinflammatory diseases, much effort has been made to determine its relationship with SAPHO syndrome. However, no genetic variants were found: we observed that *NOD2* copy number losses instead of genetic variants were involved in patients with SAPHO syndrome, and this represents significant progress in SAPHO research.

Table 3. General information on candidate genes in the nuclear family in this study.

Chr	Cytoband	Probes	Amp	Gain	Loss	Del	p	Genes
1	q31.3	3	0	0.79	0	0	2.21E-10	<i>CFHR3, CFHR1</i>
2	q13	10	0	0	-0.46	0	2.56E-13	<i>MALL, NPHP1, NCRNA00116</i>
8	p11.22	12	0	0.97	0	0	7.86E-51	<i>ADAM5P, ADAM3A</i>
9	p22.3	17	0	0	-0.53	0	5.35E-25	<i>TTC39B</i>
11	q11	4	6.54	0	0	0	4.61E-10	<i>OR4S2, OR4C6</i>
19	p12	5	0	0.74	0	0	4.53E-12	
22	q11.23-q12.1	20	0	0.83	0	0	7.48E-64	<i>IGLL3P, LRP5L, CRYBB2P1</i>
22	q13.1	3	0	0.89	0	0	3.07E-12	<i>APOBEC3A, APOBEC3B</i>
X	p22.33	17	0	0	-0.43	0	3.78E-18	<i>CSF2RA, MIR3690, IL3RA</i>

Chr: chromosome; Amp: amplification; Del: deletion.

Table 4A. Copy number variation ratio of candidate genes in patients with SAPHO syndrome and healthy controls.

	<i>PSTPIP1</i>	<i>IRAK3</i>	<i>AAGAB</i>	<i>ADAM5</i>	<i>CSF2RA</i>	<i>IL3RA</i>
Controls, n = 204	0.71 ± 0.19	1.01 ± 0.19	0.62 ± 0.12	1.96 ± 0.67	1.05 ± 0.41	0.65 ± 0.17
Cases, n = 156	0.66 ± 0.30	0.98 ± 0.40	0.52 ± 0.12	3.35 ± 3.50	0.62 ± 0.17	0.58 ± 0.32
p	0.575	0.659	0.382	0.000	0.000	0.508
ACW	0.62 ± 0.11	1.01 ± 0.58	0.50 ± 0.10	4.60 ± 5.03*	0.59 ± 0.12**	0.65 ± 0.49
ACW + peripheral skeleton	0.64 ± 0.10	0.93 ± 0.19	0.52 ± 0.10	3.28 ± 3.54*	0.60 ± 0.14**	0.55 ± 0.21
ACW + spine	0.64 ± 0.12	0.95 ± 0.12	0.51 ± 0.13	4.02 ± 3.05**	0.65 ± 0.16**	0.52 ± 0.14
ACW + spine + peripheral skeleton	0.64 ± 0.11	0.94 ± 0.19	0.52 ± 0.07	2.62 ± 2.60	0.67 ± 0.13**	0.61 ± 0.19
PPP	0.64 ± 0.10	0.98 ± 0.35	0.51 ± 0.09	3.04 ± 2.81**	0.63 ± 0.14**	0.58 ± 0.23
Psoriasis vulgaris	0.67 ± 0.13	0.92 ± 0.28	0.54 ± 0.13	4.39 ± 5.46	0.64 ± 0.18	0.65 ± 0.47
Severe acne	0.58 ± 0.13	0.86 ± 0.22	0.47 ± 0.08	4.69 ± 4.55*	0.57 ± 0.11**	0.54 ± 0.29
None	0.59 ± 0.08	0.97 ± 0.08	0.60 ± 0.15	3.08 ± 2.99	0.67 ± 0.06	0.51 ± 0.07

*p < 0.05. **p < 0.01. SAPHO: synovitis, acne, pustulosis, hyperostosis, osteitis; ACW: anterior chest wall; PPP: palmoplantar pustulosis.

Table 4B. Copy number variation ratio of candidate genes in patients with SAPHO syndrome and healthy controls.

	<i>LPIN2</i>	<i>NOD2</i>	<i>PSTPIP2</i>	<i>IGSF9</i>	<i>MEGF6</i>	<i>HRVR</i>
Controls, n = 204	1.02 ± 0.29	1.18 ± 1.36	0.86 ± 0.26	0.64 ± 0.19	0.37 ± 0.19	0.64 ± 0.17
Cases, n = 156	1.07 ± 0.52	0.55 ± 0.22	0.79 ± 0.37	0.78 ± 0.64	0.52 ± 0.25	0.59 ± 0.40
p	0.613	0.005	0.461	0.146	0.002	0.418
ACW	0.99 ± 0.22	0.57 ± 0.21	0.78 ± 0.35	0.90 ± 0.27	0.56 ± 0.21*	0.70 ± 0.50
ACW + peripheral skeleton	1.02 ± 0.17	0.53 ± 0.11**	0.73 ± 0.13	0.60 ± 0.09	0.47 ± 0.21	0.47 ± 0.14
ACW + spine	1.05 ± 0.18	0.54 ± 0.12	0.75 ± 0.15	0.72 ± 0.07	0.58 ± 0.27**	0.56 ± 0.10
ACW + spine + peripheral skeleton	1.04 ± 0.22	0.56 ± 0.11**	0.79 ± 0.11	0.58 ± 0.31	0.63 ± 0.29**	0.52 ± 0.27
PPP	1.01 ± 0.16	0.54 ± 0.14**	0.76 ± 0.14	0.63 ± 0.21	0.53 ± 0.26*	0.52 ± 0.21
Psoriasis vulgaris	1.08 ± 0.30	0.59 ± 0.16	0.76 ± 0.17	0.75 ± 0.17	0.60 ± 0.20*	0.57 ± 0.04
Severe acne	1.03 ± 0.20	0.57 ± 0.10	0.74 ± 0.40	0.76 ± 0.30	0.50 ± 0.19	0.64 ± 0.47
None	1.02 ± 0.13	0.50 ± 0.06	0.77 ± 0.14	0.70 ± 0.16	0.95 ± 0.23	0.51 ± 0.17

*p < 0.05. **p < 0.01. SAPHO: synovitis, acne, pustulosis, hyperostosis, osteitis; ACW: anterior chest wall; PPP: palmoplantar pustulosis.

We also showed that the copy numbers of *MEGF6* and *ADAM5* were higher in patients with SAPHO syndrome than in healthy controls. Although we inferred that these differences may increase the risk of acne and pustulosis, little is known about *MEGF6* and *ADAM5* proteins, so further study is required.

We established the first association between CNV in *CSF2RA*, *NOD2*, *MEGF6*, and *ADAM5* and SAPHO syndrome. These findings may provide insight into the pathogenesis of SAPHO syndrome, and perhaps a basis

for diagnosis and treatment. However, further studies are required to elucidate the pathogenesis of SAPHO syndrome.

ACKNOWLEDGMENT

We thank all participants for their contributions to the research. We thank Sarah Williams, PhD, from Liwen Bianji, Edanz Group China, for editing the English text of a draft of this report.

REFERENCES

- Magrey M, Khan MA. New insights into synovitis, acne, pustulosis, hyperostosis, and osteitis (SAPHO) syndrome. *Curr Rheumatol Rep* 2009;11:329-33.




Figure 1. CNV ratio of candidate genes in SAPHO patients and healthy controls (Part A). HC: healthy controls; CNV: copy number variations; SAPHO (synovitis, acne, pustulosis, hyperostosis, osteitis) syndrome.

- Kerrison C, Davidson JE, Cleary AG, Beresford MW. Pamidronate in the treatment of childhood SAPHO syndrome. *Rheumatology* 2004;43:1246-51.
- Matzaroglou C, Velissaris D, Karageorgos A, Marangos M,

- Panagiotopoulos E, Karanikolas M. SAPHO syndrome diagnosis and treatment: Report of five cases and review of the literature. *Open Orthop J* 2009;3:100-6.
- Hayem G. SAPHO syndrome. *Rev Prat* 2004;54:1635-6.




Figure 2. CNV ratio of candidate genes in SAPHO patients and healthy controls (Part B). HC: healthy controls; CNV: copy number variations; SAPHO (synovitis, acne, pustulosis, hyperostosis, osteitis) syndrome.

5. Ferguson PJ, Bing X, Vasef MA, Ochoa LA, Mahgoub A, Waldschmidt TJ, et al. A missense mutation in pstpip2 is associated with the murine autoinflammatory disorder chronic

multifocal osteomyelitis. Bone 2006;38:41-7.

6. Liao HJ, Chyuan IT, Wu CS, Lin SW, Chen KH, Tsai HF, et al. Increased neutrophil infiltration, IL-1 production and a

- SAPHO syndrome-like phenotype in PSTPIP2-deficient mice. *Rheumatology* 2015;54:1317-26.
7. Hurtado-Nedelec M, Chollet-Martin S, Chapeton D, Hugot JP, Hayem G, Gerard B. Genetic susceptibility factors in a cohort of 38 patients with SAPHO syndrome: a study of PSTPIP2, NOD2, and LPIN2 genes. *J Rheumatol* 2010;37:401-9.
 8. Zarrei M, MacDonald JR, Merico D, Scherer SW. A copy number variation map of the human genome. *Nat Rev Genet* 2015;16:172-83.
 9. Sebat J, Lakshmi B, Troge J, Alexander J, Young J, Lundin P. Large-scale copy number polymorphism in the human genome. *Science* 2004;305:525-8.
 10. Sharma M, Ferguson PJ. Autoinflammatory bone disorders: update on immunologic abnormalities and clues about possible triggers. *Curr Opin Rheumatol* 2013;25:658-64.
 11. Kahn MF, Hayem F, Hayem G, Grossin M. Is diffuse sclerosing osteomyelitis of the mandible part of the synovitis, acne, pustulosis, hyperostosis, osteitis (SAPHO) syndrome? Analysis of seven cases. *Oral Surg Oral Med Oral Pathol* 1994;78:594-8.
 12. Cassel SL, Janczy JR, Bing X, Wilson SP, Olivier AK, Otero JE, et al. Inflammasome-independent IL-1 β mediates autoinflammatory disease in Pstpip2-deficient mice. *Proc Natl Acad Sci USA* 2014;111:1072-7.
 13. Colina M, Pippucci T, Moro MA, Marconi C, Magini P, Ciano G, et al. Synovitis, acne, pustulosis, hyperostosis, osteitis (SAPHO) syndrome: Is PTPN22 involved? *Clin Exp Rheumatol* 2012;30:451.
 14. Ferguson PJ, Laxer RM. New discoveries in CRMO: IL-1 β , the neutrophil, and the microbiome implicated in disease pathogenesis in Pstpip2-deficient mice. *Semin Immunopathol* 2015;37:407-12.
 15. Lacey DL, Timms E, Tan HL, Kelley MJ, Dunstan CR, Burgess T, et al. Osteoprotegerin ligand is a cytokine that regulates osteoclast differentiation and activation. *Cell* 1998;93:165-76.
 16. Helena Mangs A, Morris BJ. The human pseudoautosomal region (PAR): origin, function and future. *Curr Genomics* 2007;8:129-36.
 17. Tanaka S, Takahashi N, Udagawa N, Tamura T, Akatsu T, Stanley ER, et al. Macrophage colony-stimulating factor is indispensable for both proliferation and differentiation of osteoclast progenitors. *J Clin Invest* 1993;91:257-63.
 18. Guo X, Higgs BW, Bay-Jensen AC, Wu Y, Karsdal MA, Kuziora M, et al. Blockade of GM-CSF pathway induced sustained suppression of myeloid and T cell activities in rheumatoid arthritis. *Rheumatology* 2018;57:175-84.
 19. Avci AB, Feist E, Burmester GR. Targeting GM-CSF in rheumatoid arthritis. *Clin Exp Rheumatol* 2016;34:39-44.
 20. Reynolds G, Gibbon JR, Pratt AG, Wood MJ, Coady D, Raftery G, et al. Synovial CD4+ T-cell-derived GM-CSF supports the differentiation of an inflammatory dendritic cell population in rheumatoid arthritis. *Ann Rheum Dis* 2016;75:899-907.
 21. Klimiuk PA, Yang H, Goronzy JJ, Weyand CM. Production of cytokines and metalloproteinases in rheumatoid synovitis is T cell dependent. *Clin Immunol* 1999;90:65-78.
 22. Cook AD, Pobjoy J, Steidl S, Durr M, Braine EL, Turner AL, et al. Granulocyte-macrophage colony-stimulating factor is a key mediator in experimental osteoarthritis pain and disease development. *Arthritis Res Ther* 2012;14:R199.
 23. Papatriantafyllou M. Cytokines: GM-CSF in focus. *Nat Rev Immunol* 2011;11:370-1.
 24. Firinu D, Garcia-Larsen V, Manconi PE, Del Giacco SR. SAPHO syndrome: current developments and approaches to clinical treatment. *Curr Rheumatol Rep* 2016;18:35.
 25. Hampton SL, Youssef H. Successful treatment of resistant SAPHO syndrome with anti-TNF therapy. *BMJ Case Rep* 2013;2013:bcr2012007161.
 26. Ahn S, Jeong D, Oh SJ, Ahn J, Lee SH, Chung DH. GM-CSF and IL-4 produced by NKT cells inversely regulate IL-1 β production by macrophages. *Immunol Lett* 2017;182:50-6.
 27. Berthelot JM, Corvec S, Hayem G. SAPHO, autophagy, IL-1, FoxO1, and *Propionibacterium* (*Cutibacterium*) *acnes*. *Joint Bone Spine* 2018;85:171-6.
 28. Kistowska M, Gehrke S, Jankovic D, Kerl K, Fettelschoss A, Feldmeyer L, et al. IL-1 β drives inflammatory responses to *propionibacterium acnes* in vitro and in vivo. *J Invest Dermatol* 2014;134:677-85.
 29. DeGruttola AK, Low D, Mizoguchi A, Mizoguchi E. Current understanding of dysbiosis in disease in human and animal models. *Inflamm Bowel Dis* 2016;22:1137-50.
 30. Yao Q, Li E, Shen B. Autoinflammatory disease with focus on NOD2-associated disease in the era of genomic medicine. *Autoimmunity* 2019;52:48-56.
 31. Wong CK, Chu IM, Hon KL, Tsang MS, Lam CW. Aberrant expression of bacterial pattern recognition receptor NOD2 of basophils and microbicidal peptides in atopic dermatitis. *Molecules* 2016;21:471.
 32. Hruz P, Zinkernagel AS, Jenikova G, Botwin GJ, Hugot JP, Karin M, et al. NOD2 contributes to cutaneous defense against *Staphylococcus aureus* through alpha-toxin-dependent innate immune activation. *Proc Natl Acad Sci USA* 2009;106:12873-8.