

Journal of Rheumatology

Association between centromere and topoisomerase specific immune responses and the degree of microangiopathy in Systemic Sclerosis

Nina M. van Leeuwen¹, Corrie M. Wortel¹, Cynthia M. Fehres¹, Jaap. A Bakker², Hans. U. Scherer¹, René E.M Toes¹, Tom W.J Huizinga¹, Jeska K. de Vries-Bouwstra¹

¹Department of Rheumatology, Leiden University Medical Center, Leiden, the Netherlands

²Department of Clinical Chemistry and Laboratory Medicine, Leiden University Medical Center, Leiden, the Netherlands.

Keywords:

- Systemic Sclerosis
- Pathophysiology
- Autoantibodies
- Microangiopathy

Funding: No source of support in the form of grants or industrial support was applicable for this study. Co-author C.M Wortel would like to thank ReumaNederland for her scholarship.

Disclosure statement: All authors have declared that they have no conflicts of interest.

Titles authors:

Nina M. van Leeuwen(MD) Corrie M. Wortel (MSc), Cynthia M Fehres (pHD), Jaap. A Bakker (MSc, PhD) Hans. U. Scherer (MD, PhD) René E.M Toes (Professor, MSc, PhD) Tom W.J Huizinga (Professor,MD, PhD) Jeska K. de Vries-Bouwstra (MD, PhD)

Corresponding Author:

Nina M. van Leeuwen, department of Rheumatology Leiden University Medical Center, C1-R, PO Box 9600, 2300 RC, Leiden, the Netherlands. E-mail: n.m.van_leeuwen@lumc.nl

Running head: Microangiopathy and autoantibodies

Objectives

Autoreactive antibody responses, including the use of several isotypes of autoantibodies, have been shown to associate with clinical outcome in several rheumatic autoimmune diseases. The goal of this study was 1) to evaluate whether anti-centromere antibody(ACA) and anti-topoisomerase antibody(ATA) specific isotype expression and 2) organ involvement associate with the degree of microangiopathy in SSc.

Methods

ACA and ATA IgG, IgM and IgA levels were measured in baseline serum samples of ACA IgG+ and ATA IgG+ SSc patients. The degree of microangiopathy was determined based on nailfold videocapillaroscopy images at the same time point. Logistic regression analyses with autoantibodies, clinical characteristics, isotype expression and ACA resp. ATA IgG, IgM and IgA levels as independent and NVC pattern as dependent variable were performed.

Results

In 164 patients isotype levels and degree of microangiopathy were evaluated. Logistic regression confirmed the association of the degree of microangiopathy with the presence of digital ulcers(OR 3.1 (1.4-6.6)), interstitial lung disease(OR 3.2 (1.1-9.7)) and pulmonary arterial hypertension(OR 5.25 (1.69-16.36)). ATA positivity was associated with more severe microangiopathy(OR 2.09 (1.05-4.13)). Patients that solely expressed ACA IgG showed a trend towards less severe microangiopathy compared to patients expressing also ACA IgM and/or IgA, levels of ACA IgG and ATA IgM associated with microangiopathy severity.

Conclusions

We observed an association between ACA and ATA responses and the degree of microangiopathy in SSc. These findings might indicate that the breath of the autoimmune response as reflected by autoantibody production and microvascular damage interact in the pathophysiology of SSc.

Significance and Innovations.

- We confirm that severe microangiopathy is associated with the presence of digital ulcers, pulmonary involvement and PAH in SSc.
- We confirm that anti-topoisomerase antibody positive patients have more severe microangiopathy compared to anti-centromere antibody positive patients.
- We observe an association between ATA and ACA-isotype expression and the degree of microangiopathy.
- Our observations contribute to insights in the interaction between the specific ATA and ACA expression and microangiopathy in SSc.

Introduction

Systemic Sclerosis (SSc) is characterised by the triad of microvascular damage, dysregulation of innate and adaptive immunity and generalized fibrosis (1). The pathogenesis of SSc is still not completely elucidated, and the primary cause of SSc remains to be determined (2). About 95% of SSc patients have antinuclear autoantibodies (ANAs). These autoantibodies contribute to the disease classification, are associated with specific clinical manifestations and are therefore important tools for disease prognostication (3). Anti-centromere antibodies (ACA) and anti-topoisomerase antibodies (ATA) are the two most common ANAs in patients with SSc (3). Of these, ACA associates with a relatively mild disease course, while ATA associates with more severe disease including diffuse skin and lung involvement. This clear association with a typical clinical phenotype suggests that the immune response reflected by ATA or ACA production is closely linked to disease pathophysiology. The exact pathogenicity of ATA or ACA, however, remains unclear (4).

A second important diagnostic and prognostic tool in SSc is nailfold videocapillaroscopy (NVC) which is an investigation that determines the degree of microangiopathy by using standardised magnification to visualize the capillaries in the nailfold. In SSc, specific patterns of capillary changes and the degree of these changes have been defined extensively (5). More severe microangiopathy is associated with worse disease in SSc patients, in recent studies an association between NVC pattern and organ involvement and disease progression was found (6-12). In addition, an association between NVC patterns and specific autoantibodies was also described (13). NVC therefore can be seen as an important biomarker to predict severe complications in SSc (14).

Some studies suggest that autoantibody production is secondary to vasculopathy, and as such specific autoantibodies can be viewed as bystanders in disease pathogenesis (2). However, other studies suggest that circulating autoantibodies may be directly implicated in the disease process. Higher levels of ATA have been shown to associate with the development of organ involvement (15). An association between autoantibody specific isotypes and disease severity has been shown, with higher ATA IgM levels in ATA + SSc patients who showed disease progression, and higher ACA IgG and ACA IgM levels in ACA+ SSc patients with a more severe disease (16, 17). Besides, a study performed by Ahmed et al. (18) demonstrated that SSc sera containing ACA or ATA can trigger fibrillin expression in human dermal endothelial cells and induce cell apoptosis, and Shen et al. (19) concluded that the pathognomonic ACA and ATA in SSc accelerate vascular endothelial cell senescence and functional impairment inducing Raynaud's Phenomenon (RP).

Together, these studies implicate a possible association between ACA, ATA, and possibly specific isotype levels and vasculopathy. This association has not yet been evaluated in SSc.

As the presentation of SSc can be very heterogeneous and prediction of the disease course is still very difficult, a better understanding of the interaction between the specific auto-immune response and the degree of microangiopathy could not only improve our insights in disease pathogenesis but could also contribute to more reliable disease prognostication, which is of utmost importance. In line with this, we hypothesized that an activated immune response, as reflected by higher ATA or ACA IgG levels associate with more severe microvascular damage.

Method

Study design and patients

SSc patients at the Leiden University Medical Center (LUMC) are included in an observational cohort study (Combined Care in Systemic Sclerosis; CCISS (12)), which is approved by the Ethics Committee (P09.003). The cohort study is designed in accordance with the ethical principles of the Declaration of Helsinki. All patients gave written informed consent. This standardised annual care pathway comprises extensive screening including: autoantibody testing, electrocardiography (EKG), thoracic echocardiography, high resolution computed tomography (HRCT), pulmonary function test, and NVC. An exercise test, 24 hour Holter EKG or right heart catheterization (RHC) are performed if indicated. In the current study, patients who fulfilled the ACR/EULAR 2013 classification criteria for SSc (20) and were positive for IgG either ATA or ACA, and had a clinical diagnosis of SSc were included. Use of current medication (calcium channel blockers, sildenafil, bosentan and Iloprost for vasoactive medication, and corticosteroids, cyclophosphamide, azathioprine, methotrexate, mycophenolate mofetil and azathioprine for immunosuppressive medication) at the time of blood sampling, baseline clinical characteristics and investigations were retracted from the database. Baseline characteristics were considered at time of inclusion in the cohort.

Nailfold videocapillaroscopy

NVC was performed at the same time point as the baseline characteristics and blood samples were collected. All images were obtained in the hospital in a comfortable room with a temperature of 22-25 degree Celsius. All fingers except for the thumbs from both hands were examined using a videocapillaroscope (LUMC: 2009-2015; Videocap 3.0, DS Medica; 2015 -2017, Inspectis pro; 2018 onwards) equipped with a probe with 200x magnification. NVC images were scored by trained observers and classified qualitatively as described previously; 'normal', 'non-specific' and

Accepted Article

'scleroderma pattern' (21). A "normal pattern" is defined as a pattern of typical hairpin-like capillaries with a regular distribution. A "non-specific pattern" is defined as a pattern with abnormalities without fulfilling the definition of a 'scleroderma pattern'. (21, 22) A 'scleroderma pattern' was defined according to the definition of Cutolo et al. and categorized in an 'early' 'active' or 'late' pattern (21, 23). The presence of giant capillaries, hemorrhages and avascularity are the main denominators in the definition of a 'scleroderma pattern'. A more severe degree of microangiopathy is defined as a late scleroderma pattern with capillary loss (< 4 capillaries per mm) as its main denominator. For the current evaluation the images were re-examined by a trained investigator (NvL). The inter-observer agreement was high for qualitative pattern determination (ICC 0.97).

ACA and ATA assay and measurements

Total immunoglobulin ATA, IgG, IgM and IgA and total immunoglobulin ACA, IgG, IgM and IgA levels of all the collected samples were measured in baseline samples by fluorescence enzyme-linked immune sorbent assay (FEIA), using the Phadia250 system (ThermoFisher Scientific, Nieuwegein, the Netherlands). The cut-off levels for ATA and ACA IgG were set at 7 units/ml (U/mL) according to the manufacturer's instructions. Fifty serum samples of non-rheumatic age and sex matched subjects were measured to establish cut-off values (mean + 2SD) for IgM and IgA isotypes of ACA and ATA. A cut-off for ACA IgA was determined at 37 aU/ml, for ATA IgA at 77 aU/ml, for ACA IgM at 13 aU/ml and for ATA IgM at 432 aU/ml. To evaluate specificity of the assay, 10 SSc patients, negative for ATA IgG were tested and all had ATA-IgM and -IgA levels below the defined cut-off, in addition 10 SSc patients negative for ACA IgG were tested for ACA IgM and ACA IgA, these levels were also below the defined cut-off. An expressed isotype was defined as a level above the cut-off value. Outliers were checked, and when necessary remeasured.

Organ involvement

Digital ulcers were present when there was clear visible tissue breakdown, both ischemic and mechanical (results of microtrauma and increased skin tension) ulcers were included in this definition. Interstitial lung disease was defined based on the combination of forced vital capacity $< 70\%$ and evidence for interstitial lung disease on HRCT. An experienced radiologist evaluated the HRCT for ground glass opacifications, reticulations and honeycombing. We chose to use a combined value including both pulmonary function and HRCT to make sure that we only classify patients with clinically relevant pulmonary involvement as having ILD. Pulmonary arterial hypertension (PAH) was defined as an increase in mean pulmonary arterial pressure (mPAP) ≥ 25 mmHg at rest as assessed

by RHC; including presence of pre-capillary PH, defined by a pulmonary capillary wedge pressure (PCWP) ≤ 15 mmHg and a PVR > 3 Wood units (WU) on RHC, in the absence of other causes of precapillary PH such as PH due to lung diseases, CTEPH or other rare diseases. To evaluate myocardial involvement we used different measurements. The Medsger subdomain reflecting myocardial involvement was evaluated, in which grade 0 represents a normal heart function, grade 1 denotes conduction abnormalities and a left ventricular ejection fraction (LVEF) between 44-49%, grade 2 signifies arrhythmias and a LVEF 40-45%, grade 3 indicates severe involvement with a LVEF $< 40\%$. As the Medsger scale mainly relies on the LVEF for determination of myocardial involvement, only using this parameter could lead to underestimation of presence of myocardial involvement as in SSc patients with myocardial involvement the LVEF is not always declined. Therefore, we additionally used a combined value where patients had to have at least two of the following: arrhythmias ($> 2\%$ ventricular or supraventricular arrhythmia, atrial fibrillation), conduction problems, decreased LVEF $< 50\%$, diastolic or systolic dysfunction, pericarditis or pericardial effusion.

Statistical analysis

No sample size calculation was performed due to the explorative character of this study. Analyses were performed by IBM SPSS version 23. All analyses were performed cross-sectionally. NVC patterns and clinical features, at the time of blood sample collection for autoantibody determination, were compared between ACA+ and ATA+ SSc patients using descriptive statistics and differences were tested for significance as appropriate. Disease duration was defined as duration since onset of Raynaud's Phenomenon (RP), as current SSc pathophysiology indicate that RP is a direct consequence of vasculopathy. We performed a Mann-Whitney U test to calculate the significance of the continuous variables. A Chi-square test was performed for the categorical variables. Fisher exact test was employed when appropriate. Binary logistic regression (univariable and multivariable), with autoantibodies, NVC pattern, isotype expression and ACA or ATA isotype levels as independent factor and organ involvement as dependent variables, and ordinal logistic regression analyses, with disease characteristics, autoantibodies and isotype expression and ACA or ATA IgG level as independent and NVC SSc pattern as dependent variables were performed. As age and disease duration can be a confounder for the association between organ involvement and degree of microangiopathy, we corrected for these variables in the multivariate analyses. In addition, variables with significant association in the univariate analysis were added as indicated. All isotype levels were transformed using log2. To adjust for multiple testing, Bonferroni correction was applied. P value < 0.05 was considered significant.

Results

Study group

A total of 231 SSc patients (129 ACA+ and 102 ATA+) were included. The included patients had a mean age of 55 years (SD 14) and median disease duration from onset first non-RP of 4 years (IQR 1-11). As expected, females represented the majority of the study population (186/231). ATA+ patients differed from ACA+ in sex ($p<0.001$), age ($p=0.01$), disease duration ($p<0.001$), diffuse cutaneous subset ($p<0.001$), and ILD ($p<0.001$). The main demographic and clinical data of the patients are summarized in Table 1. An important difference between the ATA and the ACA group that could have an influence on the degree of microangiopathy is the disease duration since onset RP, which is longer in the ACA positive patients (16 years vs 6 years) compared to the ATA+patients. Complete data on NVC patterns were available for 164/231 patients (100 ACA+ and 64 ATA+). The missing NVC were all from patients with baseline visits before 2013, in 2013 the NVC became an annual standard examination.

Organ involvement and the degree of microangiopathy

In the univariable analysis (Table 2), a more severe degree of microangiopathy (late SSc pattern) as shown by NVC, was associated with ILD (OR 4.7 (1.9-11.4)), PAH (OR 5.3 (1.7-16.4)), cardiac involvement (OR 2.9 (1.2-7.2)) and DU (OR 2.3 (1.2-4.5)). Multivariable analysis showed that a more severe degree of microangiopathy was significantly associated with ILD (OR 3.2 (1.1-9.7)), PAH (OR 3.6 (1.01-12.8)) and DU (OR 3.1 (1.4-6.6)) with correction for age, disease duration and ATA positivity.

NVC patterns in autoantibody subgroups

A late SSc pattern was numerically seen more often in ATA+ patients vs ACA+ patients (31% vs 18%; $p=0.05^*$ after Bonferroni correction; Table 1). The frequency of an early (10% ATA, 18% ACA) and active (58% ATA, 68% ACA) SSc pattern was comparable between ATA and ACA patients. In the multivariable analysis (Table 5 supplementary file), after adjustment for vasoactive medication, age, and disease duration, ATA positivity was associated with more severe microangiopathy (OR 2.97 (1.41-6.24)) compared to ACA positivity.

Isotype expression

In ACA IgG+ patients, 74% (n=95) were ACA IgA+ and 74% (n=95) ACA IgM+; 11% solely expressed ACA IgG, 16% of ACA patients were positive for IgG and IgM, 16% of the ACA patients were positive for IgG and IgA, and 58% were positive for ACA IgG, IgM and IgA. All ATA IgG + patients expressed more than one ATA specific isotype: ATA IgA+ was found in 98% (n=100), ATA IgM+ was found in 65% (n=66). Two percent of ATA IgG+ patients expressed also ATA IgM, in the absence of detectable IgA ATA, 33% of ATA IgG+ patients expressed also ATA IgA in the absence of detectable IgM ATA, and 65% expressed all three ATA isotypes.

Association ACA and ATA isotype expression with the degree of microangiopathy

As shown in Figure 1, the ACA IgG+ patients that only expressed ACA IgG, and no other ACA isotypes, more frequently showed an early pattern and less frequently a late pattern compared to ACA IgG+ patients expressing two or three ACA isotypes. The differences between the groups were not statistically significantly different. The ATA IgG+ patients, that concurrently expressed only ATA IgA showed less often a late pattern compared to ATA IgG+ patients expressing all three ATA isotypes. These differences were not statistically significant.

Numerically, ACA IgG levels were higher in ACA IgG+ patients with late SSc pattern compared to ACA IgG+ patients with an early SSc pattern (median 630 U/mL vs 200 U/mL), and ATA IgM levels were higher in ATA IgG+ patients with a late SSc pattern compared to ATA IgM in patients with an early pattern (median 1515 aU/mL vs 691 aU/mL). These results were not statistically significant (supplementary file). In the multivariable analysis with adjustment for disease duration and use of vasoactive medication antibody isotype levels were associated with degree of microangiopathy (see Table 3 and 4). For ACA IgG levels (OR 2.49 (1.04-5.83)) and for ATA IgM levels (OR 2.70 (1.06-4.22)) the associations were significantly different.

Discussion

In this study we evaluated the association between specific ATA and ACA responses and the degree of microangiopathy in a Dutch SSc cohort. We first confirmed the association between more severe microangiopathy and organ involvement including ILD, PAH and DU in patients with SSc. Secondly, we showed that ATA+ SSc patients more often have severe microangiopathy compared to ACA+ patients. Finally, our results indicate a possible association between characteristics of specific antibody responses and the degree of microangiopathy. After adjustment for possible confounders we observed a significant association between ACA IgG and ATA IgM levels and a more severe degree of microangiopathy.

In the association analysis we observed a trend for higher ATA IgM among ATA + patients with a late SSc pattern and higher ACA IgG among ACA+ patients with late SSc pattern on NVC. Only after correcting for possible confounders including disease duration these associations became significant with an association between a more severe degree of microangiopathy and levels of ACA IgG and between a more severe degree of microangiopathy and levels of ATA IgM. The rationale behind these findings is not fully understood and we can only hypothesize about possible explanations. As ATA+ and ACA+ patients display clearly different clinical phenotypes, one might hypothesize that behaviour of ATA and ACA specific isotypes differs which might impact on their roles in pathophysiology. Besides, ATA and ACA might bind to different cells or antigens which may be one reason for the differences between the two group. ATA bind to DNA topoisomerase I expressed by e.g. fibroblasts, and it may be that ATA is only pathogenic in case there is insufficient clearance of apoptotic bodies of endothelial cells containing DNA topoisomerase I. This would also fit with the observation that a more severe degree of microangiopathy is associated with organ involvement (24). ACA, can react against six different centromeric nucleoproteins, and could have a different ability in e.g. recruiting immune effector- or clearance mechanisms (25). Although ACA and ATA have been reported to react with endothelial cells, no data is published on differences between isotype binding and the effects on endothelial cells (18, 26). Another important factor in the pathogenesis of SSc including the endothelial cell damage, is the complement system. IgM and IgG have the ability to induce inflammation by activating complement, however, IgA is a weak activator for complement, this may be one of the explanations why no association between specific IgA and degree of microangiopathy was found (27).

In general, the production of IgM against protein-antigens is driven by short-lived plasmablasts derived from recently stimulated B cells and therefore presence of ATA and ACA-specific IgM suggests an ongoing active immune response. How IgM production is sustained in the presence of IgG against the same antigen is not fully understood, but similar observations have been reported for anti-citrullinated protein antibodies (ACPA) in rheumatoid arthritis (28). Production of IgM could also result from a failure of class-switching resulting in prolonged survival of IgM-secreting plasmablasts (29). In both the ATA+ patients (65%) and the ACA+ patients (74%) continuous IgM expression next to IgG is observed but proportionally IgM expression is more frequent in ACA+ patients.

Our results are partly in line with those reported by Markusse et al. and Caramaschi et al., who reported a relationship between organ involvement and more severe microangiopathy as assessed by NVC (12, 30). However, in these studies it was suggested that the presence of a specific IgG autoantibody is independent of the development of microangiopathy. Relations between

ATA/ACA isotype profile and isotype levels and disease severity have not yet been evaluated in large SSc cohorts. ATA IgG and ATA IgA levels, and presence of ATA IgM have previously been described to correlate with skin scores and with disease severity in small cohorts (15, 31). We are currently working on verifying these results in a multicentre study.

This study has some limitations that should be considered. As not all HRCTs were evaluated according to Goh, we were not able to discriminate between limited and extensive ILD. Therefore, we decided to apply a combined definition including presence of ILD on HRCT and FVC < 70% to make sure that only patients with clinically relevant ILD are classified as having ILD. Secondly, we only included patients positive for ATA IgG or ACA IgG at baseline we cannot fully exclude that there might be patients only positive for ACA IgM, ATA IgM, ACA IgA or ATA IgA that we did not include. This seems however unlikely as ATA IgA and IgM isotypes and ACA IgA and IgM isotype were absent in ANA+ SSc patients lacking a SSc specific antibody. To determine whether relevant ATA IgM or ATA IgA can be expected in SSc patients negative for ATA IgG, and vice versa for ACA IgM and ACA IgA, we have additionally measured expression of ATA IgM and ATA IgA in n=38 samples of ACA IgG positive and ATA IgG negative patients, and measured expression of ACA IgM and ACA IgA in n=46 samples of ATA IgG positive and ACA IgG negative patients. This showed that relevant expression of ATA/ACA IgM and ATA/ACA IgA is very rare in ATA/ACA IgG negative patients. Likewise, no conclusions can be drawn for the remaining antibody subgroups in SSc.

The current data were derived from a cohort study in a tertiary centre where patients with a (preliminary) diagnosis of SSc were included at presentation at the out-patient clinic and therefore treatment prior to inclusion was uncontrolled. Furthermore, we do not know how (previous) immunosuppressive medication might have influenced our results. However, in our study, ACA and ATA-specific isotype levels were not different for SSc patients who used immunosuppressive medication compared to SSc patients who did not use medication, and at baseline, the use of immunosuppressive medication was comparable between the ACA+ and ATA+ patients. Strikingly, in the multivariable analysis current use of vasoactive medication was associated with worse microangiopathy. Possibly, this association is reflecting confounding by indication, with patients with more severe microangiopathy more often suffering from DU. Finally, we decided not to include outcomes such as gastro-intestinal involvement and renal crisis in this analysis as endoscopies were not performed routinely, and the numbers for renal crisis was too low. In all patients, an HRCT and transthoracic echocardiography were performed routinely, regardless of risk factors for specific organ involvement.

In conclusion, in our analyses we observed associations between specific ATA and ACA responses and the degree of microangiopathy in patients with SSc indicating that dysregulated B cell

Accepted Article

responses and microvascular damage interact with each other in the pathophysiology of SSc. Further research is needed to confirm these observations, and to identify the possible mechanism behind this association.

References

1. Allanore Y, Simms R, Distler O, Trojanowska M, Pope J, Denton CP, et al. Systemic sclerosis. *Nat Rev Dis Primers*. 2015;1:15002.
2. Stern EP, Denton CP. The Pathogenesis of Systemic Sclerosis. *Rheum Dis Clin North Am*. 2015;41:367-82.
3. Steen VD. Autoantibodies in systemic sclerosis. *Semin Arthritis Rheum*. 2005;35:35-42.
4. Senecal JL, Henault J, Raymond Y. The pathogenic role of autoantibodies to nuclear autoantigens in systemic sclerosis (scleroderma). *J Rheumatol*. 2005;32:1643-9.
5. Cutolo M, Pizzorni C, Tuccio M, Burroni A, Craviotto C, Basso M, et al. Nailfold videocapillaroscopic patterns and serum autoantibodies in systemic sclerosis. *Rheumatology (Oxford)*. 2004;43:719-26.
6. Smith V, Decuman S, Sulli A, Bonroy C, Pietttte Y, Deschepper E, et al. Do worsening scleroderma capillaroscopic patterns predict future severe organ involvement? a pilot study. *Ann Rheum Dis*. 2012;71:1636-9.
7. Avouac J, Lepri G, Smith V, Toniolo E, Hurabielle C, Vallet A, et al. Sequential nailfold videocapillaroscopy examinations have responsiveness to detect organ progression in systemic sclerosis. *Semin Arthritis Rheum*. 2017;47:86-94.
8. Smith V, Riccieri V, Pizzorni C, Decuman S, Deschepper E, Bonroy C, et al. Nailfold capillaroscopy for prediction of novel future severe organ involvement in systemic sclerosis. *J Rheumatol*. 2013;40:2023-8.
9. Paxton D, Pauling JD. Does nailfold capillaroscopy help predict future outcomes in systemic sclerosis? A systematic literature review. *Semin Arthritis Rheum*. 2018;48:482-494.
10. Smith V, Vanhaecke A, Vandecasteele E, Guerra M, Paolino S, Melsens K, et al. New Kids on the Block in SSc-PAH: May We Futurely Nail It Additionally Down to Capillaroscopy? A Systematic Literature Review. *J Rheumatol*. 2019;47.
11. Sulli A, Paolino S, Pizzorni C, Ferrari G, Pacini G, Pesce G, et al. Progression of nailfold capillaroscopic patterns and correlation with organ involvement in systemic sclerosis: a 12 year study. *Rheumatology (Oxford)*. 2019.
12. Markusse IM, Meijs J, de Boer B, Bakker JA, Schippers HPC, Schouffoer AA, et al. Predicting cardiopulmonary involvement in patients with systemic sclerosis: complementary value of nailfold videocapillaroscopy patterns and disease-specific autoantibodies. *Rheumatology (Oxford)*. 2017;56:1081-8.
13. Sulli A, Ruaro B, Smith V, Pizzorni C, Zampogna G, Gallo M, et al. Progression of nailfold microvascular damage and antinuclear antibody pattern in systemic sclerosis. *J Rheumatol*. 2013;40:634-9.
14. Cutolo M, Sulli A, Secchi ME, Paolino S, Pizzorni C. Nailfold capillaroscopy is useful for the diagnosis and follow-up of autoimmune rheumatic diseases. A future tool for the analysis of microvascular heart involvement? *Rheumatology (Oxford)*. 2006;45:43-6.
15. Perera A, Fertig N, Lucas M, Rodriguez-Reyna TS, Hu P, Steen VD, et al. Clinical subsets, skin thickness progression rate, and serum antibody levels in systemic sclerosis patients with anti-topoisomerase I antibody. *Arthritis Rheum*. 2007;56:2740-6.
16. Boonstra M, Bakker JA, Grummels A, Ninaber MK, Ajmone Marsan N, Scherer HU, et al. FRI0451 Continuous presence of igm anti-topoisomerase i antibodies indicates an ongoing immune response in systemic sclerosis. [abstract]. *Ann. Rheum. Dis*. 2018;77:755.
17. N.M van Leeuwen JB, A. Grummels, C Wortel, S Jordan, O Distler, H fretheim, A Hoffman-Vold, H Scherer, R Toes, T Huizinga, J.K de Vries-bouwstra. Anticentromere Antibody Levels and

Isotypes Associate with Disease Severity in Systemic Sclerosis. [abstract]. *Arthritis Rheumatol*. 2019;71.

18. Ahmed SS, Tan FK, Arnett FC, Jin L, Geng YJ. Induction of apoptosis and fibrillin 1 expression in human dermal endothelial cells by scleroderma sera containing anti-endothelial cell antibodies. *Arthritis Rheum*. 2006;54:2250-62.
19. Shen CY, Li KJ, Lai PH, Yu CL, Hsieh SC. Anti-CENP-B and anti-TOPO-1-containing sera from systemic sclerosis-related diseases with Raynaud's phenomenon induce vascular endothelial cell senescence not via classical p53-p21 pathway. *Clin Rheumatol*. 2018;37:749-56.
20. van den Hoogen F, Khanna D, Fransen J, Johnson SR, Baron M, Tyndall A, et al. 2013 classification criteria for systemic sclerosis: an American College of Rheumatology/European League against Rheumatism collaborative initiative. *Arthritis Rheum*. 2013;65:2737-47.
21. Smith V, Herrick AL, Ingegnoli F, Damjanov N, De Angelis R, Denton CP, et al. Standardisation of nailfold capillaroscopy for the assessment of patients with Raynaud's phenomenon and systemic sclerosis. *Autoimmun Rev*. 2020;19:102458.
22. Smith V, Vanhaecke A, Herrick AL, Distler O, Guerra MG, Denton CP, et al. Fast track algorithm: How to differentiate a "scleroderma pattern" from a "non-scleroderma pattern". *Autoimmun Rev*. 2019:102394.
23. Cutolo M, Sulli A, Pizzorni C, Accardo S. Nailfold videocapillaroscopy assessment of microvascular damage in systemic sclerosis. *J Rheumatol*. 2000;27:155-60.
24. Henault J, Tremblay M, Clement I, Raymond Y, Senecal JL. Direct binding of anti-DNA topoisomerase I autoantibodies to the cell surface of fibroblasts in patients with systemic sclerosis. *Arthritis Rheum*. 2004;50:3265-74.
25. Ho KT, Reveille JD. The clinical relevance of autoantibodies in scleroderma. *Arthritis Res Ther*. 2003;5:80-93.
26. Mostmans Y, Cutolo M, Giddelo C, Decuman S, Melsens K, Declercq H, et al. The role of endothelial cells in the vasculopathy of systemic sclerosis: A systematic review. *Autoimmun Rev*. 2017;16:774-86.
27. Okrój M, Johansson M, Saxne T, Blom AM, Hesselstrand R. Analysis of complement biomarkers in systemic sclerosis indicates a distinct pattern in scleroderma renal crisis. *Arthritis Res Ther*. 2016;18:267.
28. Suwannalai P, Willemze A, van Toorn L, Stoeken-Rijsbergen G, Levarht N, Drijfhout JW, et al. The fine specificity of IgM anti-citrullinated protein antibodies (ACPA) is different from that of IgG ACPA. *Arthritis Res Ther*. 2011;13:R195.
29. Boes M. Role of natural and immune IgM antibodies in immune responses. *Mol Immunol*. 2000;37:1141-9.
30. Caramaschi P, Canestrini S, Martinelli N, Volpe A, Pieropan S, Ferrari M, et al. Scleroderma patients nailfold videocapillaroscopic patterns are associated with disease subset and disease severity. *Rheumatology (Oxford)*. 2007;46:1566-9.
31. Hu PQ, Fertig N, Medsger TA, Jr., Wright TM. Correlation of serum anti-DNA topoisomerase I antibody levels with disease severity and activity in systemic sclerosis. *Arthritis Rheum*. 2003;48:1363-73.
32. Cutolo M, Melsens K, Herrick AL, Foeldvari I, Deschepper E, De Keyser F, et al. Reliability of simple capillaroscopic definitions in describing capillary morphology in rheumatic diseases. *Rheumatology (Oxford)*. 2018;57:757-9.

Figure legends

Figure 1. Presence of one, two or three autoantibody specific isotypes and the distribution of early, active or late SSc pattern.

Legend: The significant values are on group level; 1A:group ACA IgG+ vs. ACA IgG+ and IgM+ p=0.37 group ACA IgG + vs ACA IgG+ and IgM+ p=0.25, group ACA IgG+ vs ACA IgG+, IgM+ and IgA+ p=0.28 (no significant difference in NVC pattern prevalence),. 1B; All ATA IgG+ patients expressed at least one additional ATA isotype. Two patients expressed ATA IgG+ and ATA IgM+ both with an active NVC pattern. Prevalence of NVC patterns was not significantly different between ATA IgG+ and IgA+ vs ATA IgG+, IgM+ and IgA+ patients (p= value 0.39).

Figure 2. Nailfold videocapillaroscopy (NVC) images made with the Inspectis pro.

Legend: Examples of NVC images with a 1 mm grid. # hemorrhages, \$ dilation > 30uM, * giant (dilation > 50um), @ 'abnormal shapes' (= 'neoangiogeneses') (32).




Figure 1. Presence of one, two or three autoantibody specific isotypes and the distribution of early, active or late SSc pattern.

The significant values are on group level; 1A: group ACA IgG+ vs. ACA IgG+ and IgM+ $p=0.37$ group ACA IgG+ vs ACA IgG+ and IgM+ $p=0.25$, group ACA IgG+ vs ACA IgG+, IgM+ and IgA+ $p=0.28$ (no significant difference in NVC pattern prevalence),.

1B; All ATA IgG+ patients expressed at least one additional ATA isotype. Two patients expressed ATA IgG+ and ATA IgM+ both with an active NVC pattern. Prevalence of NVC patterns was not significantly different between ATA IgG+ and IgA+ vs ATA IgG+, IgM+ and IgA+ patients ($p=0.39$).




Figure 2. Examples of nailfoldvideocapillaroscopy images with a 1 mm grid. # hemorrhages, \$ dilation > 30uM, * giant (dilation > 50um), @ 'abnormal shapes' ('neoangiogenesis') (32).

Table 1. Baseline characteristics of included ACA+ and ATA+ SSc patients

	Total group	ACA+	ATA+
Demographic	n= 231	n= 129	n= 102
Female, n (%)	186 (81)	116 (90)	70 (69)
Age, mean (SD)	55 (14)	58 (13)	51 (14)
Smoking, ever, n (%)	138 (60)	76 (59)	62 (61)
Disease duration			
Since RP, median (IQR)	10 (4-20)	16 (6-26)	6 (2-13)
Since non RP, median (IQR)	4 (1-11)	5 (1-12)	3 (1-9)
Organ involvement			
DcSSc, n (%)	49 (21)	2 (2)	47 (46)
Puffy fingers, n (%)	74 (33)	38 (30)	36 (37)
Sclerodactylie, n (%)	151 (67)	71 (55)	80 (82)
mRSS, median (IQR)	4 (2-6)	3 (1-5)	6 (2-12)
Pitting Scars, n (%)	106 (46)	59 (46)	47 (46)
Teleangiectasia, n (%)	152 (66)	108 (84)	44 (44)
Digital ulcers, n (%)	43 (19)	30 (23)	13 (13)
DLCO % predicted mean (SD)	65 (17)	70 (17)	62 (17)
FVC % predicted, mean (SD)	92 (21)	92 (21)	91 (20)
ILD on HRCT, n (%)	55 (24)	11 (9)	44 (43)
PAH, n(%)	20 (13)	10 (8)	10 (10)
Nailfold videocapillaroscopy			
NVC early, n (%)	22 (15)	16 (18)	6 (10)
NVC active, n (%)	90 (63)	56 (68)	34 (58)
NVC late, n (%)*	33 (23)	15 (18)	18 (31)
NVC capillary loss < 7mm, n (%)	114 (70)	64 (65)	50 (78)
Medication			
CYC, n (%)	39 (17)	19 (15)	20 (20)
HSCT, n (%)	14 (7)	7 (7)	7 (7)
Iloprost/bosentan, n (%)	17 (12)	14 (11)	13 (13)
Methotrexate ever, n (%)	44 (19)	23 (18)	21 (20)
ACA or ATA characteristics			
IgA positivity, n (%)	-	95 (74)	100 (98)
IgA level (aU/mL), median (IQR)	-	73 (34-146)	2778 (933-8368)
IgM positivity, n (%)	-	95 (74)	66 (65)
IgM level (aU/mL), median (IQR)	-	65 (4-561)	822 (286-2162)
IgG level (U/mL), median (IQR)	-	478 (186-1031)	484 (170-934)

Baseline characteristics. CYC=cyclophosphamide, dcSSc= diffuse cutaneous SSc, DLCO= carbon monoxide diffusing capacity, FVC= forced vital capacity, HSCT= stem cell transplantation, ILD= interstitial lung disease, mRSS= Modified Rodnan Skin Score, NVC= nailfold video capillaroscopy, PAH= pulmonary arterial hypertension, RP= Raynaud phenomenon.

Table 2. ATA and more severe microangiopathy is associated with organ involvement

	Univariable OR (95% CI)				Multivariable OR (95% CI)			
	Interstitial lung disease	Cardiac involvement	PAH	DU	Interstitial lung disease	Cardiac involvement	PAH	DU
Male	1.71 (0.84-3.48)	0.59 (0.26-1.34)	1.04 (0.33-3.27)	1.32 (0.55-3.20)	-	-	-	-
Age	1.01 (0.99-1.04)	1.07 (1.03-1.10)	1.08 (1.03-1.13)	0.99 (0.97-1.02)	1.01 (0.95-1.07)	1.06 (1.01-1.11)	1.09 (1.02-1.18)	0.98 (0.95-1.01)
Disease duration since RP	0.99 (0.95-1.01)	1.04 (0.98-1.03)	1.00 (0.99-1.05)	1.01 (0.99-1.03)	0.98 (0.89-1.07)	1.01 (0.97-1.05)	1.02 (0.94-1.07)	0.99 (0.96-1.03)
Disease duration since NR	1.03 (0.99-1.06)	1.01 (0.97-1.05)	1.01 (0.97-1.07)	1.04 (1.01-1.08)	-	-	-	-
ATA	6.68 (1.87-23.94)	0.96 (0.47-1.95)	1.29 (0.52-3.24)	2.07 (1.02-4.22)	13.34 (2.87-52.61)	1.28 (0.38-4.31)	3.92 (0.68-22.46)	0.36 (0.14-0.92)
NVC SSc pattern	3.59 (1.75-15.91)	2.95 (1.20-7.23)	5.85 (1.90-18.65)	2.28 (1.17-4.47)	3.41 (1.11-10.61)	2.17 (0.86-5.48)	5.58 (2.05-17.81)	3.07 (1.43-6.60)
Immunosuppressiva	4.64 (1.57-13.66)	1.08 (0.50-2.31)	0.53 (0.26-1.99)	0.46 (0.20-1.07)	1.79 (0.45-7.01)	-	-	-

In the multivariable logistic regression autoantibody, age, disease duration (since onset RP) and parameters with significant association in univariate analysis were included. Interstitial lung disease (ILD) was defined as ILD on HRCT and a FVC < 70% of predicted. NVC pattern was entered as ordinal variable, in order: early, active or late.

ATA=anti-topoisomerase, CI=confidence interval, DU=digital ulcers, NR=non-Raynaud, NVC= nailfold videocapillaroscopy, OR= odds ratio, PAH= pulmonary arterial hypertension, RP= Raynaud Phenomenon, SSc= systemic sclerosis .

This accepted article is protected by copyright. All rights reserved.

Table 3. ACA IgG levels are associated with more severe microangiopathy after adjustment for disease duration and vasoactive medication.

	Univariable OR (95% CI)	Multivariable OR (95% CI)
	NVC SSc pattern	NVC SSc pattern
Male	0.69 (0.17-2.77)	-
Age	1.02 (0.99-1.05)	-
Disease duration since RP	1.01 (0.97-1.04)	1.01 (0.98-1.05)
Disease duration since NR	1.01 (0.95-1.07)	
Autoantibody specific IgM positive	1.21 (0.42-3.52)	-
Autoantibody specific IgA positive	1.13 (0.45-2.81)	-
Autoantibody specific IgG levels, U/ml	2.24 (0.99-5.05)	2.46 (1.04-5.83)
Autoantibody specific IgM levels, aU/ml	0.99 (0.60-1.65)	-
Autoantibody specific IgA levels, aU/ml	1.51 (0.67-3.37)	-
Autoantibody specific isotype expression	0.80 (0.33-1.91)	-
Immunosuppressive medication	0.64 (0.22-1.88)	-
Vasoactive medication	3.33 (1.03-12.11)	1.81 (1.07-2.87)

Univariable and multivariable logistic regression for **anti-centromere positive** patients.

NVC pattern was entered as ordinal variable, in order: early, active or late. ACA=anti-centromere antibody, ATA= anti-topoisomerase antibody, CI=confidence interval, NR=non-Raynaud, NVC=nailfold videocapillaroscopy, OR= odds ratio, RP= Raynaud Phenomenon, SSc= systemic sclerosis.

Accepted Article

Table 4. ATA IgM levels are associated with more severe microangiopathy after adjustment for age and disease duration.

	Univariable OR (95% CI)	Multivariable OR (95% CI)
	NVC SSc pattern	NVC SSc pattern
Male	2.20 (0.64-7.53)	-
Age	1.06 (1.02-1.10)	1.05 (1.01-1.10)
Disease duration since RP	1.06 (1.01-1.11)	1.05 (1.0-1.11)
Disease duration since NP	1.21 (0.88-1.67)	-
Autoantibody specific IgM positive	1.29 (0.43-3.82)	-
Autoantibody specific IgA positive	1.00 (1.00-1.00)	-
Autoantibody specific IgG levels, U/ml	1.24 (0.41-3.77)	-
Autoantibody specific IgM levels, aU/ml	2.24 (0.97-5.18)	2.70 (1.06-4.22)
Autoantibody specific IgA levels, aU/ml	0.96 (0.42-2.18)	-
Autoantibody specific isotype expression	1.29 (0.43-3.82)	-
Immunosuppressive medication	1.65 (0.56-4.86)	-
Vasoactive medication	1.33 (0.30-5.96)	-

Univariable and multivariable logistic regression for **anti-topoisomerase positive** patients.

NVC pattern was entered as ordinal variable, in order: early, active or late. ACA=anti-centromere antibody, ATA= anti-topoisomerase antibody, CI=confidence interval, NR=non-Raynaud, NVC=nailfold videocapillaroscopy, OR= odds ratio, RP= Raynaud Phenomenon, SSc= systemic sclerosis.