

Longterm Effect on Leisure Time Physical Activity Level in Individuals with Axial Spondyloarthritis: Secondary Analysis of a Randomized Controlled Trial

Silje Halvorsen Sveaas, Hanne Dagfinrud, Melissa Woll Johansen, Elisabeth Pedersen, Ole-Martin Wold, and Annelie Bilberg

ABSTRACT. Objective. To explore the longterm effect of a 3-month exercise program on leisure time physical activity level in individuals with axial spondyloarthritis (axSpA).

Methods. A secondary analysis was performed on data from 100 individuals with axSpA who were included in a randomized controlled trial. The exercise group (EG) participated in a 3-month exercise program while the control group (CG) received no intervention. Physical activity during leisure time was measured with a questionnaire (physically active: ≥ 1 h/week with moderate/vigorous intensity physical activity). Disease activity was measured with the Ankylosing Spondylitis Disease Activity Scale (ASDAS; higher score = worst). Statistical analyses were performed on an intention-to-treat basis using chi-square tests, logistic regression, and mixed models.

Results. At the 12-month followup, significantly more individuals in the EG than in the CG were physically active [29 (67%) vs 13 (30%), $p < 0.001$] and exercised 2–3 times/week [25 (58%) vs 15 (34%), $p = 0.02$], and fewer exercised at light intensity [3 (8%) vs 14 (44%), $p = 0.002$]. “Participation in the EG” (OR 6.7, 95% CI 2.4–18.6, $p < 0.001$) and “being physically active at baseline” (OR 4.7, 95% CI 1.4–15.8, $p = 0.01$) were the factors most associated with being physically active. There were no differences between the groups in ASDAS ($p = 0.79$).

Conclusion. A 3-month exercise program had a beneficial longterm effect on leisure time physical activity in individuals with axSpA, thus indicating a more beneficial health profile. Still, few individuals continued the intensive program, and there was no difference between the groups in disease activity after 12 months. (ClinicalTrials.gov: NCT02356874) (J Rheumatol First Release June 15 2020; doi:10.3899/jrheum.190317)

Key Indexing Terms:

SPONDYLOARTHRITIS

EXERCISE

PHYSICAL ACTIVITY

From the National Advisory Unit on Rehabilitation in Rheumatology, Department of Rheumatology, Diakonhjemmet Hospital, Oslo; Department of Physiotherapy, Martina Hansens Hospital, Bærum; Department of Physiotherapy, University Hospital of North Norway, Tromsø; Department of Coaching and Psychology, Norwegian School of Sport Sciences, Oslo; The Norwegian Rheumatology Association, Norway; Institute of Neuroscience and Physiology, Section of Health and Rehabilitation, Physiotherapy, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden.

Supported by The Norwegian Fund for Post-Graduate Training in Physiotherapy and the Norwegian Rheumatology Association.

S.H. Sveaas, PhD, National Advisory Unit on Rehabilitation in Rheumatology, Diakonhjemmet Hospital; H. Dagfinrud, PhD, National Advisory Unit on Rehabilitation in Rheumatology, Diakonhjemmet Hospital; M.W. Johansen, MSc, Department of Physiotherapy, Martina Hansens Hospital; E. Pedersen, PT, the Department of Physiotherapy, University Hospital of North Norway; O.M. Wold, MSc, Department of Coaching and Psychology, Norwegian School of Sport Sciences, and The Norwegian Rheumatology Association; A. Bilberg, PhD, Institute of Neuroscience and Physiology, Section of Health and Rehabilitation, Physiotherapy, Sahlgrenska Academy, University of Gothenburg.

Address correspondence to S.H. Sveaas, National Advisory Unit on Rehabilitation in Rheumatology, Department of Rheumatology, Diakonhjemmet Hospital, PO Box 23 Vinderen, 0319 Oslo, Norway. E-mail: s.h.sveaas@medisin.uio.no

Accepted for publication November 4, 2019.

Axial spondyloarthritis (axSpA) is a chronic inflammatory rheumatic disease that mainly affects the axial skeleton¹. The disease is characterized by inflammatory back pain, and may also lead to reduced spinal mobility, arthritis, enthesitis, stiffness, and fatigue¹, as well as an increased risk of cardiovascular (CV) diseases².

Physical activity is recommended as an important part of the management of axSpA³, but there is concern that exercise at a vigorous intensity might exacerbate axSpA disease activity, and patients have generally been advised to engage in light-intensity exercises^{4,5}. However, in recently published recommendations, individuals with inflammatory arthritis are advised to adhere to the general physical activity recommendations⁶. These recommendations state that adults should perform moderate intensity aerobic activity for a minimum of 30 min on 5 days per week or vigorous intensity aerobic activity for a minimum of 20 min on 3 days per week (or a combination), and engage in strength exercises 2–3 days per week. Hence, these recommendations represent a shift toward a more active approach with cardiorespiratory and strength exercises.

Exercise reduces the disease burden in individuals with axSpA because it has beneficial effects on disease activity, physical function, stiffness^{7,8,9}, and CV health¹⁰. Despite this, individuals with axSpA tend to be less physically active than recommended^{11,12,13}, and they report that they engage less in activities of moderate and vigorous intensity^{14,15}. The most frequent exercise modes are pool exercises, stretching, and walking^{13,16,17}, and a previous cross-sectional study reported that fewer than one-third of individuals with axSpA engage in regular aerobic activities¹³. Hence, it is important to investigate how physical activity can be implemented into daily life for this group. Staying physically active over time is important to secure health benefits¹⁸, but few studies have investigated the longterm effect of exercise programs on physical activity level in axSpA. The aim of this study was therefore to examine longterm physical activity levels after a 3-month exercise program in individuals with axSpA.

MATERIALS AND METHODS

Design. This is a secondary analysis of a multicenter randomized controlled trial comparing the effects of 12 weeks of supervised exercise with usual care. The trial was conducted at outclinic rheumatology departments in Norway [Diakonhjemmet Hospital (DH), Martina Hansen Hospital (MHH), and the University Hospital of North Norway (UNN)] and in Sweden [Sahlgrenska University Hospital (SUH)]. The study was approved by the Regional Committee for Medical and Health Research Ethics (REK South East 2015/86) in Norway and the Regional Ethical Review Board Gothenburg in Sweden (032-16). All procedures followed the Declaration of Helsinki, and all participants gave written and oral informed consent before entering. The study protocol is registered at ClinicalTrials.gov (NCT02356874).

Participants. Participants were recruited from outpatient rheumatology departments as well as through various social media channels. The inclusion criteria were fulfillment of the Assessment of Spondyloarthritis international Society criteria for axSpA¹⁹, age 18–70 years, no change in tumor necrosis factor inhibitor use during the last 3 months, and moderate to high disease activity [Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) ≥ 3.5]. In addition, participants should not have participated in regular exercises (> 1 h per week) during the last 6 months with the aim of increasing cardiorespiratory fitness or muscular strength²⁰. Even though they could have been physically active in activities such as pool exercises, walking, and stretching, the program should have the potential to further increase their cardiorespiratory fitness and muscular strength. Exclusion criteria were established or symptoms of coronary heart disease, other comorbidity involving reduced exercise capacity, inability to participate in weekly exercise sessions, and pregnancy.

Exercise group (EG): exercise program. The EG had access to supervised sessions during a 3-month period. A physiotherapist with experience in the field of rheumatology and trained in the exercise protocol supervised the sessions twice a week. The program followed the American College of Sports Medicine (ACSM) recommendations for cardiorespiratory and strength exercises (Sveaas, *et al*, 2019 for details²¹). Some pain was tolerated during the exercises (≤ 5 on a scale 0–10), but the exercises were adapted if the pain got worse the day after.

The cardiorespiratory exercise was performed 3 times per week for ≥ 40 min. Two times per week the EG performed intervals on a treadmill or a cycle ergometer at vigorous intensity level [10 min warmup, thereafter 4 min at 90–95% of maximal heart rate (HR) followed by 3 min of active resting at 70% of maximal HR repeated 4 times]²². Maximal HR was determined at baseline and monitored by a pulse-watch. Once a week,

participants in the EG performed ≥ 40 min on a moderate intensity level ($> 70\%$ of maximal HR) on their own.

The strength exercises were performed twice a week with 8–10 repetitions maximum in 2–3 sets. The exercises were individually adapted and focused on major muscle groups (squat, leg press, deadlifts, rows to chest, bench press, shoulder press, pull-downs, and sit-ups).

Behavior change techniques. Several behavior change techniques²³ were used in the delivery of the intervention. The most important was the use of supervision that gave the participants a detailed plan, individual feedback, and information. Two of the hospitals organized the exercise program as group sessions (UNN and SUH), while the physiotherapists were available for the participants during a fixed timepoint at the 2 others (DH and MHH). Equal for all the participants in the EG was that they had committed to exercise, the intensity was monitored during the sessions, and adherence was recorded by the physiotherapist as well as self-reported in an exercise diary. Participants in both the EG and the control group (CG) tested their physical fitness before and after the intervention period.

After the 3-month period, the exercise intervention ended, and participants were not given any instructions or reminders regarding physical activity.

Program for CG. Participants in the CG received no intervention and were asked to continue their usual physical activity habits during the intervention period. All included participants received standard outpatient care from their respective hospitals, but before inclusion in the study it was specified that no change in medication before the 3-month assessment was desirable. After the 3-month assessment, participants in the CG were not given any instruction in physical activity.

Assessments. All participants underwent a clinical examination and filled out questionnaires at baseline and 3 months after inclusion. After 12 months, a questionnaire was sent by postal mail to all participants together with a prepaid envelope. Along with the questionnaire, participants were requested to visit the local study center to give blood samples to be analyzed for C-reactive protein (CRP) and erythrocyte sedimentation rate. Up to 2 reminders were given by phone calls or text messages.

Variables such as age, educational level, working status, disease characteristics, and medication were obtained from the questionnaires. Cardiorespiratory fitness was tested using a maximal walking test on treadmill for estimation of peak oxygen uptake ($\text{VO}_{2\text{peak}}$) at baseline. Spinal mobility was assessed by the Bath Ankylosing Spondylitis Metrology Index (0–10, 10 = worst)²⁴.

Physical activity level. To assess physical activity we used the questions from the Nord-Trøndelag Health Study (HUNT1)²⁵ and the International Physical Activity Questionnaire Short Form (IPAQ-SF)²⁶. The HUNT1 questions are considered valid to assess leisure time physical activity because higher values of the summary index correlate positively with higher values of $\text{VO}_{2\text{max}}$ and also with accelerometer data. Further, HUNT1 questions are reported to have high test-retest reliability with correlation coefficients from 0.76 to 0.87. Participants were asked, “how often do you exercise? (never, less than once a week, once a week, 2–3 times a week and almost every day),” with the text, “give an average, exercise means going for walks, skiing, swimming and training/sports”. If they exercised ≥ 1 week, they were asked about the intensity [no sweating/not out of breath (light), sweating/out of breath (moderate), or almost exhausted (vigorous)] and average duration (< 15 min, 16–30 min, > 30 min–1 h, or > 1 h). A product of frequency, intensity, and duration gave a summary index²⁵. To categorize individuals into physical activity levels (physically active or inactive), total minutes per week were calculated by multiplying frequency and duration (frequency: never = 0, less than once a week = 0, once a week = 1, 2–3 times per week = 2.5, almost every day = 7; duration: < 15 min = 0, 16–30 min = 23 min, > 30 min–1 h = 45 min, > 1 h = 60 min)²⁷. Thereafter, total minutes per week and intensity were used to categorize individuals as either physically inactive (0–420 min with light physical activity or 0–59 min with moderate/vigorous physical

activity per week) and physically active (defined as ≥ 60 min per week with moderate/vigorous physical activity)²⁷.

IPAQ-SF is reported to have acceptable criterion validity when compared with an activity tracker and to have good test-retest reliability with a correlation coefficient of 0.80²⁶. IPAQ-SF consists of 6 questions about physical activity during the last 7 days. Based on the data, metabolic equivalent (MET) scores were calculated, and 1 MET represents the body's resting energy expenditure (www.ipaq.ki.se). All the calculations and data cleaning were performed according to the official scoring protocol, but missing values were coded as no activity on the respective activity level. The MET scores were calculated by multiplying the number of minutes per day by the number of days per week by the activity MET (vigorous intensity = 8 MET, moderate intensity = 4 MET, and walking = 3.3 MET). Total physical activity level was calculated by summarizing the MET scores from the 3 activity levels.

Participants were also asked to list the exercise modes they had been active in during the last year. If they reported at the 12-month followup that they exercised ≥ 1 /week, the listed exercise modes were categorized into relevant categories for analytical purposes.

Disease activity and physical function. Disease activity was measured with the Ankylosing Spondylitis Disease Activity Score (ASDAS)²⁸ and the BASDAI²⁹. ASDAS is a composite score of CRP and self-reported variables: (1) neck/back/hip pain, (2) peripheral joint pain, (3) duration of morning stiffness, and (4) global assessment. All self-reported variables are reported on an 11-point numeric rating scale (NRS). ASDAS gives a continuous variable (inactive disease < 1.3, low disease activity 1.3 to < 2.1, high disease activity 2.1–3.5, and very high disease activity > 3.5). The BASDAI is a self-reported index of 5 symptoms (fatigue, neck-back-hip pain, peripheral joint pain, tenderness, and degree/length of morning stiffness)²⁹. Physical function was assessed with the Bath Ankylosing Spondylitis Functional Index (BASFI), which is a disease-specific index³⁰. Each question in BASDAI and BASFI was answered on an 11-point NRS, and a sum score from 0 to 10 (10 = worst) was calculated.

Statistical analyses. Data are presented as mean (SD), median (min–max), and number (%) as appropriate. All statistical analyses were performed on an intention-to-treat basis. For comparisons between groups, the chi-square test was used to analyze differences in categorical data and independent sample t test or Mann-Whitney U test was used to analyze differences in continuous variables as appropriate.

To examine factors associated with being physically active at the 12-month followup, physically active participants were compared with physically inactive participants according to background variables and variables thought to be associated with physical activity level. Thereafter, a multivariate logistic regression analysis was performed to calculate the odds for being physically active at the 12-month followup. Candidate variables for the multivariable logistic regression analysis were sex, age, and variables with p values < 0.1 in simple analyses.

A linear mixed model was used to assess differences between the groups in disease activity and physical function at 12-month followup, with adjustments for baseline values and study center and the interaction between treatment and time.

The linear mixed model analyses were performed in Stata (StataCorp.), and all other statistical analyses were performed in SPSS (IBM Corp.). The level of statistical significance was set at $p < 0.05$.

RESULTS

Participants. Flow of participants is shown in Figure 1. A total of 97 (97%) and 88 (88%) of 100 participants completed the assessment at 3 and 12 months, respectively. Background variables for the EG and CG are shown in Table 1.

Adherence and adverse events. A total of 38 (76%) participants in the EG followed $\geq 80\%$ of the prescribed exercise protocol (≥ 29 of 36 sessions registered by the

physiotherapist or in the exercise diaries), while 4 (8%) participants did not attend more than a few sessions. Two participants reported persistent pain during the exercise period but completed the prescribed exercise protocol. In addition, 1 participant experienced chest pain and nausea during the exercises and completed the intervention at moderate intensity after advice from a cardiologist.

Longterm effect on physical activity level. The EG had a significantly higher exercise summary index at the 12-month followup ($p = 0.01$; Table 2). Further, significantly more individuals in the EG were physically active (≥ 1 h per week with moderate/vigorous physical activity) compared to the CG ($p < 0.001$; Figure 2). At the 12-month followup, only 17 of 43 (40%) individuals in the EG performed both cardiorespiratory and strength exercises.

A total of 28 of 88 (32%) participants had missing items on the IPAQ. Although not significant, there was a tendency toward more MET at a vigorous intensity level and less walking in the EG compared to the CG. Further, more individuals in the CG were physically active on a moderate level than in the EG ($p = 0.02$).

Factors associated with being physically active at the 12-month followup. The adjusted logistic regression analysis (Table 3) showed that having received the exercise intervention ($p < 0.001$) and being physically active at baseline ($p = 0.01$) were the only factors that were significantly associated with being physically active at the 12-month followup. The OR for being physically active was 6.0 times higher in the EG than in the CG.

In the EG, physical activity level at 12-month followup was not associated with adherence to the exercise program (no. sessions) or study center.

Longterm effect on disease activity and physical function. The significant beneficial effect of the intervention seen at 3 months' followup (Sveaas, *et al*, 2019²¹), was no longer present at 12 months' followup; no statistical differences were seen between the groups in disease activity (ASDAS, $p = 0.79$; BASDAI, $p = 0.37$) or physical function (BASFI; $p = 0.821$; Figure 3).

DISCUSSION

The results indicate that participation in an exercise program increases the chance of staying physically active at a health-enhancing level over time in individuals with axSpA. At the 12-month followup, almost 70% of the individuals in the EG were physically active during leisure time (≥ 1 h per week with moderate/vigorous intensity activity) compared to 30% in the CG. Hence, individuals in the EG were 6 times more likely to be physically active at the 12-month followup than those who did not receive the intervention. Still, few individuals continued with the exercise program, and the beneficial effect on disease activity and physical function found immediately after the exercise program had declined at the 12-month followup.




Figure 1. Flow of participants throughout the randomized controlled trial.

Physical activity and exercise are recognized as important in the management of axSpA³. However, previous research has shown that individuals with axSpA are less physically active than recommended^{11,12,13,31}. The longterm improved physical activity level shown in our study is therefore important, because there are indisputable health effects of even a small enhancement in leisure time physical activity¹⁸.

To the best of our knowledge, this is the first study to

demonstrate a prolonged beneficial effect on physical activity level after an exercise program in individuals with axSpA. There are few comparable studies, and previous research is conflicting. Three studies, one in individuals with axSpA¹⁰, one in individuals with osteoarthritis³², and one in individuals with fibromyalgia³³ concluded with no effect of an exercise program on longterm activity, while a study of elderly individuals with rheumatoid arthritis (RA)³⁴

Table 1. Baseline characteristics of all participants, the exercise group, and the control group.

Characteristics	All, n = 100	Exercise Group, n = 50	Control Group, n = 50
Age, yrs, mean (min–max)	46.2 (23–69)	45.1 (23–68)	47.2 (24–69)
Sex, male, n (%)	47 (47)	25 (50)	22 (44)
Radiographic axSpA, n (%)	70 (70)	38 (76)	32 (64)
Married/cohabitant, n (%)	76 (76)	39 (78)	37 (74)
In work, n (%)	81 (81)	42 (78)	39 (78)
Current smoking, n (%)	12 (12)	5 (10)	7 (14)
Height, cm, mean (SD)	172 (11)	172 (11)	172 (11)
Weight, kg, mean (SD)	82.9 (17.9)	81.5 (19.4)	83.1 (19.5)
Medication			
NSAID, n (%)	71 (71)	38 (76)	33 (66)
TNFi, n (%)	44 (44)	23 (46)	21 (42)
Disease characteristics			
Disease activity (ASDAS-CRP), mean (SD)*	2.6 (0.7)	2.6 (0.8)	2.7 (0.6)
Disease activity (BASDAI), mean (SD)	5.1 (1.6)	4.9 (1.6)	5.3 (1.5)
CRP, mg/l, median (min–max)	2 (2–28)	2 (2–28)	2 (2–13)
ESR, mm/h, median (min–max)	8 (1–67)	8 (2–67)	8 (1–28)
Physical function (BASFI), median (min–max)	3.2 (0.2–9.1)	2.6 (0.2–6.7)	3.0 (0.4–9.1)
Spinal flexibility (BASMI), mean (SD)	2.8 (1.3)	2.9 (1.3)	2.6 (1.3)
Cardiorespiratory fitness (VO _{2peak}), mean (SD)	35.7 (36.3)	36.0 (5.9)	35.4 (6.9)

*ASDAS-CRP: C-reactive protein–based Ankylosing Spondylitis Disease Activity Score (inactive disease < 1.3, low disease activity 1.3 to < 2.1, high disease activity 2.1–3.5, and very high disease activity > 3.5). axSpA: axial spondyloarthritis; BASDAI: Bath Ankylosing Disease Activity Index; BASFI: Bath Ankylosing Spondylitis Functional Index; BASMI: Bath Ankylosing Spondylitis Metrology Index (All BAS instruments: 0–10, 10 = worst); ESR: erythrocyte sedimentation rate; NSAID: nonsteroidal antiinflammatory drugs; TNFi: tumor necrosis factor inhibitor; VO_{2peak}: peak oxygen uptake.

Table 2. Comparison of physical activity level at 12-month followup between the exercise and the control groups.

Activity Level	Exercise Group, n = 44	Control Group, n = 44	p
Exercise summary index* (0–15, 15 = high), median (min–max)	3.8 (0–15)	2.3 (0–15)	0.01
Exercise frequency			
Never (0)	3 (7)	2 (5)	0.63
< 1 time per week (0.5)	2 (5)	10 (23)	0.02
1 time per week (1)	7 (16)	8 (18)	0.81
2–3 times per week (2.5)	25 (58)	15 (34)	0.02
Almost every day (5)	6 (14)	9 (21)	0.42
Exercise intensity†	n = 36	n = 31	
Not out of breath or sweating (1)	2 (6)	13 (42)	0.001
Out of breath and sweating (2)	25 (69)	15 (48)	
Almost exhausted (3)	9 (25)	3 (10)	
Exercise duration†	n = 38	n = 31	
< 15 min (0.10)	0 (0)	0 (0)	
16–30 min (0.38)	1 (3)	6 (19)	0.07
30–60 min (0.75)	27 (71)	19 (61)	
> 1 h (1.0)	10 (26)	6 (19)	
Exercise mode	n = 43	n = 44	
Cardiorespiratory	26 (60)	6 (14)	< 0.001
Muscular strength	21 (49)	5 (11)	< 0.001
Cardiorespiratory and muscular strength	17 (39)	3 (7)	< 0.001
Walking	18 (42)	23 (52)	0.30
Pool exercises	8 (19)	14 (32)	0.14
Physical activity level (MET)	n = 44	n = 44	
Total MET, median (min–max)	1886 (0–17,892)	1386 (0–9600)	0.83
MET vigorous	720 (0–10,080)	0 (0–7200)	0.15
MET moderate	120 (0–5040)	160 (0–5040)	0.95
MET walking	363 (0–4158)	495 (0–4158)	0.21

Values are n (%) unless otherwise specified. * Calculated based on exercise frequency, duration, and intensity with the scores for each response given in the parentheses. † Only participants exercising ≥ 1 per week were asked about intensity/duration of exercise. MET: metabolic equivalent.




Figure 2. The longterm effect of the supervised exercise program on physical activity level. Bar graphs represent the frequency of physically inactive and physically active participants at baseline, 3-month, and 12-month followup in the exercise group (n = 43) and the control group (n = 44). Physically active: ≥ 60 min with moderate to vigorous activity. Physically inactive: 0–420 min per week with light activity or 0–59 min with moderate to vigorous activity.

Table 3. Factors associated with being physically active at 12-month followup. A total of 42 of 87 (48%) participants were physically active.

Variables	Crude Estimates, OR (95% CI)	p	Adjusted Estimates ³ , OR (95% CI)	p
Age				
Continuous	1.00 (0.96–1.04)	0.88	0.99 (0.95–1.04)	0.79
Sex				
Male	Ref		Ref	
Women	1.38 (0.59–3.20)	0.46	1.51 (0.57–4.02)	0.41
Physical activity at baseline*				
Inactive at baseline	Ref		Ref	
Physically active at baseline	2.57 (0.95–6.92)	0.062	4.73 (1.42–15.75)	0.01
Intervention				
Control group	Ref		Ref	
Exercise group	4.94 (1.99–12.26)	0.001	6.72 (2.42–18.63)	< 0.001

*Physically active was defined as ≥ 1 h per week with moderate/vigorous physical activity at baseline, and physically inactive was defined as 0–420 min with light activity and < 60 min with moderate/vigorous activity.


found a positive longterm effect on physical activity. The conflicting results might be explained by several reasons. First, the timepoint for measurement differed between the studies; one of the studies that found no effect on longterm activity¹⁰ measured physical activity right after completion of the intervention. However, it has been reported that exercise adherence seems to diminish over time³⁵, hence it is not likely that the physical activity level had been higher at a later timepoint.

Second, the difference in results might be explained by differences in exercise modes and intensities. It is well known that intensity is crucial for achieving the effect of exercise¹⁸ and that perceived benefits of exercise are


important for adherence³⁶. The study in RA³⁴ and our study³⁷ included cardiorespiratory interval exercises (with an intensity of 70–89% and 90–95% of maximal HR, respectively), and both studies demonstrated large effects on general health immediately after the exercise intervention. The experience of immediate benefits may have contributed to the prolonged effect on physical activity level observed in both the RA study³⁴ and the present study.

Motivational and educational interventions have shown beneficial effects on physical activity level in individuals with rheumatic diseases^{38,39,40}. Even if these interventions are different from our exercise program, they are similar regarding inclusion of well-known facilitators for physical

Disease activity (ASDAS-CRP)



Disease activity (BASDAI)



Physical function (BASFI)




Figure 3. The longterm effect of the 3-month supervised exercise program on disease activity and physical function. ASDAS-CRP: C-reactive protein-based Ankylosing Spondylitis Disease Activity Score (inactive disease < 1.3, low disease activity 1.3 to < 2.1, high disease activity 2.1–3.5, and very high disease activity > 3.5); BASDAI: Bath Ankylosing Spondylitis Disease Activity Index (0–10, 10 = worst); BASFI: Bath Ankylosing Spondylitis Functional Index (0–10, 10 = worst).

activity, such as health professionals providing information^{41,42}, a focus on motivation for exercise^{36,43,44}, and use of reminders⁴⁵. These factors may be of importance for improving physical activity level in individuals with axSpA.

In line with this, a recent systematic review concluded that education and supervision are important factors for adherence to exercise in individuals with axSpA³⁵. Obviously, motivational or educational interventions are more cost-effective than supervised exercise programs, but it should be noticed that the magnitude of the clinical effect was poor in one of the educational studies⁴⁰. Brophy, *et al*⁴⁴ stated that adding motivational strategies to the delivery of exercise programs was most effective in increasing physical activity, hence a combination of motivational and practical programs is probably important to increase physical activity level in patients with axSpA.

We found that physical activity status at baseline and participation in the supervised exercise program were the factors that were the most strongly associated with physical activity status at 12 months' followup. Previous research has shown that lower disease activity^{31,44,46}, better physical function, and quality of life is associated with being physically active in people with axSpA³¹. However, none of these factors were associated with physical activity level at the 12-month followup in our present study, and the difference in results may be explained by unequal study designs^{31,44,46} and larger study populations in previous studies^{31,44}. Our result is in line with the result of a systematic review stating that among several factors, low levels of physical activity at baseline were associated with poor adherence to physiotherapy treatment⁴⁷. Exercise experience is known to be essential for achieving exercise self-efficacy⁴⁵. In our study,

the participants in the EG got practical training in exercise, which may perhaps compensate for lack of previous exercise experience.

The diminished effects on disease activity and physical function at 12 months' followup are in accordance with findings in previous studies^{32,33,34,48}. Despite the beneficial effect on physical activity level, few participants continued with the exercise program. Our result is in line with earlier research reporting that it is more difficult for people to adhere to exercises at vigorous intensity, and that exercise-induced adaptations are reversed over time without adherence to the program¹⁸. Hence, the clinical implication is that vigorous-intensity exercise should be recommended as important in the treatment of individuals with axSpA owing to the beneficial effects on disease activity. However, as the effect slowly declines, future studies should investigate whether intermittent booster exercise sessions may increase participation in vigorous exercises.

Strengths of our present study are the study design, the relatively large sample size, the low dropout rate, and the longterm followup period. Further, the exercise program was based on the ACSM recommendations. The generalization of the study is probably high, because the study was carried out in outpatient clinics at 4 different hospitals and several physiotherapists supervised the exercise program. We have previously shown that the exercise program was safe and well tolerated in a group of individuals with high disease activity³⁷. Self-reporting of physical activity level may be considered as a limitation, because participants may have under- or overreported their physical activity⁴⁹. Further, participants were not blinded for group assignment, a factor that is reported to exaggerate subjective outcomes⁵⁰. Nevertheless, it is a strength that physical activity was measured with standardized, frequently used questionnaires²⁷. And, as the numbers of inactive individuals in the CG were unchanged during the study period, we argue that this strengthens the validity of the physical activity questions. Likewise, a limitation is how the exercise mode questions were formulated. Although only responses from individuals exercising ≥ 1 /week were included, individuals in the EG might have included activities from the intervention period.

A supervised exercise program seems to increase the chance of staying physically active over time, thereby maintaining a beneficial health profile. Still, few individuals continued the exercise program, and the beneficial effect on disease activity and physical function at the end of the exercise program had declined at 12 months' followup.

ACKNOWLEDGMENT

We thank the patient representatives for their contribution in the study, as well as all the participants. Further, we also thank the physiotherapists Emma Klittmar, Linn A. Haukland, Mary E. Deighan Hanssen, and Kim Reier Nielsen Martinsen for their help with exercise supervision, and The Norwegian Fund for Post-Graduate Training in Physiotherapy for financial support.

REFERENCES

1. Sieper J, Poddubnyy D. Axial spondyloarthritis. *Lancet* 2017;390:73-84.
2. Mathieu S, Soubrier M. Cardiovascular events in ankylosing spondylitis: a 2018 meta-analysis. *Ann Rheum Dis* 2018;78:e57.
3. van der Heijde D, Ramiro S, Landewe R, Baraliakos X, Van den Bosch F, Sepriano A, et al. 2016 update of the ASAS-EULAR management recommendations for axial spondyloarthritis. *Ann Rheum Dis* 2017;76:978-91.
4. Iversen MD, Scanlon L, Frits M, Shadick NA, Sharby N. Perceptions of physical activity engagement among adults with rheumatoid arthritis and rheumatologists. *Int J Clin Rheumatol* 2015;10:67-77.
5. Lundberg IE, Nader GA. Molecular effects of exercise in patients with inflammatory rheumatic disease. *Nat Clin Pract Rheumatol* 2008;4:597-604.
6. Rausch Osthoff AK, Niedermann K, Braun J, Adams J, Brodin N, Dagfinrud H, et al. 2018 EULAR recommendations for physical activity in people with inflammatory arthritis and osteoarthritis. *Ann Rheum Dis* 2018;77:1251-60.
7. Pecourneau V, Degboe Y, Barnette T, Cantagrel A, Constantin A, Ruysen-Witrand A. Effectiveness of exercise programs in ankylosing spondylitis: a meta-analysis of randomized controlled trials. *Arch Phys Med Rehabil* 2018;99:383-9.
8. O'Dwyer T, O'Shea F, Wilson F. Exercise therapy for spondyloarthritis: a systematic review. *Rheumatol Int* 2014;34:887-902.
9. van den Berg R, Baraliakos X, Braun J, van der Heijde D. First update of the current evidence for the management of ankylosing spondylitis with non-pharmacological treatment and non-biologic drugs: a systematic literature review for the ASAS/EULAR management recommendations in ankylosing spondylitis. *Rheumatology* 2012;51:1388-96.
10. Niedermann K, Sidelnikov E, Muggli C, Dagfinrud H, Hermann M, Tamborini G, et al. Effect of cardiovascular training on fitness and perceived disease activity in people with ankylosing spondylitis. *Arthritis Care Res* 2013;65:1844-52.
11. O'Dwyer T, O'Shea F, Wilson F. Physical activity in spondyloarthritis: a systematic review. *Rheumatol Int* 2015;35:393-404.
12. Jacquemin C, Servy H, Molto A, Sellam J, Foltz V, Gandjbakhch F, et al. Physical activity assessment using an activity tracker in patients with rheumatoid arthritis and axial spondyloarthritis: Pprospective observational study. *JMIR Mhealth Uhealth* 2018;6:e1.
13. Fabre S, Molto A, Dadoun S, Rein C, Hudry C, Kreis S, et al. Physical activity in patients with axial spondyloarthritis: a cross-sectional study of 203 patients. *Rheumatol Int* 2016;36:1711-8.
14. Swinnen TW, Scheers T, Lefevre J, Dankaerts W, Westhovens R, de Vlam K. Physical activity assessment in patients with axial spondyloarthritis compared to healthy controls: a technology-based approach. *PLoS One* 2014;9:e85309.
15. O'Dwyer T, O'Shea F, Wilson F. Decreased physical activity and cardiorespiratory fitness in adults with ankylosing spondylitis: a cross-sectional controlled study. *Rheumatol Int* 2015;35:1863-72.
16. Passalent LA, Soever LJ, O'Shea FD, Inman RD. Exercise in ankylosing spondylitis: discrepancies between recommendations and reality. *J Rheumatol* 2010;37:835-41.
17. Sundström B, Ekergard H, Sundelin G. Exercise habits among patients with ankylosing spondylitis. A questionnaire based survey in the county of Vasterbotten, Sweden. *Scand J Rheumatol* 2002;31:163-7.
18. Garber CE, Blissmer B, Deschenes MR, Franklin BA, LaMonte MJ, Lee IM, et al. American College of Sports Medicine position stand.

- Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc* 2011;43:1334-59.
19. Rudwaleit M, van der Heijde D, Landewe R, Listing J, Akkoc N, Brandt J, et al. The development of Assessment of Spondyloarthritis international Society classification criteria for axial spondyloarthritis (part II): validation and final selection. *Ann Rheum Dis* 2009;68:777-83.
 20. Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness: Definitions and distinctions for health-related research. *Public Health Rep* 1985;100:126-31.
 21. Sveaas SH, Bilberg A, Berg IJ, Provan SA, Rollefstad S, Semb AG, et al. High intensity exercise for 3 months reduces disease activity in axial spondyloarthritis (axSpA): A multicentre randomised trial of 100 patients. *Br J Sports Med* 2020;54:292-7.
 22. Helgerud J, Høydal K, Wang E, Karlsen T, Berg P, Bjerkaas M, et al. Aerobic high-intensity intervals improve vo2max more than moderate training. *Med Sci Sports Exerc* 2007;39:665-71.
 23. Michie S, Richardson M, Johnston M, Abraham C, Francis J, Hardeman W, et al. The behavior change technique taxonomy (v1) of 93 hierarchically clustered techniques: building an international consensus for the reporting of behavior change interventions. *Ann Behav Med* 2013;46:81-95.
 24. van der Heijde D, Landewe R, Feldtkeller E. Proposal of a linear definition of the Bath Ankylosing Spondylitis Metrology Index (BASMI) and comparison with the 2-step and 10-step definitions. *Ann Rheum Dis* 2008;67:489-93.
 25. Kurtze N, Rangul V, Hustvedt BE, Flanders WD. Reliability and validity of self-reported physical activity in the Nord-Trøndelag Health Study: HUNT 1. *Scand J Public Health* 2008;36:52-61.
 26. Craig CL, Marshall AL, Sjoström M, Bauman AE, Booth ML, Ainsworth BE, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 2003;35:1381-95.
 27. Kurtze N, Gundersen KT, Holmen J. Self-reported physical activity in Norwegian studies - a methodological problem. [Article in Norwegian.] *Nor Epidemiol* 2003;13:163-70.
 28. van der Heijde D, Lie E, Kvien TK, Sieper J, Van den Bosch F, Listing J, et al; Assessment of SpondyloArthritis international Society (ASAS). ASDAS, a highly discriminatory ASAS-endorsed disease activity score in patients with ankylosing spondylitis. *Ann Rheum Dis* 2009;68:1811-8.
 29. Garrett S, Jenkinson T, Kennedy LG, Whitelock H, Gaisford P, Calin A. A new approach to defining disease status in ankylosing spondylitis: The Bath Ankylosing Spondylitis Disease Activity Index. *J Rheumatol* 1994;21:2286-91.
 30. Calin A, Garrett S, Whitelock H, Kennedy LG, O'Hea J, Mallorie P, et al. A new approach to defining functional ability in ankylosing spondylitis: The development of the Bath Ankylosing Spondylitis Functional Index. *J Rheumatol* 1994;21:2281-5.
 31. Haglund E, Bergman S, Petersson IF, Jacobsson LT, Strombeck B, Bremander A. Differences in physical activity patterns in patients with spondylarthritis. *Arthritis Care Res* 2012;64:1886-94.
 32. Svege I, Nordsletten L, Fernandes L, Risberg MA. Exercise therapy may postpone total hip replacement surgery in patients with hip osteoarthritis: a long-term follow-up of a randomised trial. *Ann Rheum Dis* 2015;74:164-9.
 33. Larsson A, Palstam A, Lofgren M, Ernberg M, Bjersing J, Bileviciute-Ljungar I, et al. Resistance exercise improves muscle strength, health status and pain intensity in fibromyalgia--a randomized controlled trial. *Arthritis Res Ther* 2015;17:161.
 34. Lange E, Kucharski D, Svedlund S, Svensson K, Bertholds G, Gjørtsson I, et al. Effects of aerobic and resistance exercise in older adults with rheumatoid arthritis: a randomized controlled trial. *Arthritis Care Res* 2019 Jan;71:61-70.
 35. McDonald MT, Siebert S, Coulter EH, McDonald DA, Paul L. Level of adherence to prescribed exercise in spondyloarthritis and factors affecting this adherence: a systematic review. *Rheumatol Int* 2019;39:187-201.
 36. Ehrlich-Jones L, Lee J, Semanik P, Cox C, Dunlop D, Chang RW. Relationship between beliefs, motivation, and worries about physical activity and physical activity participation in persons with rheumatoid arthritis. *Arthritis Care Res* 2011;63:1700-5.
 37. Sveaas SH, Bilberg A, Berg IJ, Provan S, Rollefstad S, Semb AG, et al. High intensity exercise for 3 months reduces disease activity in axial spondyloarthritis (axSpA): a multicenter randomized trial of 100 patients. *Br J Sports Med* 2020;54:292-7.
 38. O'Dwyer T, Monaghan A, Moran J, O'Shea F, Wilson F. Behaviour change intervention increases physical activity, spinal mobility and quality of life in adults with ankylosing spondylitis: a randomised trial. *J Physiother* 2017;63:30-9.
 39. Thomsen T, Aadahl M, Beyer N, Hetland ML, Loppenthin K, Midtgaard J, et al. The efficacy of motivational counselling and SMS-reminders on daily sitting time in patients with rheumatoid arthritis: a nationwide randomized controlled trial. *Ann Rheum Dis* 2017;76:1603-6.
 40. Rodriguez-Lozano C, Juanola X, Cruz-Martinez J, Pena-Arrebola A, Mulero J, Gratacos J, et al; Spondyloarthropathies Study Group of the Spanish Society of Rheumatology. Outcome of an education and home-based exercise programme for patients with ankylosing spondylitis: a nationwide randomized study. *Clin Exp Rheumatol* 2013;31:739-48.
 41. Madsen M, Jensen KV, Esbensen BA. Men's experiences of living with ankylosing spondylitis: a qualitative study. *Musculoskeletal Care* 2015;13:31-41.
 42. Manning VL, Hurley MV, Scott DL, Bearne LM. Are patients meeting the updated physical activity guidelines? Physical activity participation, recommendation, and preferences among inner-city adults with rheumatic diseases. *J Clin Rheumatol* 2012;18:399-404.
 43. Niedermann K, Nast I, Ciurea A, Vliet Vlieland T, van Bodegom-Vos L. Barriers and facilitators of vigorous cardiorespiratory training in axial spondyloarthritis: surveys among patients, physiotherapists, rheumatologists. *Arthritis Care Res* 2019;71:839-51.
 44. Brophy S, Cooksey R, Davies H, Dennis MS, Zhou SM, Siebert S. The effect of physical activity and motivation on function in ankylosing spondylitis: a cohort study. *Semin Arthritis Rheum* 2013;42:619-26.
 45. Bandura A. Self-efficacy. The exercise of control. 11th ed. New York: W H Freeman/Times Books/Henry Holt & Co.;1997.
 46. Plasqui G, Boonen A, Geusens P, Kroot EJ, Starmans M, van der Linden S. Physical activity and body composition in patients with ankylosing spondylitis. *Arthritis Care Res* 2012;64:101-7.
 47. Jack K, McLean SM, Moffett JK, Gardiner E. Barriers to treatment adherence in physiotherapy outpatient clinics: a systematic review. *Man Ther* 2010;15:220-8.
 48. van Baar ME, Dekker J, Oostendorp RA, Bijl D, Voorn TB, Bijlsma JW. Effectiveness of exercise in patients with osteoarthritis of hip or knee: nine months' follow up. *Ann Rheum Dis* 2001;60:1123-30.
 49. Prince SA, Adamo KB, Hamel ME, Hardt J, Connor Gorber S, Tremblay M. A comparison of direct versus self-report measures for assessing physical activity in adults: a systematic review. *Int J Behav Nutr Phys Act* 2008;5:56.
 50. Wood L, Egger M, Gluud LL, Schulz KF, Juni P, Altman DG, et al. Empirical evidence of bias in treatment effect estimates in controlled trials with different interventions and outcomes: meta-epidemiological study. *BMJ* 2008;336:601-5.