

Letter

Famous Artists Who Suffer(ed) From Rheumatic Diseases: A Systematic Review

To the Editor:

Rheumatic diseases (RD) occur at a relatively high frequency in the population. We hypothesize that some of these diseases may have affected some artists and possibly influenced their works. In this article, a systematic review of all studies that described the occurrence of rheumatic diseases in famous artists in the world was performed. A PEO format (P = population, E = exposure, O = outcome) to elaborate the research question, "Famous artists (P), with rheumatic diseases (E), have their works changed (O) due to these diseases?" was used.

An extensive literature search in Pubmed/MEDLINE, Scielo, and LILACS, following the Preferred Reporting Items for Systematic reviews and Meta-Analyses guidelines, was performed without language restriction, from 1965 to June 2020. After the review of titles and abstracts, 116 out of 1026 articles were selected for reading the full texts, of which 68 were selected for this review. We have identified 20 famous artists who had RD. Table 1 is a summary of all data regarding the artists $^{1-20}$. Most of them had rheumatoid arthritis (RA) as a confirmed or presumptive diagnosis (n = 8), followed by gout (n = 3), systemic sclerosis (SSc; n = 2), reactive arthritis (n = 2), juvenile idiopathic arthritis (n = 1), ankylosing spondylitis (AS; n = 1), fibromyalgia (FM; n = 1), pycnodysostosis (n = 1), pustulotic arthro-osteitis syndrome (n = 1), and Susac syndrome (n = 1). Of note, some artists had 2 or more potential diagnoses: Lisboa ("Aleijadinho") with syphilis, leprosy, RA, or porphyria cutanea tarda; Rubens with gout; and Joyce with AS.

A confirmed diagnosis was verified in 8 of 20 and a presumptive diagnosis in 11 of 20. All of them continued to perform their artistic activities despite the limitations secondary to RD. Importantly, most (n = 15/20)of these artists had an art style change after the beginning of the RD. For only one (1/20) artist (Michelangelo), the RD seemed to not interfere with his artistic abilities, and for 4/20 (Aleijadinho, Barrie Cook, Benvenuto Cellini, and Erasmus of Rotterdam), we do not have information regarding this aspect. Further, artists with RA had polyarthritis (n = 5/8) and some of them had extraarticular manifestations (n = 3/8). Frida Kahlo had a medical history that was tragic and complex, including scoliosis, poliomyelitis, spina bifida, and multiple fractures after a car accident (needing more than 30 surgeries). These diagnostic hypotheses of postpoliomyelitis and posttraumatic FM also need to be considered²¹. One artist (Michelangelo) had probable saturnine gout secondary to lead paint exposure. Reactive arthritis was seen in 2 artists; they had suspected sexual contact as well as ocular inflammation followed by arthritis. SSc was a presumptive diagnosis for Aleijadinho, who had phalanx amputations. Treatments varied from nonpharmacological therapies such as diets (n = 3), spas (n = 4), exercises/ physical therapy (n = 4), surgeries (n = 2), occupational therapy (n = 2), synoviorthesis with yttrium (n = 1) and use of antiinflammatory drugs (n = 5), cortisone (n = 2), methotrexate (n = 1), antimalarials (n = 1), gold (n = 2), colchicine (n = 1), homeopathy (n = 2), and biologicals (n = 1).

Of note, RD in other artists were reported by Appelboom¹⁸, but articles were not found about them in our literature review. These artists are Marquise de Sevigné (writer, probable RA), Samuel Johnson (writer, gout), Mozart (composer, rheumatic fever), Niccolò Paganini (composer and violinist, Marfan or Ehlers-Danlos), Paul Scarron (poet, AS), Paul Verlaine (poet, knee arthritis), and Flannery O'Connor (writer, systemic lupus erythematosus).

A possible explanation for RA being the most frequent RD in these patients is that the occupational exposure to the color inks as well as

heavy smoking may have had a role in the development of RA in Renoir, Jawlensky, Niki, and Dufy. In addition, vapor from smoking contains several heavy metals such as cadmium. Metals are used in ink composition, such as in chrome yellow, molybdenum orange, and cadmium red; iron black and copper-zinc alloy powder (gold bronze) are used in novel silver and gold inks

It has been reported that mercury, as well as gold, induces autoimmunity in genetically susceptible animals. Delayed hypersensitivity to metals was already evaluated in patients with RA, and significant amounts of mercury, titanium, and palladium were observed in these patients.

Rubens, Renoir, Dufy, and Klee used significantly brighter and clearer colors based on toxic heavy metals, and fewer earth colors containing harmless iron and carbon compounds. There is evidence of occupational exposure to cadmium by Barrie Cook. Another relevant risk factor for the development of RD is vitamin D deficiency. For those people who work indoors, it is a concern, as many artistic activities are developed in an indoor environment.

When we discovered that the majority of the artists that had RD changed their style due to their rheumatic conditions, we were able to conclude that there is a close relationship between disease and art²². In the same way that the pain and functional limitation caused by RD evoked sadness in these artists, that pain also provoked them to create beauty in their artistic works. Therefore, these diseases may be considered a "therapeutic method"²². We must remember that the arts are a way of expressing deep feelings, including suffering.

Jozélio Freire de Carvalho¹, Associate Professor Felipe Freire da Silva², Rheumatology Resident Carlos Augusto Ferreira de Andrade^{3,4}, Rheumatologist José Dirson Argolo⁵, Associate Professor Licia Maria Henrique da Mota⁶, Associate Professor ¹Institute for Health Sciences from Federal University of Bahia, Salvador, Bahia;

²Rheumatology Division, Hospital das Clínicas, Faculty of Medicine, University of São Paulo, São Paulo-SP;

³Department of Epidemiology and Quantitative Methods in Health, Sergio Arouca National School of Public Health, Oswaldo Cruz Foundation, Rio de Janeiro;

⁴Faculty of Medicine, Vassouras University, Rio de Janeiro; ⁵School of Fine Arts of Federal University of Bahia and Art Studio Argolo,

⁶Service of Rheumatology, University Hospital of Brasilia, Posgraduate Program in Medical Sciences, Faculty of Medicine, University of Brasilia, Brasilia-DF, Brazil.

Address correspondence to Dr. J.F. de Carvalho, Rua das Violetas, 42, ap. 502, Pituba, Salvador, Bahia, Brazil. Email: jotafc@gmail.com.

REFERENCES

- Mota LMH, Neubarth F, Diniz LR, Carvalho JF, Santos Neto LL. Pierre-Auguste Renoir (1841-1919) and rheumatoid arthritis. J Med Biogr 2012;20:91-2.
- Appelboom T, De Boelpaepe C, Famaey JP, Ehrlich GE, Famaey JP. Rubens and the question of antiquity of rheumatoid arthritis. JAMA 1981;245:483-6.
- Zeidler H. ["Memories of my sick hands": life and medical history of the painter Alexej von Jawlensky.] [Article in German] Z Rheumatol 2011;70:336-57.
- Castillo-Ojugas A. [The effect of rheumatoid arthritis in the life and work of 3 great painters: Auguste Renoir, Raoul Dufy and Alexis Jawlensky.] [Article in Spanish] An R Acad Nac Med 1992; 109:673-83.

1

Table 1. Summary of famous artists and their rheumatic diseases, clinical features, treatments used, and art style changes.

Artist (Birth–Death), Specialization, Country of Birth	Age at Disease Onset, yrs	Confirmed (CD) or Presumptive Diagnosis (PD)	Potential Differential Diagnosis (PDD)	Clinical Features	Treatment	Continued to Work After the Disease	Disease Changed the Artist's Style	Reference, First Author, Yr
Pierre-Auguste Renoir (1841–1919), painter and sculptor, France	25	RA (CD)	No PDD	Severe polyarticular synovitis, deformity on hands, feet, and coxofemoral joint Fixed flexion of knees, destruction and ankylosis of his right shoulder, and ruptures in several extensor tendons of the hands Extraarticular: rheumatoid nodules, pleural effusion, facial paralysis, rheumatoid cachexia, necrosis of the distal phalanges of the fifth finger (vasculitis)	· Thermal baths · Exercises · Devices of occupational therapy (orthesis) · Purgatives · Phenazone	Yes	· Yes · Small and rapid strokes · Invented the moving canvas or picture roll · More interest in the human body instead of landscapes · More vivid colors	Mota, 2012 ¹
Peter Paul Rubens (1577–1640), painter, Germany	33	RA (PD)	Gout	· Chronic involvement of feet, knees, and hands (symmetrical), complicated by flares · Extraarticular: pleuritis	N/A	Yes	· Yes · Represented joint alterations in his paintings	Appelboom, 1981 ²
Alexej von Jawlensky (1864–1941), painter, Russia	65	RA (CD)	No PDD	Feet, knee, ankles, jaw, shoulders, elbows, and cervical spine evolving to hand deformities Complete immobilization due to cervical compression Extraarticular: pleuritis, anemia and dry mouth (possible secondary Sjögren syndrome)	Diet (vegetarian, raw food, no milk. Physical therapy. Spas. Painkillers (pyramidone, phenacetin plus codeine). Gold injections. Irradiations (deep radiograph, radiu water drinks, radiu compresses, and in. Injections of an undrug in and arounded the space of the sp	m im ijections) nknown d joints cine oal y, nom ie injections) as to treat and		Zeidler, 2011 ³
Raoul Dufy (1877–1953), painter, France	55	RA (CD)	No PDD	· Polyarthritis · ESR 33 mm/h	· ACTH · Cortisone · Gold · Aspirin · Physiotherapy	Yes	· Yes · Reduced size of paintings · Plots were less accurate and subjects were restricted to his imagination or what he could contemp without moving	Castillo-Ojugas 1992 ⁴ blate

Artist (Birth–Death), Specialization, Country of Birth	Age at Disease Onset, yrs	Confirmed (CD) or Presumptive Diagnosis (PD)	Potential Differential Diagnosis (PDD)	Clinical Features	Treatment	Continued to Work After the Disease	Disease Changed the Artist's Style	Reference, First Author, Yr
							Represented joint alterations in his late paintings Entitled one of his most colorful canvases "La Cortisone"	
Paul Klee (1879–1940), painter and poet, Switzerland	55	SSc after measles or heavy metal toxicity (CD)	N₀ PDD	Raynaud phenomenon Skin alterations with contractures and ulcerations Fatigue, exhaustion, dysphagia, weight loss, dyspnea Arthritic pain, disability, difficulty in holding paintbrushes Heart failure	· Medical recommendation of stopping smoki and visit a spa to improve his well- Medical records were destroyed by a fire at the hospital after his death · There is no indication either of how his illness was managed		· Yes · Simplicity, more intensity, and use of rough materials (e.g., burlap and newspaper) · More somber and contracted forms, drawing on black and brown · Broad brush strokes, heavy, black crayonlike lines, and dull colors · Paintings depicted agonies that he suffered suffering, death, and value of the suffering of the	ed fear,
Antoni Gaudí i Cornet (1852–1926), architect, Spain	66	Juvenile RA (PD)	Rheumatic fever	Arthralgia, mainly involving the ankles Difficulty walking during disease flare	 Vegetarian diet, plenty of liquid Homeopathic therapy Spa 	Yes	· Yes · Influenced his observational power and nature analysis	da Mota, 2012
Manolo Hugué (1872–1945), painter and sculptor, Spain	55	RA (CD)		Polyarthritis Deformity of fingers	· Diathermia · Physical therapy	Yes	· Yes · Partially abandoned sculpture and devoted himself to painting and poetry	Pou, 2011 ⁷
Antônio Francisco Lisboa "Aleijadinho" (1730–1814), sculptor and architect, Brazil	47	SSc (PD)	· Syphilis · Leprosy · RA · Porphyria cutanea tarda	Progressive disabling disease with hand deformity (phalanx amputations) Paraplegy Paraparesis	N/A	Yes	N/A	Azevedo, 2008
Max Slevogt (1868–1932), painter, Germany	27	Gout (CD)	N₀ PDD	Arthritis (feet and knees)	Bed rest Colchicine, iodite, aminofena potassium ointme Vegetarian diet, dietary measures Multiple spa trea	ent other	· Yes · Many of his attacks of gouty arthritis are illustrated in drawings that document involveme of feet and knees	
Michelangelo di Lodovico Buonarroti Simoni (1475–1564), painter, sculptor, poet, and architect, Italy	80	Gout (PD)	· Osteoarthritis · Saturnine gout · Deficiency of hypoxanthine- guanine phosphoribosyl transferase	· Arthritis, tophi · Urinary stones (uric acid) · Changes in the first carpometacarpal joint · Difficulty writing · Depression (or bipolar manic-depressive illness)	N/A	Yes	Probably not His articular illness seems to have had little effect on his artistic productivity Some of his paintings mirror his depression as seen in "Jeremiah in the Sistine Chapel"	Pinals, 2015 ¹⁰

Letter 3

Table 1. Continued.

Artist (Birth–Death), Specialization, Country of Birth	Age at Disease Onset, yrs	Confirmed (CD) or Presumptive Diagnosis (PD)	Potential Differential Diagnosis (PDD)	Clinical Features	Treatment	Continued to Work After the Disease	Disease Changed the Artist's Style	Reference, First Author, Yr
James Augustine Aloysius Joyce (1882–1941), poet and writer, Ireland	23	ReA or AS (PD)	No PDD	Recurrent iritis and oligo/polyarthritis Frequented prostitutes several times	Dental treatment Iristomy and iridectomy 8 ocular surgeries Poor adhesion to treatments	Yes	· Yes · Written works abound with references to doctors, symptoms, and diseases of all kinds	Ventura, 2008 ¹¹
Barrie Cook (1929–), painter, UK	44	RA after chronic cadmium exposure from pain and smoking (CD)		· RF 149 IU/mL · anti-CCP > 500 U/mL	Multiple biologicals	Yes	N/A	Cates, 2016 ¹²
Benvenuto Cellini (1500–1571), sculptor and writer, Italy	29	ReA (PD)	· Syphilis · Malaria	Painful ocular inflammation Skin rash Chronic arthritis Back pain History of suspected sexual contact	· Lignum vitae · Mercury (reported to cure syphilis)	Yes I	N/A	Anderson, 1989 ¹³
Albrecht Dürer (1471–1528), painter, Germany	20	RA (PD)	No PDD	Joint swelling and deformity	N/A	Yes	Possibly Discovered own individuality and introduced his portrabiography into Germa Polyarthritis did not necessarily interfere whis artistic activities	an arts
Jean-Baptiste- Camille Corot (1796–1875), painter, France	70	Gout (PD)	No PDD	Podagra; arthritis of right foot or both feet Walking was affected	N/A	Yes	· Possibly · Portrayed specific anatomic abnormaliti his work (e.g., the han of the "Gypsy Girl") representing gout/arti	d
Niki de Saint Phalle (1930–2002), painter and sculptor, France	50	RA (CD)	No PDD	Erosive RA Positive RF Also suffered from various other diseases: depression, chronic lung disease attributed to polystyrene exposure, selective IgA deficiency	· Antimalarials · Prednisolone · Methotrexate · Synoviorthesis wit yttrium	Yes :h	· Yes · Reflected anger and violence after disease	Zeidler, 2013 ¹⁶
Frida Kahlo (1907–1954), painter, Mexico	18	Posttraumatic fibromyalgia (PD)	· Spina bifida · Postpolio syndrome	Chronic generalized pain and profound fatigue Poliomyelitis Injuries from vehicle accident with 30 orthopedic surgeries; back pain started after accident at 18 yrs	Painkillers Orthopedic devices Metal and plaster corset Surgeries	Yes	Yes Work is completely influenced by her diseases Anguish and pain are common themes Work reflected both anger and violence as a product of her sul	
Henri de Toulouse-Lautrec (1864–1901), painter, France	Birth	Pycnodysostosis (PD)	· Osteogenesis imperfecta · Achondroplasia · Polyepiphyseal dysplasia	Shortness of stature Bone brittleness (fractures in both legs from minor trauma during his childhood) Shortness of hands Craniofacial deformity with absence of knitting of the fontanel and autosomal recessive transmission	N/A	Yes	· Yes · Physical appearance limitations deeply influenced his artworl · Used art as a way to fully express his repressed energy	Appelboom, 1989 ¹⁸

Artist (Birth–Death), Specialization, Country of Birth	Age at Disease Onset, yrs	Confirmed (CD) or Presumptive Diagnosis (PD)	Potential Differential Diagnosis (PDD)	Clinical Features	Treatment	Continued to Work After the Disease	Disease Changed the Artist's Style	Reference, First Author, Yr
Rotterdam (1466–1536), philosopher and scholar, the Netherlands	32	Pustulotic arthro-osteitis syndrome (PD)	· Gout · Syphilitic arthritis · Enteric rheumatism	· Periods of malaise, of gastrointestinal disturbances, fever with systemic manifestations, spine problems suggestive of spondylitis, pustulotic skin problems suggestive of vasculitis, joint problems typical of arthritis, and chronic osteitis · Retroperitoneal fibrosis (secondary to longstanding inflammatory process)	N/A	Yes	N/A	Dequeker, 1991 ¹⁹
Francisco Jose de Goya y Lucientes (1746–1828), painter, Spain	47	Susac syndrome	· Syphilis · Malaria · Measles · Vogt-Koyanagi-Harada disease · Vasculitis · Plumbism · Cinchonism · Cerebral arteriosclerosis · Mumps encephalitis · Cogan syndrome	"Noises in his head," and then total deafness Impairment of vision, difficulty maintaining his balance Eventually recovered his faculties, including his eyesight	Electric therapy attempt Sign language	Yes	· Yes · Turned from the pastoral to the horrif · Most importantly, hegan to draw everyous scenes, from life or from memory · One effect of Goya deafness on his art was to stimulate him to experiment with new themes and subjects in noncommissioned.	ne Hay 'S

Names in bold are those in which the artist is best known. ACTH: adrenocorticotropic hormone; anti-CCP: anticyclic citrullinated peptide antibodies; AS: ankylosing spondylitis; ESR: erythrocyte sedimentation rate; N/A: not available; RA: rheumatoid arthritis; ReA: reactive arthritis; RF: rheumatoid factor; SSc: systemic sclerosis.

- Suter H. Case report on the illness of Paul Klee (1879-1940). Case Rep Dermatol 2014;6:108-13.
- da Mota LM. On mosaics and consensus: Gaudí, Brazil and rheumatoid arthritis. Rev Bras Reumatol 2012;52:133-4.
- Pou MA, Díaz-Torné C, Azevedo VF. [Manolo Hugué: from sculpture to painting due to arthritis.] [Article in Spanish] Reumatol Clin 2011;7:135-6.
- 8. Azevedo VF. [Could Aleijadinho have suffered from scleroderma.] [Article in Italian] Reumatismo 2008;60:230-3.
- Zeidler H. [Rheumatism and art: Max Slevogt. The bon viveur and artist plagued by gout.] [Article in German] Z Rheumatol 2020;79:491-6.
- Pinals RS, Schlesinger N. Did Michelangelo have gout? J Clin Rheumatol 2015;21:364-7.
- Ventura L. [Portrait of the artist as a sick man. Rheumatological pathography of James Joyce (1882-1941).] [Article in Italian] Reumatismo 2008;60:150-8.
- 12. Cates M, Harrington C, Taylor A, Hutchinson D. "Quiet reflections" by an artist with seropositive rheumatoid arthritis, highlighting the potential role of cadmium inhalation in the workplace as a trigger for rheumatoid arthritis. J Clin Rheumatol 2016;22:276.
- Anderson B. Did Benvenuto Cellini (1500-1571) have Reiter's disease? Sex Transm Dis 1989;16:47-8.

- Weisz GM. Arthritis in the Durer family. Wien Klin Wochenschr 2007;119:553-6.
- Panush RB, Caldwell JR, Panush RS. Corot's 'gout' and a 'gipsy' girl. JAMA 1990;264:1136-8.
- 16. Zeidler H. Niki de Saint Phalle's Lifelong Dialogue Between Art and Diseases: Psychological trauma of sexual abuse, transient selective IgA deficiency, occupational exposure to toxic plastic material, chronic lung disease, rheumatoid arthritis. Joint Bone Spine 2013;80:332-7.
- 17. Courtney CA, O'Hearn MA, Franck CC. Frida Kahlo: portrait of chronic pain. Phys Ther 2017;97:90-6.
- 18. Appelboom T. The past: a gallery of arthritics. Clin Rheumatol 1989;8:442-52.
- Dequeker J. Art, history, and rheumatism: the case of Erasmus of Rotterdam 1466-1536 suffering from pustulotic arthro-osteitis. Ann Rheum Dis 1991;50:517-21.
- Hertzano R, Tomlinson JA, Mackowiak PA. Goya's lost hearing: a twenty-first century perspective on its cause, effects and possible treatment. Am J Med Sci 2019;357:275-9.
- 21. Martínez-Lavín, M; Amigo M-C; Coindreau, J; Canoso, J. Fibromyalgia in Frida Kahlo's life and art. Arthritis Rheum 2000;43:708-9.
- Chatzidionysiou K. Rheumatic disease and artistic creativity. Mediterr J Rheumatol 2019;30:103-9.

Letter 5