




Evaluation of a Patient-reported Frailty Tool in Women With Systemic Lupus Erythematosus

Sarah B. Lieber¹ , Musarrat Nahid², Stephen Paget¹, Jessica R. Berman¹, Medha Barbhaiya¹, Lisa R. Sammaritano¹, Kyriakos Kirou¹, John A. Carrino¹, Mangala Rajan² , Dina Sheira³, and Lisa A. Mandl¹ 

ABSTRACT. Objective. Frailty is associated with mortality in systemic lupus erythematosus (SLE), but how best to measure frailty is unclear. We aimed to compare 2 frailty metrics, the self-reported Fatigue, Resistance, Ambulation, Illnesses, and Loss of weight (FRAIL) scale (FS) and the Fried phenotype (FP), in SLE to evaluate differences between frail and nonfrail women and whether frailty is associated with self-reported disability.

Methods. Adult women aged < 70 years with validated SLE and mild/moderate disease enrolled in this cross-sectional study between August 2018 and October 2019. Correlation and agreement between the FS and the FP were determined. Differences in sociodemographic and disease characteristics, patient-reported outcome measures (PROMs), and biomarkers between frail and nonfrail participants were evaluated, as well as the association of frailty with Valued Life Activities disability.

Results. Of 67 participants, 27% and 18% were frail according to the FS and the FP, respectively. Correlation ($r = 0.51$; $P < 0.0001$) and agreement ($\kappa = 0.46$; $P = 0.0004$) between the FS and the FP were significant. Frail women had greater disease damage, high-sensitivity C-reactive protein, and interleukin 6, and worse PROMs according to both frailty definitions. Both frailty measures were associated with self-reported disability after adjustment for age, comorbidity, and disease activity and damage; this relationship was attenuated for the FP.

Conclusion. Frailty prevalence was high in this cohort of women with SLE using both frailty definitions, suggesting that frailty may be accelerated in women with SLE, particularly when based exclusively on self-report. Frailty remained associated with self-reported disability in adjusted analyses. The FS may be an informative point-of-care tool to identify frail women with SLE.

Key Indexing Terms: frailty, patient-reported outcome measures, systemic lupus erythematosus

This work was supported by a Scientist Development Award from the Rheumatology Research Foundation and a Michael D. Lockshin Fellowship from the Barbara Volcker Center for Women and Rheumatic Diseases at Hospital for Special Surgery. Research reported in this publication was supported by the National Center For Advancing Translational Sciences of the National Institutes of Health (NIH) under Award Number UL1TR002384. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

¹S.B. Lieber, MD, MS, Assistant Professor, S. Paget, MD, Professor, J.R. Berman, MD, Associate Professor, M. Barbhaiya, MD, MPH, Assistant Professor, L.R. Sammaritano, MD, Professor, K. Kirou, MD, Associate Professor, J.A. Carrino, MD, MPH, Professor, L.A. Mandl, MD, MPH, Assistant Research Professor, Hospital for Special Surgery, and Weill Cornell Medicine, New York, New York; ²M. Nahid, MSc, M. Rajan, MBA, Research Associate, Weill Cornell Medicine, New York, New York; ³D. Sheira, BA, Stanford School of Medicine, Palo Alto, California, USA.

JAC declares the following conflicts of interest: Covera Health, Image Analysis Group, Image Biopsy Lab, Pfizer, Simplify Medical, and the Arthritis & Rheumatology and Osteoarthritis Imaging journals. LAM declares the following conflicts of interest: Regeneron Pharmaceuticals. All other authors declare no conflicts of interest relevant to this article.

Address correspondence to Dr. S.B. Lieber, Hospital for Special Surgery, 535 East 70th Street, New York, NY 10021, USA. Email: liebers@hss.edu.

Accepted for publication August 17, 2021.

Frailty is a syndrome reflecting decreased homeostatic reserve and has been shown to be an independent risk factor for increased morbidity and mortality.¹ Frailty is commonly defined according to the Fried phenotype (FP), encompassing 5 domains: unintended weight loss, weakness, fatigue, slowness, and low activity.² Frailty is present when at least 3 criteria are met.

Frailty has been studied in rheumatoid arthritis,³⁻¹² and some data are available in systemic lupus erythematosus (SLE).^{13,14,15,16} Frailty has been associated with increased mortality in SLE, using the FP and an SLE-specific cumulative deficit definition of frailty, the Systemic Lupus International Collaborating Clinics Frailty Index (SLICC-FI).^{13,14,15,16} This association remained after adjusting for potential confounding factors such as age and physician-reported SLE-associated damage.¹³ Frailty measured by the FP has been associated with self-reported disability, after adjusting for age and self-reported SLE disease activity and damage.¹⁶ These observations suggest that frailty is an independent risk factor for important health outcomes in SLE. However, it may be challenging to obtain the FP in clinical practice, as it requires measurement of grip strength and walking speed as well as the completion of surveys on physical activity and depression.

The Fatigue, Resistance, Ambulation, Illnesses, and Loss of weight (FRAIL) scale (FS) is a simple 5-item self-report questionnaire.¹⁷ To our knowledge, no self-reported frailty tool has been compared with the FP in SLE.

The aims of this study were to evaluate the correlation of the FS with the FP in women with SLE and to compare differences between frail and nonfrail women, as measured with each frailty instrument. We hypothesized that the FS in addition to the FP would be associated with patient-reported disability in women with SLE. We also explored whether either definition of frailty was associated with serum biomarkers that reflect increased systemic inflammation.

METHODS

Participants and study visit. Women between the ages of 18 and 70 years meeting American College of Rheumatology (ACR) 1997 SLE criteria who were seen at the Hospital for Special Surgery (HSS) at least twice in the 12-month period preceding enrollment and who could complete surveys in English were eligible.¹⁸ Enrollment occurred between August 2018 and October 2019. Women with severe SLE activity, defined by the Safety of Estrogens in Lupus Erythematosus National Assessment–Systemic Lupus Erythematosus Disease Activity Index (SELENA-SLEDAI), at the visit prior to enrollment were excluded.¹⁸ Criteria for severe disease activity included new or worse central nervous system SLE, vasculitis, nephritis, myositis, anemia, or thrombocytopenia; use of prednisone or prednisone equivalent > 0.5 mg/kg/day; or start of a new medication for SLE apart from hydroxychloroquine. This decision was made to facilitate direct comparisons with a previous study of frailty in women with SLE with similar enrollment criteria and to avoid potential confounding of frailty by disease severity in this relatively small study.¹⁶ Women on dialysis who were pregnant, who had active malignancy (apart from nonmelanomatous skin cancer), who had overlapping autoimmune inflammatory disease (apart from Sjögren syndrome or antiphospholipid syndrome), or who had recent surgery or injury were excluded. The decision to exclude active malignancy, despite presence of malignancy among the comorbid conditions included in the FS, was made to avoid potential confounding of FP by malignancy, given the known association of frailty with malignancy.¹⁹

Patients with SLE were identified based on at least 1 International Classification of Diseases, 10th revision code (M32) for SLE. SLE diagnosis was confirmed by medical record review. Participant consent was obtained by an institutional review board (IRB)-approved study team member. Grip strength, 4-meter walk test, and blood draws were performed during a single study visit. SLE disease activity and damage were measured by the treating rheumatologist during the most recent clinical visit.^{20,21} Questionnaires were administered via REDCap (Research Electronic Data Capture), a secure Web-based survey tool,²² at the time of the study visit. Participants were permitted to complete questionnaires on their own device during or following the study visit. Approval to conduct this study was obtained from the HSS IRB (#2017-1061).

Frailty. Frailty, according to Fried's definition and largely consistent with a previous study of frailty in women with SLE,^{2,16} was outlined as follows: (1) unintended weight loss: BMI < 18.5 kg/m² or self-reported unintended weight loss of ≥ 10 pounds over the past year; (2) weakness: hand grip normalized for BMI (Jamar hydraulic hand dynamometer, JLW Instruments); (3) fatigue: affirmative response to "Everything I did was an effort" or "I could not get 'going'" on the Center for Epidemiologic Studies Depression (CES-D) scale²³; (4) slowness: time to walk 4 meters (normalized for height); and (5) low activity: < 600 MET-minutes/week according to the International Physical Activity Questionnaire (Supplementary Table 1, available with the online version of this article).²⁴ Participants are considered frail if they meet ≥ 3 criteria.

The self-report FS, an alternate method of measuring frailty proposed on the basis of systematic review of the literature and expert opinion²⁵ and validated in middle-aged adults,¹⁷ was administered during the same study visit (Supplementary Table 1, available with the online version of this article).¹⁷ This 5-item questionnaire gives participants 1 point for each question answered in the affirmative. Participants are asked about fatigue, resistance (difficulty climbing steps), ambulation, illnesses (comorbidity), and loss of weight. Participants who score ≥ 3 points are defined as frail.

Sociodemographic and clinical characteristics. Date of birth and Charlson Comorbidity Index²⁶ were extracted from the medical record. Race/ethnicity, education level, cigarette smoking (never/past/current), corticosteroid use (current dose), immunosuppressive medication use (within the past year), and SLE duration were self-reported during the study visit. Participants were asked if they had been diagnosed with fibromyalgia (FM), as secondary FM is common in SLE.²⁷

Disease activity and damage. SLE disease activity and damage were scored using the SELENA-SLEDAI²⁰ and the SLICC/ACR Damage Index for SLE (SDI).²¹ The SELENA-SLEDAI range is 0–105, with higher scores indicating worse disease activity. The SDI range is 0–46, with higher scores indicating worse damage. Both disease activity and damage were reported as continuous variables.

Patient-reported outcomes. The following Patient Reported Outcomes Measurement Information System (PROMIS) computerized adaptive tests (CATs) were assessed: physical function (v2.0), mobility (v2.0), pain behavior (v1.0), pain interference (v1.1), fatigue (v1.0), anxiety (v1.0), and depression (v1.0).²⁸ These generic instruments have been validated in women with SLE.²⁹ PROMIS CAT domains are scored using *t*-scores, where high values represent more of the domain being measured. A score of 50 represents the population mean and a difference of 5 (a half SD) is generally accepted as being clinically meaningful.

The Lupus Quality of Life (LupusQoL) is a validated SLE-specific patient-reported outcome measure (PROM) that includes 34 questions across 8 domains.³⁰ Score range is 0–100, with higher scores indicating better health-related QOL (HRQOL). Disability was assessed using the Valued Life Activities (VLA) instrument, which includes 21 items, each rated on a 0–3 scale, and has been validated in SLE.³¹ Score range is 0–3, with higher scores indicating greater disability. Depression was additionally assessed using the CES-D scale, which includes 20 items, each rated on a 0–3 scale.²³ Score range is 0–60, with higher scores indicating greater depressive symptoms. PROMs were reported as continuous variables.

Inflammatory and metabolic biomarkers. Erythrocyte sedimentation rate (ESR), high-sensitivity C-reactive protein (hsCRP), interleukin 6 (IL-6), insulin-like growth factor 1 (IGF-1), and soluble tumor necrosis factor receptor-1 (sTNFR1) were measured at a central laboratory. IL-6 and sTNFR1 were used to calculate a previously validated inflammatory index, which has been associated with increased mortality in community-dwelling elderly³²; higher scores have been associated with higher mortality. IGF-1³³ has been proposed as an additional biomarker for frailty.

Analysis. A cross-sectional analysis of baseline measurements of the prospective cohort was performed on all observed data. Participants who completed both the FP and the FS were included in this analysis. Frequencies or medians with IQR were determined for demographic characteristics and compared between frail and nonfrail subjects using Fisher exact or Wilcoxon rank-sum tests. PROMs and laboratory biomarkers were compared between frail and nonfrail participants using Wilcoxon rank-sum tests. Logistic regression was used to assess whether frailty was associated with disability, defined as the highest quartile of VLA disability, including after adjustment for multiple potential confounding factors, which were selected a priori and informed by prior data.¹⁶ Correlation between the FS and the FP was determined using Spearman correlation and agreement using a κ statistic. Sensitivity and specificity of the FS as compared to the FP also were determined.

RESULTS

Sample characteristics. Of 417 potential participants with validated SLE, 172 were eligible at the time of enrollment, with the remainder ineligible due to transient factors such as severe disease activity ($n = 111$) or fixed factors such as overlap conditions ($n = 134$). Seventy-two women with SLE were enrolled during the predefined study period of August 2018 to October 2019. Median (IQR) age was 44.5 (31.0–58.0) years, and the median duration of SLE was 13.0 (7.0–23.0) years. Participants were 31.0% White, 33.8% Black or African American, 7.0% Asian, and 28.6% Hispanic or Latino. Median (IQR) SELENA-SLEDAI score was 3.5 (0–4.0), SDI score was 0 (0–2.0), and duration between study visits and assessment of disease activity and damage was 15 (0–36.5) days. There were no statistically significant differences in sociodemographic features or disease characteristics between those who did ($n = 67$) and did not ($n = 5$) complete the FS, potentially due to survey fatigue.

Frailty classifications. Of the 67 participants who completed both the FS and the FP, significantly more women were classified as frail according to the FS than the FP (26.9% vs 17.9%, $P < 0.01$; Table 1). Nine participants were classified as frail according to both the FS and the FP. There was a statistically significant correlation between the FS and the FP ($r = 0.51$; $P < 0.0001$), as well as statistically significant agreement ($\kappa = 0.46$; $P = 0.0004$). Using the FP as the standard for identifying frailty, the FS had a sensitivity of 0.75 and specificity of 0.84 (data not shown).

Using the FS, fatigue (52.2%), ambulation (35.8%), and resistance (34.3%) were the most commonly self-reported components of frailty (Table 1). Using the FP to define frailty, fatigue (46.3%) and weakness (41.8%) were the most common components. Significantly more participants self-reported fatigue ($P = 0.03$) and ambulation difficulty ($P < 0.01$) based on the FS, compared to the fatigue and slow gait criteria of the FP.

Frail women according to both definitions had significantly greater SLE disease damage than nonfrail women (Table 2). When classified by the FS, frail women were significantly older ($P = 0.04$) with less educational attainment ($P < 0.01$). Compared with nonfrail women, frail women classified by the FP had significantly greater comorbidity burden ($P = 0.01$). There was no difference in SLE disease activity, current prednisone dose, or immunosuppressive medication use between frail and nonfrail women using either frailty definition.

Patient-reported outcomes.

· Generic instruments: Frail women, whether classified using the FS or the FP, reported clinically meaningful and statistically worse PROMIS mobility, physical function, pain interference, and fatigue as compared to nonfrail participants (Table 3). Pain behavior was statistically worse, but of borderline clinical significance according to the FP. Clinically meaningful and statistically worse PROMIS depression ($P = 0.02$) was observed in frail women according to the FS. Statistically significantly greater depressive symptoms based on the CES-D scale, as well as increased prevalence of disability, were found in frail women according to both definitions.

· SLE-specific instruments: Frail women reported statistically significantly worse physical health, pain, planning, and fatigue according to the LupusQoL, regardless of which frailty definition was used (Table 3). Frail women according to the FP also described less satisfactory intimate relationships ($P < 0.01$).

Associations with disability. Odds of disability were significantly higher in frail women with SLE compared with nonfrail women using both frailty definitions (Table 4). Using the FS definition resulted in a nearly 3-fold higher OR of disability compared with the FP (OR 17.2, 95% CI 4.5–66.2, $P < 0.01$ vs OR 6.2, 95% CI 1.6–23.5, $P < 0.01$). This association remained significant after adjustment for age, comorbidity burden, and disease activity (FP: $P = 0.03$; FS: $P < 0.01$). After adjusting further for disease damage, the association remained statistically significant when using the FS (OR 17.8, 95% CI 3.4–94.2, $P < 0.01$), but the association was attenuated when using the FP (OR 4.2, 95% CI 0.8–21.7, $P = 0.09$; Table 4).

Inflammatory biomarkers. Frail women according to either definition had significantly greater hsCRP (FP: $P < 0.01$; FS: $P = 0.02$) and IL-6 ($P < 0.01$; Table 5). Frail women also had a higher inflammatory index score (FP: $P < 0.01$; FS: $P = 0.02$). Compared with nonfrail women, frail women classified by the FP also had a higher ESR ($P < 0.01$).

DISCUSSION

Frailty was present in 27% and 18% of this cohort of women with SLE according to the FS and the FP, respectively. These estimates exceed the 10.7% prevalence of frailty seen in community-dwelling elderly.³⁴ This is despite the low median

Table 1. Prevalence of frailty categories and individual components by frailty definition.

Component	FRAIL Scale, n = 67		Fried Phenotype, n = 67		P
	n (%)	n (%)	Component	n (%)	
Weight loss	14 (20.9)		Weight loss	10 (14.9)	0.20
Fatigue	35 (52.2)		Fatigue	31 (46.3)	0.03
Ambulation	24 (35.8)		Slow gait	12 (17.9)	< 0.01
Resistance	23 (34.3)		Weakness	28 (41.8)	0.12
NA	NA		Inactivity	16 (23.9)	NA
Illnesses	3 (4.5)		NA	NA	NA
Classification	n (%)		Classification	n (%)	NA
Frail	18 (26.9)		Frail	12 (17.9)	< 0.01

FRAIL: Fatigue, Resistance, Ambulation, Illnesses, and Loss of weight; NA: not applicable.

Table 2. Characteristics of women by frailty classification.

	FRAIL Scale, n = 67			Fried Phenotype, n = 67		
	Nonfrail, n = 49	Frail, n = 18	P	Nonfrail, n = 55	Frail, n = 12	P
Age, yrs	42.0 (29.0–56.0)	55.0 (37.0–64.0)	0.04	41.0 (31.0–57.0)	57.0 (52.5–62.0)	0.05
Race, n (%)			0.70			0.27
Asian	5 (10.4)	0 (0)		5 (9.1)	0 (0)	
Black or African American	15 (31.3)	7 (38.9)		15 (27.3)	7 (63.6)	
White	15 (31.3)	6 (33.3)		19 (34.6)	2 (18.2)	
Other	11 (22.9)	5 (27.8)		14 (25.5)	2 (18.2)	
Declined to state	2 (4.2)	0 (0)		2 (3.6)	0 (0)	
Ethnicity, n (%)			0.10			0.71
Hispanic	11 (23.4)	8 (44.4)		15 (27.8)	4 (36.4)	
Non-Hispanic	36 (76.6)	10 (55.6)		39 (72.2)	7 (63.6)	
Education, n (%)			< 0.01			0.25
High school or less	2 (4.1)	8 (44.4)		6 (10.9)	4 (33.3)	
Some college	12 (24.5)	4 (22.2)		13 (23.6)	3 (25.0)	
College	22 (44.9)	3 (16.7)		22 (40.0)	3 (25.0)	
Graduate/professional school	13 (26.5)	3 (16.7)		14 (25.5)	2 (16.7)	
SELENA-SLEDAI ^a	2.0 (0.0–4.0)	4.0 (2.0–7.0)	0.11	1.0 (0.0–7.5)	4.0 (0.0–4.0)	0.59
SDI ^b	0.0 (0.0–1.0)	2.0 (1.0–4.0)	< 0.01	0.0 (0.0–2.0)	3.0 (1.0–5.0)	< 0.01
CCI	3.0 (2.0–6.0)	3.0 (1.0, 5.0)	0.35	2.0 (1.0, 3.0)	3.5 (2.5, 6.0)	0.01
Current prednisone dose, mg	5.0 (4.0–8.0)	5.0 (4.5–6.5)	0.88	5.0 (4.0–9.0)	5.0 (5.0–5.0)	0.81
Immunosuppressive medication use, n (%) ^c	32 (65.3)	15 (83.3)	0.23	36 (65.5)	11 (91.7)	0.09
BMI, kg/m ²	24.1 (22.1–29.7)	28.3 (23.2–31.2)	0.09	24.1 (22.1–30.8)	27.0 (25.7–33.9)	0.08
Current smoking, n (%)	2 (4.1)	2 (11.1)	0.29	2 (3.6)	2 (16.7)	0.14
Self-reported fibromyalgia, n (%)	6 (12.2)	6 (33.3)	0.07	8 (14.6)	4 (33.3)	0.21

Values are expressed as median (IQR) unless otherwise stated. ^a Score range of 0–105, with higher scores indicating greater disease activity. ^b Score range of 0–46, with higher scores indicating greater damage. ^c Immunosuppressive medications: azathioprine, belimumab, cyclophosphamide, cyclosporine, methotrexate, mycophenolate mofetil, mycophenolic acid, rituximab, tacrolimus, tocilizumab. Missing values: race (n = 1); ethnicity (n = 2); prednisone (n = 26). CCI: Charlson Comorbidity Index; FRAIL: Fatigue, Resistance, Ambulation, Illnesses, and Loss of weight; SELENA-SLEDAI: Safety of Estrogens in Lupus Erythematosus National Assessment–Systemic Lupus Erythematosus Disease Activity Index; SDI: Systemic Lupus International Collaborating Clinics/American College of Rheumatology Damage Index.

SELENA-SLEDAI score of 2–4, representing minimal SLE disease activity.³⁵ These are surprisingly high prevalence values for women with a median age of 44.5 years, suggesting that frailty may be accelerated in women with SLE. Our FP prevalence is comparable to the 20% prevalence of FP-calculated frailty reported in what is, to our knowledge, the only other cohort of women with SLE in which frailty according to the FP has been evaluated. These women had similar sociodemographic and disease characteristics, providing face validity for our findings.¹⁶

Although the FP is the most widely used disease-agnostic definition of frailty, it is challenging to implement at point of care due to multiple non-standard-of-care performance-based measures and PROMs. The prevalence of frailty in our cohort was higher according to the FS (27% vs 18%, $P < 0.01$), with statistically significant correlation between the 2 instruments ($r = 0.51$; $P < 0.0001$). Educational attainment differed between frail and nonfrail participants according to the FS, but not the FP, whereas comorbidity burden differed between frail and nonfrail participants according to the FP, but not the FS. These differences suggest that although similar, these 2 metrics measure slightly different constructs. Frailty according to the FP has been associated with higher comorbidity burden, similar to our findings.³⁶ In contrast with our findings, frailty according to the FP has been associated with lesser educational attainment.³⁷ Further

study is needed to understand the performance characteristics of both metrics in diverse patients with SLE.

The proportion of subjects who reported fatigue and slow gait/ambulation differed significantly between the FS and the FP. There was a higher prevalence of fatigue in the FS compared to the FP (52.2% vs 46.3%; $P = 0.03$). Although both are self-report, the Fried definition asks about fatigue over the past week, whereas the FS asks about fatigue over the past month; therefore, the FS gives participants a longer time in which to consider themselves fatigued. The FP asks about fatigue tangentially, whereas the FS asks directly if the subject felt “tired.” The FS wording may be more likely to elicit a positive response. Whether this is specific to patients with SLE or might differ by age, sex, or presence of concurrent FM is not known. Ambulation difficulty was 2 times higher when self-reported in the FS compared with the objective measurement of slow gait in the FP (35.8% vs 17.9%; $P < 0.01$). This may relate to the walking distance in each definition: the FS assesses participant comfort with walking several hundred yards, whereas the FP requires a 4-m walk test. Although self-reported walking speed has been found to be strongly associated with observed walking speed in older adults and both measures have been associated with mortality in this population,³⁸ it is possible that a subjective self-report of ambulation difficulty is not a good proxy for gait speed. A

Table 3. Patient-reported outcome measures by frailty classification.

	FRAIL Scale, n = 67			Fried Phenotype, n = 67		
	Nonfrail, n = 49	Frail, n = 18	P	Nonfrail, n = 55	Frail, n = 12	P
PROMIS domains^a						
Mobility	46.6 (42.0–49.7)	35.2 (33.0–37.7)	< 0.01	46.4 (40.2–49.7)	34.1 (31.9–38.1)	< 0.01
Physical function	44.8 (40.9–50.1)	32.7 (28.8–37.5)	< 0.01	43.9 (39.8–48.5)	32.7 (26.7–35.5)	< 0.01
Pain behavior	55.4 (48.5–59.0)	62.1 (58.5–63.6)	< 0.01	56.6 (49.7–59.8)	60.5 (57.6–63.1)	< 0.01
Pain interference	54.3 (46.6–57.3)	65.7 (58.5–68.2)	< 0.01	54.3 (46.6–60.1)	62.7 (58.4–67.6)	< 0.01
Fatigue	55.4 (48.5–62.4)	65.9 (62.4–73.9)	< 0.01	55.6 (49.1–62.7)	72.8 (64.0–73.9)	< 0.01
Depression	51.2 (44.6–55.0)	56.8 (49.9–67.6)	0.02	51.3 (44.7–57.5)	56.5 (48.1–69.5)	0.12
Anxiety	53.7 (49.9–61.3)	60.2 (52.9–65.1)	0.17	54.1 (50.6–61.5)	61.2 (48.1–69.1)	0.33
LupusQoL domains^b						
Physical health	75.0 (56.3–84.4)	39.1 (25.0–56.3)	< 0.01	71.9 (53.1–84.4)	32.8 (17.2–51.6)	< 0.01
Pain	83.3 (50.0–91.7)	45.8 (25.0–50.0)	< 0.01	75.0 (50.0–91.7)	45.8 (25.0–50.0)	< 0.01
Planning	75.0 (66.7–91.7)	50.0 (16.7–83.3)	< 0.01	75.0 (66.7–91.7)	45.8 (16.7–75.0)	< 0.01
Intimate relationships	75.0 (50.0–100.0)	12.5 (0–75.0)	0.05	75.0 (50.0–100.0)	0.0 (0.0–37.5)	< 0.01
Burden to others	75.0 (50.0–83.3)	62.5 (50.0–83.3)	0.27	75.0 (58.3–83.3)	62.5 (29.2–83.3)	0.48
Emotional health	83.3 (58.3–91.7)	56.3 (50.0–83.3)	0.06	83.3 (58.3–91.7)	52.1 (37.5–91.7)	0.20
Body image	90.0 (50.0–91.7)	61.3 (50.0–83.3)	0.19	80.0 (50.0–91.7)	59.2 (43.8–79.2)	0.12
Fatigue	56.3 (37.5–75.0)	43.8 (18.8–56.3)	0.02	56.3 (37.5–75.0)	40.6 (15.6–56.3)	0.04
VLA ^c	0.5 (0.1–0.8)	1.2 (1.0–1.7)	< 0.01	0.5 (0.2–0.9)	1.2 (1.1–1.8)	< 0.01
CES-D ^d	17.0 (15.0, 22.0)	22.5 (19.0, 28.0)	< 0.01	17.0 (15.0–22.0)	25.0 (20.5–33.5)	< 0.01

Values are expressed as median (IQR) unless otherwise stated. ^a Scored using a *t*-score metric, with 50 representing the population mean and a difference of 5 considered clinically significant. ^b Score range of 0–100, with higher scores indicating higher health-related quality of life. ^c Score range of 0–3, with higher scores indicating greater disability. ^d Score range of 0–60, with high scores indicating greater depression. Missing values: intimate relationships (n = 18); VLA (n = 1). CES-D: Center for Epidemiologic Studies Depression Scale; FRAIL: Fatigue, Resistance, Ambulation, Illnesses, and Loss of weight; LupusQoL: Lupus Quality of Life; PROM: patient-reported outcome measure; PROMIS: Patient Reported Outcomes Measurement Information System; VLA: Valued Life Activities.

Table 4. Cross-sectional association of frailty with self-report disability by frailty classification.

Model	FRAIL Scale, n = 66			Fried Phenotype, n = 66		
	Nonfrail, n = 48 ^a	Frail, n = 18 ^b	P	Nonfrail, n = 54 ^c	Frail, n = 12 ^d	P
	OR	95% CI		OR	95% CI	
Unadjusted ^e	17.2	4.5–66.2	< 0.01	6.2	1.6–23.5	< 0.01
Adjusted for age	15.1	3.8–59.6	< 0.01	4.7	1.2–18.8	0.03
Adjusted for age and CCI	16.5	4.0–68.5	< 0.01	6.6	1.4–32.2	0.02
Adjusted for age, CCI, and disease activity	20.8	4.1–105.8	< 0.01	6.0	1.2–29.6	0.03
Adjusted for age, CCI, disease activity, and disease damage	17.8	3.4–94.2	< 0.01	4.2	0.8–21.7	0.09

^a Disability: n = 5; no disability: n = 43. ^b Disability: n = 12; no disability: n = 6. ^c Disability: n = 10; no disability: n = 44. ^d Disability: n = 7; no disability: n = 5. ^e Odds of VLA score in the top quartile in frail vs nonfrail women. CCI: Charlson Comorbidity Index; FRAIL: Fatigue, Resistance, Ambulation, Illnesses, and Loss of weight; VLA: Valued Life Activities.

person's perception of ambulation difficulty may be tied more closely to real-world function, and thus morbidity and mortality, rather than how quickly a person can walk a short distance when required by a study investigator. How best to operationalize gait speed and ambulation to optimize the predictive value for frailty studies is an important area for future research.

Both definitions of frailty were associated with poor HRQOL using global and disease-specific instruments. While poor physical function and fatigue are components of frailty, and thus would be expected to be reported more frequently by frail women with SLE, pain was also significantly more common in frail women as compared to nonfrail women. This is consistent

with prior observations in women with SLE, suggesting that pain is increased in frail vs nonfrail women.¹⁶ Persistent pain is a risk factor for frailty in the general population.³⁹ Further study of the longitudinal relationship of pain with frailty in SLE is needed to better understand causal relationships. Although self-reported FM was about 3 times more common in frail vs nonfrail women, the difference was not statistically significant. FM was present in only one-third of frail women, calling into question the potential critique that frailty is simply a proxy for FM.

In unadjusted models, frailty, according to both the FS and the FP, was associated with significantly increased odds of patient-reported disability, confirming the association

Table 5. Inflammatory and metabolic biomarkers by frailty classification.

	FRAIL Scale, n = 67			Fried Phenotype, n = 67		
	Nonfrail, n = 49	Frail, n = 18	P	Nonfrail, n = 55	Frail, n = 12	P
ESR, mm	14.0 (8.0–24.0)	20.0 (13.0–32.0)	0.06	13.0 (8.0–22.0)	28.0 (20.0–52.0)	< 0.01
hsCRP, mg/L	1.4 (0.6–4.1)	4.6 (2.0–10.2)	0.02	1.6 (0.6–4.2)	8.3 (2.1–10.2)	< 0.01
IGF-1, ng/mL	137.3 (102.0–206.6)	107.5 (78.3–131.1)	0.06	128.2 (100.0–206.5)	109.4 (78.1–173.3)	0.28
IL-6, pg/mL	0.8 (0.4–1.3)	1.6 (1.2–2.2)	< 0.01	0.8 (0.4–1.3)	2.1 (1.6–3.0)	< 0.01
sTNFR1, pg/mL	1141.3 (983.2–1367.1)	1275.9 (1136.2–1524.1)	0.11	1148.5 (1012.8–1361.9)	1390.7 (1229.4–1992.5)	0.06
Inflammatory index score ^a	4.7 (4.4–4.8)	4.9 (4.7–5.2)	0.02	4.7 (4.4–4.8)	5.1 (4.8–5.4)	< 0.01

Values are expressed as median (IQR) unless otherwise stated. ^aInflammatory index score = 1/3 log(IL-6) + 2/3 log(sTNFR1). Higher inflammatory index score is worse. Missing values: ESR (n = 2); hsCRP (n = 1); IGF-1 (n = 1); IL-6 (n = 10); sTNFR1 (n = 4); inflammatory index score (n = 13). ESR: erythrocyte sedimentation rate; FRAIL: Fatigue, Resistance, Ambulation, Illnesses, and Loss of weight; hsCRP: high-sensitivity C-reactive protein; IGF-1: insulin-like growth factor-1; IL-6: interleukin 6; sTNFR1: soluble tumor necrosis factor receptor-1.

previously observed in a different cohort of women with SLE, in which frailty was defined according to the FP.¹⁶ This association was attenuated for the FP, but not the FS definition of frailty, after adjustment for multiple confounders, though CIs were wide. The stronger association between VLA disability and the FS as compared to the FP may reflect the collinearity between 2 exclusively self-report instruments. Whether the FS is also more strongly associated with objectively measured adverse health outcomes in SLE, or is a better predictor of downstream morbidity and mortality, requires further study.

We found higher hsCRP and IL-6 in frail as compared to nonfrail women with SLE, using either frailty definition. Higher ESR was observed among frail women with SLE as defined by the FP. Frail women using both definitions had higher inflammatory index scores, which have been associated with increased mortality in community-dwelling elderly, although this is not validated in SLE.³² While inflammatory biomarkers often reflect SLE disease activity, in our study disease activity did not differ between frail and nonfrail women and would be unlikely to explain differences in inflammatory markers between these groups. Larger studies are needed to explore this relationship.

There has been limited prior study of frailty in SLE.^{13,14,15,16} In a longitudinal cohort of 152 women with SLE, frailty, defined according to the FP, was present in 20% of patients with SLE despite low self-reported disease activity and damage.¹⁶ Frailty was associated with greater self-reported functional decline and higher mortality in patients with SLE, after adjustment for age, disease duration, and self-reported disease damage over a period of up to 8.8 years. In a cohort of 1683 patients with SLE, using the SLICC-FI, an SLE-related frailty index, higher frailty scores at baseline were independently associated with increased risk of mortality.¹³ These studies did not explore associations of frailty with inflammatory or metabolic biomarkers, and only evaluated a limited number of patient-reported outcomes.

Our study has some limitations. Our sample size was limited, and our results, generated through multiple statistical comparisons, should be viewed as hypothesis-generating. Our results cannot be generalized to men or patients with high SLE disease activity, and frailty may be underrepresented in our study compared to the general SLE population due to multiple

exclusion criteria. To ensure homogeneity of our sample, we excluded those whose SLE diagnosis could not be validated according to ACR criteria¹⁸; hence, our subjects are unlikely representative of all patients treated as SLE in clinical practice. It was beyond the scope of this study to enroll non-English speakers. Our study was likely underpowered to evaluate confounding or effect modification by age, SLE disease characteristics, and steroid and immunosuppressive medication use, and there may be residual confounding by measured and unmeasured factors. We excluded participants who did not complete the FS, potentially introducing selection bias, though those who did and did not complete the FS did not differ significantly in terms of socio-demographic characteristics. Although we classified participants as frail according to 2 previously validated phenotypic definitions of frailty in widespread use, there are multiple additional frailty constructs, notably including the SLICC-FI,^{13,40} and it is possible that some participants might have been misclassified in the absence of a single definitive frailty definition. Since our study was cross-sectional, a causal pathway between frailty and disability could not be established.

This study has several strengths. Our study cohort was diverse in terms of race, ethnicity, and educational attainment. All participants were validated according to ACR criteria for SLE. Our study is the first, to our knowledge, in which the FS, a self-reported frailty instrument that could be feasibly integrated into clinical and research settings, was evaluated in a cohort of women with SLE.

Frailty was present in up to a quarter of our cohort and may represent a distinct subset of physiologically vulnerable women with SLE at risk of worse outcomes, independent of disease severity, organ damage, or comorbidity. Targeting behavioral or pharmacologic interventions for frailty is a rich area for future study and may be an especially attractive complementary therapeutic approach for patients with SLE.

ONLINE SUPPLEMENT

Supplementary material accompanies the online version of this article.

REFERENCES

1. Hanlon P, Nicholl BI, Jani BD, Lee D, McQueenie R, Mair FS.

- Frailty and pre-frailty in middle-aged and older adults and its association with multimorbidity and mortality: a prospective analysis of 493 737 UK Biobank participants. *Lancet Public Health* 2018;3:e323-e32.
2. Bandeen-Roche K, Xue QL, Ferrucci L, Walston J, Guralnik JM, Chaves P, et al. Phenotype of frailty: characterization in the women's health and aging studies. *J Gerontol A Biol Sci Med Sci* 2006;61:262-6.
 3. Andrews JS, Trupin L, Yelin EH, Hough CL, Covinsky KE, Katz PP. Frailty and reduced physical function go hand in hand in adults with rheumatoid arthritis: a US observational cohort study. *Clin Rheumatol* 2017;36:1031-9.
 4. Andrews JS, Trupin L, Wysham KD, Hough CL, Yelin EH, Katz PP. The impact of frailty on changes in physical function and disease activity among adults with rheumatoid arthritis. *ACR Open Rheumatol* 2019;1:366-72.
 5. Barile-Fabris LA, Pérez-Cristobal M, Merlos-López R], Xibillé-Friedman D. [Frailty syndrome in patients with rheumatoid arthritis]. [Article in Spanish] *Rev Med Inst Mex Seguro Soc* 2016;54 Suppl 2:S210-5.
 6. Haider S, Grabovac I, Berner C, Lamprecht T, Fenzl KH, Erlacher L, et al. Frailty in seropositive rheumatoid arthritis patients of working age: a cross-sectional study. *Clin Exp Rheumatol* 2019;37:585-92.
 7. Li G, Chen M, Li X, Cesta A, Lau A, Thabane L, et al. Frailty and risk of osteoporotic fractures in patients with rheumatoid arthritis: data from the Ontario Best Practices Research Initiative. *Bone* 2019;127:129-34.
 8. Oetsma S, Boonen A, Starmans M, Peeters R, van Onna M. Validation of two frailty questionnaires in older patients with rheumatoid arthritis: a cross-sectional study. *Clin Exp Rheumatol* 2020;38:523-8.
 9. Salaffi F, Di Carlo M, Farah S, Di Donato E, Carotti M. Prevalence of frailty and its associated factors in patients with rheumatoid arthritis: a cross-sectional analysis. *Clin Rheumatol* 2019; 38:1823-30.
 10. Salaffi F, Di Carlo M, Farah S, Carotti M. The Comprehensive Rheumatologic Assessment of Frailty (CRAF): development and validation of a multidimensional frailty screening tool in patients with rheumatoid arthritis. *Clin Exp Rheumatol* 2020;38:488-99.
 11. Wysham KD, Shoback DM, Andrews JS, Katz PP. Sex differences in frailty and its association with low bone mineral density in rheumatoid arthritis. *Bone Rep* 2020;12:100284.
 12. Tada M, Yamada Y, Mandai K, Hidaka N. Correlation between frailty and disease activity in patients with rheumatoid arthritis: data from the CHIKARA study. *Geriatr Gerontol Int* 2019;19:1220-5.
 13. Legge A, Kirkland S, Rockwood K, Andreou P, Bae SC, Gordon C, et al. Evaluating the properties of a frailty index and its association with mortality risk among patients with systemic lupus erythematosus. *Arthritis Rheumatol* 2019;71:1297-307.
 14. Legge A, Kirkland S, Rockwood K, Andreou P, Bae S-C, Gordon C, et al. Construction of a frailty index as a novel health measure in systemic lupus erythematosus. *J Rheumatol* 2020;47:72-81.
 15. Legge A, Kirkland S, Rockwood K, Andreou P, Bae SC, Gordon C, et al. Prediction of damage accrual in systemic lupus erythematosus using the Systemic Lupus International Collaborating Clinics Frailty Index. *Arthritis Rheumatol* 2020;72:658-66.
 16. Katz PP, Andrews J, Yazdany J, Schmajuk G, Trupin L, Yelin E. Is frailty a relevant concept in SLE? *Lupus Sci Med* 2017;4:e000186.
 17. Morley JE, Malmstrom TK, Miller DK. A simple frailty questionnaire (FRAIL) predicts outcomes in middle aged African Americans. *J Nutr Health Aging* 2012;16:601-8.
 18. Hochberg MC. Updating the American College of Rheumatology revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum* 1997;40:1725.
 19. Handforth C, Clegg A, Young C, Simpkins S, Seymour MT, Selby PJ, et al. The prevalence and outcomes of frailty in older cancer patients: a systematic review. *Ann Oncol* 2015;26:1091-101.
 20. Petri M, Kim MY, Kalunian KC, Grossman J, Hahn BH, Sammaritano LR, et al; OC-SELENA Trial. Combined oral contraceptives in women with systemic lupus erythematosus. *N Engl J Med* 2005;353:2550-8.
 21. Gladman D, Ginzler E, Goldsmith C, Fortin P, Liang M, Urowitz M, et al. The development and initial validation of the Systemic Lupus International Collaborating Clinics/American College of Rheumatology Damage Index for systemic lupus erythematosus. *Arthritis Rheum* 1996;39:363-9.
 22. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009;42:377-81.
 23. Radloff LS. The CES-D scale: a self-report depression scale for research in the general population. *Appl Psych Meas* 1977; 1:385-401.
 24. Craig CL, Marshall AL, Sjöström M, Bauman AE, Booth ML, Ainsworth BE, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 2003;35:1381-95.
 25. Abellan van Kan G, Rolland Y, Bergman H, Morley JE, Kritchevsky SB, Vellas B. The I.A.N.A Task Force on frailty assessment of older people in clinical practice. *J Nutr Health Aging* 2008;12:29-37.
 26. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 1987;40:373-83.
 27. Staud R. Are patients with systemic lupus erythematosus at increased risk for fibromyalgia? *Curr Rheumatol Rep* 2006;8:430-5.
 28. HealthMeasures. Intro to PROMIS [Internet. Accessed September 16, 2021]. Available from: www.healthmeasures.net/explore-measurement-systems/promis/intro-to-promis
 29. Kasturi S, Szymonifka J, Burket JC, Berman JR, Kirou KA, Levine AB, et al. Validity and reliability of Patient Reported Outcomes Measurement Information System computerized adaptive tests in systemic lupus erythematosus. *J Rheumatol* 2017;44:1024-31.
 30. Jolly M, Pickard AS, Wilke C, Mikolaitis RA, Teh LS, McElhone K, et al. Lupus-specific health outcome measure for US patients: the LupusQoL-US version. *Ann Rheum Dis* 2010;69:29-33.
 31. Katz P, Morris A, Trupin L, Yazdany J, Yelin E. Disability in valued life activities among individuals with systemic lupus erythematosus. *Arthritis Rheum* 2008;59:465-73.
 32. Varadhan R, Yao W, Matteini A, Beamer BA, Xue QL, Yang H, et al. Simple biologically informed inflammatory index of two serum cytokines predicts 10 year all-cause mortality in older adults. *J Gerontol A Biol Sci Med Sci* 2014;69:165-73.
 33. Leng SX, Hung W, Cappola AR, Yu Q, Xue QL, Fried LP. White blood cell counts, insulin-like growth factor-1 levels, and frailty in community-dwelling older women. *J Gerontol A Biol Sci Med Sci* 2009;64:499-502.
 34. Collard RM, Boter H, Schoevers RA, Oude Voshaar RC. Prevalence of frailty in community-dwelling older persons: a systematic review. *J Am Geriatr Soc* 2012;60:1487-92.
 35. Zen M, Iaccarino L, Gatto M, Saccon F, Larosa M, Ghirardello A, et al. Lupus low disease activity state is associated with a decrease in damage progression in Caucasian patients with SLE, but overlaps with remission. *Ann Rheum Dis* 2018;77:104-10.
 36. Belga S, Majumdar SR, Kahlon S, Pederson J, Lau D, Padwal RS, et al. Comparing three different measures of frailty in medical

- inpatients: multicenter prospective cohort study examining 30-day risk of readmission or death. *J Hosp Med* 2016;11:556-62.
37. Hoogendijk EO, van Hout HP, Heymans MW, van der Horst HE, Frijters DH, Broese van Groenou MI, et al. Explaining the association between educational level and frailty in older adults: results from a 13-year longitudinal study in the Netherlands. *Ann Epidemiol* 2014;24:538-44.e2.
 38. Syddall HE, Westbury LD, Cooper C, Sayer AA. Self-reported walking speed: a useful marker of physical performance among community-dwelling older people? *J Am Med Dir Assoc* 2015;16:323-8.
 39. Saraiva MD, Suzuki GS, Lin SM, de Andrade DC, Jacob-Filho W, Suemoto CK. Persistent pain is a risk factor for frailty: a systematic review and meta-analysis from prospective longitudinal studies. *Ageing* 2018;47:785-93.
 40. Legge A, Kirkland S, Rockwood K, Andreou P, Bae SC, Gordon C, et al. Construction of a Frailty Index as a novel health measure in systemic lupus erythematosus. *J Rheumatol* 2020;47:72-81.