Post SARS-CoV-2 Vaccine Chilblains-like Lesions

To the Editor:

We would like to share ideas on the article, "A Case of Chilblains-like Lesions Post SARS-CoV-2 Vaccine?"¹ Meara et al reported a female case and concluded, "The chilblains-like lesions suggest that she had an antibody response from either 1 vaccine dose or a previous PCR-negative COVID-19 [coro-navirus disease 2019] infection."¹ The existence of antibodies might or might not explain occurrence of chilblains-like lesions. An exact pathomechanism is unclear. In addition to abnormal immune response, the process might be related to an increased blood viscosity after vaccination.² After COVID-19 vaccination, there might be an occurrence of a hyperviscosity problem,² which is reported as an uncommon cause of chilblains-like lesions.³

Rujittika Mungmunpuntipantip¹, PhD Viroj Wiwanitkit², MD, Honorary Professor ¹Private Academic Consultant, Bangkok, Thailand;

²DY Patil University, Pune, India.

The authors declare no conflicts of interest relevant to this article.

The Journal of

Rheumatolog

Address correspondence to Dr. R. Mungmunpuntipantip, Private Academic Consultant, 11 Bangkok 122, Bangkok 103300, Thailand. Email: rujittika@gmail.com.

REFERENCES

- 1. Meara AS, Silkoski M, Quin K, Jarjour W. A case of chilblains-like lesions post SARS-CoV-2 vaccine? J Rheumatol 2021;48:1754.
- Joob B, Wiwanitkit V. Expected viscosity after COVID-19 vaccination, hyperviscosity and previous COVID-19. Clin Appl Thromb Hemost 2021;27:10760296211020833.
- Millard LG, Rowell NR. Chilblain lupus erythematosus (Hutchinson). A clinical and laboratory study of 17 patients. Br J Dermatol 1978;98:497-506.