

Infliximab induced T lymphocyte apoptosis in Crohn's disease.

Jan van den Brande, Daan W Hommes and Maikel P Peppelenbosch

J Rheumatol 2005;74;26-30

<http://www.jrheum.org/content/74/26>

1. Sign up for TOCs and other alerts
<http://www.jrheum.org/alerts>
2. Information on Subscriptions
<http://jrheum.com/faq>
3. Information on permissions/orders of reprints
http://jrheum.com/reprints_permissions

The Journal of Rheumatology is a monthly international serial edited by Earl D. Silverman featuring research articles on clinical subjects from scientists working in rheumatology and related fields.

Infliximab Induced T Lymphocyte Apoptosis in Crohn's Disease

JAN M.H. Van den BRANDE, DAAN W. HOMMES, and MAIKEL P. PEPPELENBOSCH

ABSTRACT. Crohn's disease (CD) is a chronic inflammatory disorder of the gastrointestinal tract of unknown origin. Therapies include immune modulating agents, biological therapies, and surgery. The activity and efficacy of the anti-tumor necrosis factor (TNF) therapies infliximab and etanercept have proved to be different: infliximab is effective to induce and maintain remission in refractory CD, while etanercept is not. This brief review considers the question of whether this disparity can be explained by the different structure of the proteins, their different binding affinities, or the subsequent effects on T lymphocytes. (J Rheumatol 2005;32 Suppl 74:26-30)

Key Indexing Terms:

TUMOR NECROSIS FACTOR
ETANERCEPT

APOPTOSIS

CROHN'S DISEASE
INFLIXIMAB

EFFICACY OF TNF ANTAGONISTS IN CROHN'S DISEASE

Crohn's disease (CD) is a chronic inflammatory disorder of the gastrointestinal (GI) tract of unknown origin. Available therapies include immune modulating drugs, therapy with biological compounds, or surgery. Anti-tumor necrosis factor (TNF) therapy has become a drug of major importance in remission and endoscopic remission induction, which was not shown previously with available therapies. Two anti-TNF molecules are approved for clinical therapy of inflammatory disorders. Infliximab (Remicade), a chimeric anti-TNF- α antibody, is registered for both rheumatoid arthritis and Crohn's disease. Etanercept, a recombinant TNF receptor/IgG fusion protein, is registered for rheumatoid arthritis only.

Interestingly, the activity and efficacy profiles of infliximab and etanercept in Crohn's disease have proved to be different. Infliximab is effective to induce and maintain remission in steroid-refractory CD; it is also used to treat fistulizing disease. Targan, *et al* showed in a multicenter, randomized, double-blind, placebo-controlled trial that infliximab induced a clinical response by 61% of patients at Week 2, 65% at Week 4, and 41% at Week 12 in patients with moderate to severe, treatment-resistant CD compared to response rates in the placebo group of 17%, 17%, and 12%, respectively¹.

Rutgeerts, *et al* showed that response can be maintained with repeated administration. In a randomized,

double-blind, placebo-controlled study, 62% of patients treated with infliximab maintained clinical response compared to 37% of patients who received placebo every 8 weeks for 36 weeks².

In contrast, etanercept is not effective in the treatment of CD. Response rates following the administration of etanercept 25 mg or placebo subcutaneously twice weekly for 8 weeks in a randomized, double-blind, placebo-controlled trial were similar at all timepoints (2, 4, and 8 weeks). Clinical response at Week 4 (the primary study endpoint) was observed in 39% of patients in the etanercept group and 45% of the placebo group³.

The question arises whether these disparities can be explained by the different structure of the proteins, the different binding affinities, and subsequent effects on T lymphocytes.

T CELL APOPTOSIS IN CD

Defective T cell apoptosis appears to play a role in the development of inflammation in patients with CD. Lamina propria T cells isolated from patients with CD show a resistance to apoptosis that is based on several disturbances compared to controls. Indeed, these lamina propria T cells show increased interleukin 2 (IL-2) dependent proliferation and are relatively resistant to apoptosis induced by CD2 activation, IL-2 depletion, nitric oxide induced apoptosis, and engagement of Fas^{4,5}. This phenomenon is in part explained by an increased ratio of intracellular expression of the proteins Bax (pro-apoptotic) and Bcl-2 (anti-apoptotic)⁶.

BINDING OF TNF ANTAGONISTS TO MEMBRANE-BOUND TNF- α

Infliximab and etanercept bind differently to transmembrane TNF. Using a transfection model, Scallon, *et al* showed that infliximab has a higher binding affinity to transfected cells with the uncleavable form of transmembrane TNF than etanercept. Moreover, in their experiments infliximab was shown to bind both soluble and membrane-bound TNF, whereas etanercept bound primarily to soluble TNF⁷.

From the Department of Gastroenterology, Academic Medical Centre, Amsterdam; and Department of Cell Biology, Academic Medical Centre Groningen, Groningen, The Netherlands.

J.M.H. Van den Brande, MD; D.W. Hommes, MD, PhD, Department of Gastroenterology, Academic Medical Centre, Amsterdam; M.P. Peppelenbosch, Department of Cell Biology, Academic Medical Centre Groningen, Groningen, The Netherlands.

Address reprint requests to Dr. J.M.H. Van den Brande, Department of Gastroenterology, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands. E-mail: J.vandenbrande@amc.uva.nl




Figure 1. Infliximab (A) and etanercept (B) neutralized recombinant human TNF- α . From Van den Brande, et al. *Gastroenterology* 2003;124:1774-85; with permission.




Figure 2A and B. Only infliximab bound transmembrane TNF- α on activated lamina propria lymphocytes from patients with Crohn's disease. From Van den Brande, et al. *Gastroenterology* 2003;124:1774-85; with permission.




Figure 3A. Only infliximab induced apoptosis in peripheral lymphocytes activated by mixed lymphocyte reaction. *Gastroenterology* 2003;124:1774-85; with permission.




Figure 3B. Infliximab but not etanercept induced apoptosis in activated Crohn's lamina propria mononuclear cells in a mixed lymphocyte reaction. *Gastroenterology* 2003;124:1774-85; with permission.

INFLIXIMAB INDUCED APOPTOSIS

Apoptosis induction by infliximab has been suggested in 2 earlier reports. Luger and colleagues demonstrated significant infliximab binding to membrane-bound TNF, dose-dependent increases in apoptosis, and caspase-3 proteolytic activity in monocytes from patients with chronic active CD⁸. Caspase-3 activation and apoptosis following infliximab administration occurred rapidly and were significantly increased as early as 4 hours after treatment ($p = 0.002$ and $p = 0.003$ vs before treatment, respectively). Interestingly F(ab)₂ fragments of infliximab (which lack the Fc domain) also induced dose-dependent apoptosis. Second, ten Hove, *et al* showed a significant increase in TUNEL-positive lamina propria (CD3+) T cells in tissue taken from patients 24 hours

after infliximab infusion⁹.

WHY IS INFLIXIMAB, BUT NOT ETANERCEPT, EFFECTIVE IN CROHN'S DISEASE?

This background information raises a number of questions. Do infliximab and etanercept inhibit activation of target cells by neutralization of soluble TNF- α and subsequently prevent translocation of nuclear factor- κ B (NF- κ B) and DNA transcription of cytokines? Or do infliximab and etanercept bind to membrane TNF- α on target cells and alter function by inducing apoptosis? We performed a number of experiments in an attempt to answer some of these questions.

Do infliximab and etanercept differ in the capacity to neutralize TNF- α ?





Figure 4A and B. Induction of cleaved caspase-3 upon treatment with infliximab. *Gastroenterology* 2003;124:1774-85; with permission.

We assayed TNF- α neutralizing potential of these TNF antagonists with respect to TNF- α dependent transactivation of a NF- κ B-driven reporter construct. After stimulation with TNF- α , green fluorescent protein (GFP) is transcribed along with NF- κ B and can be detected by FACS analysis. Infliximab-neutralized TNF- α induced GFP expression in NF- κ B reporter construct transfected HeLa cells at a concentration of 1 g/ml, and etanercept neutralized TNF- α at a concentration of 0.25 g/ml (Figure 1)¹⁰. Thus, both infliximab and etanercept neutralized TNF- α effectively in this assay.

Do infliximab and etanercept differ in the capacity to bind to memTNF- α ?

To investigate the difference between infliximab and etanercept with respect to their capacity to bind to transmembrane TNF- α (memTNF- α) expressed by activated

lymphocytes, we used a binding assay with FITC-labeled secondary or tertiary antibodies for detection with FACS analysis. We tested the binding of these TNF antagonists to CD3/CD28-activated peripheral blood lymphocytes (PBL) from healthy volunteers and PMA-ionomycin-activated isolated lamina propria mononuclear cells from surgical specimens from patients with active Crohn's disease. A chimeric antibody directed against EpCAM, c17-1a, was added as a nonspecific control antibody. Histograms from FACS analysis showed no nonspecific binding in the group treated with secondary antibody only. Addition of infliximab resulted in a positive cell population that was not seen in the etanercept-treated group or in the control antibody group. In addition we double-stained lamina propria cells for CD3 and confirmed that the majority of the infliximab-positive cells were T lymphocytes (CD3-positive), in contrast to the etanercept condition (Figure 2a and b).

Does binding to memTNF have functional consequences for the T cell?

We used monocyte derived dendritic cells to activate PBL and lamina propria cells (obtained from patients with active CD) in a mixed lymphocyte reaction. These activated cells were incubated for 24 h in separate conditions with infliximab, etanercept, or the above mentioned isotype control antibody. Next, lymphocyte apoptosis was determined using a well established apoptosis assay with annexin V-7AAD double-staining. In brief, annexin V binds to externalized phosphatidyl serine, an early marker of impermeable apoptotic cells, and 7AAD stains the nucleus of permeable necrotic cells. In this assay, the annexin V-positive/7AAD-negative cells represent the apoptotic cells and the double-positive cells the necrotic cells. Addition of infliximab induced a significant increase of apoptosis of the activated lymphocytes (compared to the medium control), whereas etanercept and the isotype control were inactive in this respect (Figure 3a and b). Apparently, the binding of infliximab to memTNF- α on activated T cells correlates with the ability to induce apoptosis of these clinically relevant immune cells.

To further investigate markers of apoptosis we investigated the activation state of caspase-3. Caspase-3 is a downstream cleaving protein that can be activated by cleavage itself either through the intrinsic (mitochondrial)




Figure 5. Infiximab-induced apoptosis is not mediated via the antibody-dependent cytotoxicity reaction (ADCR). *Gastroenterology* 2003;124:1774-85; with permission.

or by the extrinsic (caspase-8) cell death pathway. Figure 4 shows a representative experiment of a Western blot of cells in a mixed lymphocyte reaction treated with infiximab, etanercept, and a chimeric control antibody at 24 and 48 h after treatment. Strong induction of cleaved caspase-3 can be seen upon treatment with infiximab. These results show that the induction of apoptosis by infiximab correlates with activation of caspase-3.

THE COMPLEMENT ISSUE

Our experiments involved immune cells in heat-inactivated fetal calf serum-supplemented medium; that is, we considered the complement system to be deactivated. Indeed, we showed that replacing fetal calf serum with complement-competent human serum did not increase infiximab effects on apoptosis, as measured with iodine blue staining (Figure 5A) and by granzyme-B ELISA (Figure 5B).

CONCLUSION

In our assays both infiximab and etanercept neutralize soluble TNF effectively. However, in activated lamina propria lymphocytes, only infiximab binds to membrane TNF, induces apoptosis, and activates caspase-3.

FUTURE PLANS

We intend to perform SPECT scans of patients treated with infiximab to visualize apoptosis. Response to infiximab, about two-thirds in steroid-refractory CD, might be correlated to the degree of apoptosis induction. Characterizing the nonresponder population could increase cost-effectiveness.

The role of apoptosis in remission induction and maintenance should be investigated in powerful therapy regimens such as combinations of immune modulators methotrexate and infiximab.

REFERENCES

- Targan SR, Hanauer SB, van Deventer SJ, et al. A short-term study of chimeric monoclonal antibody cA2 to tumor necrosis factor alpha for Crohn's disease. Crohn's Disease cA2 Study Group. *N Engl J Med* 1997;337:1029-35.
- Rutgeerts P, D'Haens G, Targan S, et al. Efficacy and safety of retreatment with anti-tumor necrosis factor antibody (infiximab) to maintain remission in Crohn's disease. *Gastroenterology* 1999;117:761-9.
- Sandborn WJ, Hanauer SB, Katz S, et al. Etanercept for active Crohn's disease: a randomized, double-blind, placebo-controlled trial. *Gastroenterology* 2001;121:1088-94.
- Boirivant M, Marini M, Di Felice G, et al. Lamina propria T cells in Crohn's disease and other gastrointestinal inflammation show defective CD2 pathway-induced apoptosis. *Gastroenterology* 1999;116:557-65.
- Itoh J, de La Motte C, Strong SA, Levine AD, Fiocchi C. Decreased Bax expression by mucosal T cells favours resistance to apoptosis in Crohn's disease. *Gut* 2001;49:35-41.
- Ina K, Itoh J, Fukushima K, et al. Resistance of Crohn's disease T cells to multiple apoptotic signals is associated with a Bcl-2/Bax mucosal imbalance. *J Immunol* 1999;163:1081-90.
- Scallon BJ, Moore MA, Trinh H, Knight DM, Ghraeyeb J. Chimeric anti-TNF-alpha monoclonal antibody cA2 binds recombinant transmembrane TNF-alpha and activates immune effector functions. *Cytokine* 1995;7:251-9.
- Lugering A, Schmidt M, Lugering N, Pauels HG, Domschke W, Kucharzik T. Infiximab induces apoptosis in monocytes from patients with chronic active Crohn's disease by using a caspase-dependent pathway. *Gastroenterology* 2001;121:1145-57.
- ten Hove T, van Montfrans C, Peppelenbosch MP, van Deventer SJ. Infiximab treatment induces apoptosis of lamina propria T lymphocytes in Crohn's disease. *Gut* 2002;50:206-11.
- Van den Brande JM, Braat H, van den Brink GR, et al. Infiximab but not etanercept induces apoptosis in lamina propria T-lymphocytes from patients with Crohn's disease. *Gastroenterology* 2003;124:1774-85.