Online supplement to Cardiovascular Disease Prevention in Rheumatoid Arthritis: Compliance with Diabetes Screening Guidelines, *The Journal of Rheumatology*, doi:10.3899/jrheum.170973

Supplementary Table 1. Description of plasma glucose testing.

	RA cohort (n = 22,624)	General Pop (n = 22,579)
Number of tests per patient year, [median (Q1, Q3)]	0.49 (0.25, 0.78)*	0.46 (0.22, 0.73)*
Time from index date to first PG test, days [median (25 th , 75 th)]	603 (251, 1148)**	597 (270, 1157)**
Time between PG tests in those with ≥ 2 tests, days [median (25 th , 75 th)]	307 (109, 538)*	366 (178,585)*

Abbreviations: SD: Standard deviation; 25Q: first quartile; 75Q: third quartile *p value comparing RA vs. general population <0.0001; **p value = 0.0036

Online supplement to Cardiovascular Disease Prevention in Rheumatoid Arthritis: Compliance with Diabetes Screening Guidelines, *The Journal of Rheumatology*, doi:10.3899/jrheum.170973

Supplementary Table 2. Compliance with PG test for RA individuals stratified by categorical variables.

Predictive Variable	Categories	Compliance*
Sex	Male	70.5%
	Female	71.9%
Age**	40-50	62.4%
	51-60	70.4%
	61-70	75.3%
	71-80	75.3%
	81-90	72.3%
	90+	57%
Urban vs Rural	Rural	71.5%
	Urban	71.3%
Health Authority	Interior	70.1%
	Fraser	73.4%
	Coastal	70.1%
	Island	72.5%
	Northern	69.0%
SES Index (Quartiles)	Lowest 25%	73.6%
	25%-50%	70.6%
	50%-75%	72.6%
	75%+	68.7%
SES Index, highest quartile vs others	Lowest 75%	72.3%
	Upper 25%	68%
Romano comorbidity score, ≥ 1 vs 0	0	71.6%
	1 and above	71.4%
Having a hospitalization in the prior year‡	No	70.9%

Online supplement to Cardiovascular Disease Prevention in Rheumatoid Arthritis: Compliance with Diabetes Screening Guidelines, *The Journal of Rheumatology*, doi:10.3899/jrheum.170973

Predictive Variable	Categories	Compliance*
	Yes	73.4%
Seeing a rheumatologist in the prior 5 years ^{††}	No	73.6%
	Yes	64%
Rate of physician visits in the prior year ‡‡	11 or less visits	68.9%
	More than 11 visits	74.3%
Timing of periods, starting ≥ 2001 vs. $<2001^{\dagger}$	Before 2001	60.6%
	After2001	77.2%
GC use within eligibility period	Yes	75.7%
	No	70.5%

Abbreviations: SES: Socioeconomic status; GC: Glucocorticosteroids

^{*}Compliance measured with periods as the unit of analysis, i.e., as the number of periods with ≥1 glucose test divided by the number of eligible periods in RA cohort, converted to a percentage.

^{**}Age: Stratified into groups of 10 years, measured at the start of each new period †Compliance per period was stratified according to whether the date at the beginning of the

Compliance per period was stratified according to whether the date at the beginning of the period was before or after 01 Jan 2001.

^{††}measured in the 5 years prior to the start of period

[‡]measured in the year prior to start of period

^{‡‡} rate of physician visits dichotomized at the median of 11 visits per year per individual